

Community Plan of Mississippi

Quick reference guide

Opioid overutilization prevention and opioid use disorder treatment programs for UnitedHealthcare Community Plan of Mississippi

In response to the U.S. opioid epidemic, we've developed programs to help our members receive the care and treatment they need in safe and effective ways. We've based our measures on the Centers for Disease Control and Prevention's (CDC) opioid treatment guidelines to help prevent overuse of short-acting and long-acting opioid medications. Please use this quick reference guide to learn more about what we offer.

Concurrent Drug Utilization Review (cDUR) programs

The cDUR program uses the pharmacy claims processing system to screen all prescriptions at the point-of-service and checks for possible inappropriate drug prescribing and utilization, as well as potentially dangerous medical implications or drug interactions. The program includes communication to the dispensing pharmacy at point-of-service through claims edits and messaging to the dispensing pharmacy at point-of-service. The pharmacist will need to address the clinical situation at the point-of-sale before entering appropriate National Council for Prescription Drug Programs (NCPDP) codes to receive an approved claim, unless otherwise stated below.

THERDOSE Acetaminophen	<ul style="list-style-type: none">• Combination opioids plus acetaminophen (APAP) limit• Prevents doses of APAP greater than 4 grams per day
Drug-Drug Interaction – Opioids and Medication-Assisted Treatment (MAT)	<ul style="list-style-type: none">• Point-of-sale alert for concurrent use of opioids and MAT drugs
Drug-Drug Interaction – Opioids and Carisoprodol	<ul style="list-style-type: none">• Point-of-sale alert for concurrent use of opioids and carisoprodol
Drug-Drug Interaction – Opioids and Sedative Hypnotics	<ul style="list-style-type: none">• Point-of-sale alert for concurrent use of opioids and sedative hypnotics
Duplicate Therapy – Short-Acting Opioids (SAOs)	<ul style="list-style-type: none">• Alerts to concurrent use of multiple SAOs
Duplicate Therapy – Long-Acting Opioids (LAOs)	<ul style="list-style-type: none">• Alerts to concurrent use of multiple LAOs
Drug-Inferred Health State – Opioids and Prenatal Vitamins and Medications Used in Pregnancy	<ul style="list-style-type: none">• Enhanced point-of-sale alert for concurrent use of opioids and prenatal vitamins and for concurrent use of opioids and medications used in pregnancy (e.g., doxylamine/pyridoxine)• This custom message does not require the pharmacist to enter appropriate NCPDP codes to receive an approved claim

Retrospective Drug Utilization Review (rDUR) programs

The rDUR program analyzes claims on a daily basis and sends communications to prescribers.

Abused Medications DUR Program	<ul style="list-style-type: none"> • Daily identification of members who are getting multiple opioid prescriptions from multiple prescribers and/or filling at multiple pharmacies • Also identifies members with chronic early refill attempts, overlapping LAOs, overlapping SAOs, high daily doses of opioids, a diagnosis of opioid overdose with an opioid fill, overlapping opioid and MAT medications, overlapping opioid, muscle relaxant and benzodiazepine and overlapping opioid and opioid potentiator • Patient-specific information sent to all prescribers with medication fill history for the last 4 months
Pharmacy Lock-In Program	<ul style="list-style-type: none"> • Pharmacy lock-in programs vary by state; however, all include filling of opioids by multiple prescribers at multiple pharmacies as an inclusion criterion for the program • Members chosen for the program will be locked into 1 pharmacy for all of their medications for a period of 1 year. Lock-in periods vary by state.

Utilization Management (UM) programs

UM programs promote appropriate use, help reduce costs and, ultimately, help improve the health status of members.

New to Therapy (NTT) use of LAO	<ul style="list-style-type: none"> • Prior authorization required for opioid-naïve members (no opioid claims in the last 90 days) • Step through short-acting opioid (non-cancer or non-sickle cell pain); step through preferred LAOs
Cumulative 90 Morphine Milligram Equivalents (MME) Limit	<ul style="list-style-type: none"> • Point-of-sale dosage limit for all opioid products up to 90 MME • Prevents cumulative opioid doses above the preset threshold from processing • Prior authorization required for doses above the preset threshold
NTT SAO Edit	<ul style="list-style-type: none"> • For opioid-naïve members (no opioid claims in the last 90 days), point-of-sale limits include a maximum of a 7-day supply for initial fill and a maximum of two 7-day supplies in a rolling 30-day period • Prior authorization required to exceed these quantities for opioid-naïve members
SAO Duration Limits	<ul style="list-style-type: none"> • Point-of-sale limits include a maximum of 62 units of a SAO in a rolling 31-day period • Prior authorization required to exceed these limits
Cough and Cold Products Containing Opioid Components	<ul style="list-style-type: none"> • Quantity per fill of 120 mL (units) is applied, as well as a 30-day maximum quantity of 360 mL (units) • Prior authorization is required for those under age 18
Drug-Drug Interaction – Opioids and Benzodiazepines	<ul style="list-style-type: none"> • Prior authorization required for concurrent use of opioids and benzodiazepines • If a member is currently taking chronic opioids, the member can obtain a maximum 3-day supply of a benzodiazepine without requiring a prior authorization • If a member is currently taking chronic benzodiazepines, the member can obtain a maximum 3-day supply of an opioid without

Utilization Management (UM) programs

UM programs promote appropriate use, help reduce costs and, ultimately, help improve the health status of members.

requiring a prior authorization

Transmucosal Fentanyl Product Prior Authorization

Prior authorization requires:

- Documentation of pain due to cancer and patient is already receiving opioids

Overdose Prevention (Naloxone)

- No prior authorization is required for preferred naloxone products (generic naloxone injection, Narcan® Nasal Spray)

Evidence-Based Prescribing programs

Focuses on outreach to prescribers identified as outliers.

Fraud/Waste/Abuse Evaluation

- Retrospective controlled substance claims analysis
- Identifies outlier opioid prescribers

Miscellaneous

Substance Use Disorder Help Line

- 24/7 Help Line: Call **855-780-5955**. For members or caregivers, staffed by licensed behavioral health providers.
- Reference: liveandworkwell.com

Miscellaneous – Drug Enforcement Agency (DEA) License Edit

- Verifies DEA is active and matches scheduled medication in the claim

Miscellaneous – Refill-Too-Soon Threshold

- Increases the refill-too-soon threshold to 90% on opioids and other controlled substances CII-V

Abbreviations

APAP	Acetaminophen	MME	Morphine Milligram Equivalent
CDC	Centers for Disease Control and Prevention	PA	Prior Authorization
cDUR	Concurrent Drug Utilization Review	rDUR	Retrospective Drug Utilization Review
DEA	Drug Enforcement Agency	SAOs	Short-Acting Opioids
LAOs	Long-Acting Opioids	UM	Utilization Management
MAT	Medication-Assisted Treatment		

We're here to help

For more information, please call Provider Services at **888-362-3368**.

How to submit prior authorizations

- **Online:** Use the Prior Authorization and Notification tool in Link. For more information, go to UHCprovider.com/paan.
- **Phone:** Call **800-310-6826**
- **Fax:** Fax your completed form to 866-940-7328
- Pharmacy Prior Authorization forms are available at UHCprovider.com > Menu > Health Plans by State – choose your state > Medicaid (Community Plan) > Pharmacy Resources and Physician Administered Drugs > **Pharmacy Prior Authorization**

