

Neuropsychological Testing Under the Medical Benefit (for Tennessee Only)

Policy Number: CS083TN.O
Effective Date: October 1, 2023

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	3
Applicable Codes	3
Description of Services	3
Clinical Evidence	5
U.S. Food and Drug Administration	25
References	26
Policy History/Revision Information	31
Instructions for Use	32

Related Policies
None

Application

This Medical Policy applies to Medicaid and CoverKids in the state of Tennessee.

Coverage Rationale

Neuropsychological testing is proven and medically necessary for evaluating individuals with the following conditions when the results of testing will be used to support a diagnosis, prognosis, or treatment plan:

- Attention-deficit/hyperactivity disorder (ADHD) when all of the following are present:
 - Specific neurocognitive behavioral deficits related to ADHD need to be evaluated; and
 - Testing has been recommended by a physician and is related or secondary to a known or suspected organic-medical condition resulting from brain injury or disease process (e.g., concussion, intractable seizure disorder, cancer treatment effects, genetic disorders, inborn errors of metabolism)

Note: The scope of these criteria is applicable only to neuropsychological testing that is covered by the medical benefit. These criteria do not apply to evaluate or determine educational interventions.

- Confirmed space-occupying brain lesion including but not limited to the following:
 - Brain abscess
 - Brain tumors
 - Arteriovenous malformations within the brain
- Demyelinating disorders including multiple sclerosis
- Intellectual disability or intellectual developmental disorder when all of the following are present:
 - The intellectual disability or intellectual developmental disorder is associated with a known or suspected medical cause (e.g., Traumatic Brain Injury, in utero toxin exposure, early seizure disorder, sickle cell disease, genetic disorders); and

- The intellectual disability or intellectual developmental disorder meets all of the following criteria (DSM-5):
 - Deficits in intellectual function, such as reasoning, problem solving, planning, abstract thinking, judgment, academic learning, and learning from experience, confirmed by both clinical assessment and individualized, standardized intelligence testing
 - Deficits in adaptive functioning that result in failure to meet developmental and sociocultural standards for personal independence and social responsibility. Without ongoing support, the adaptive deficits limit functioning in one or more activities of daily life, such as communication, social participation, and independent living across multiple environments, such as home, school, work, and community; and
 - Onset of intellectual and adaptive deficits during the developmental period

Note: The scope of these criteria is applicable only to neuropsychological testing that is covered by the medical benefit. These criteria do not apply to evaluate or determine educational interventions.

- Encephalopathy including acquired immunodeficiency syndrome (AIDS) encephalopathy, human immunodeficiency virus (HIV) encephalopathy, hepatic encephalopathy, Lyme disease encephalopathy including neuroborreliosis, Wernicke's encephalopathy and systemic lupus erythematosus (SLE) encephalopathy
- Neurocognitive Disorders including Mild Cognitive Impairment (MCI), Dementia or symptoms of dementia such as memory impairment or memory loss (including Alzheimer's and extrapyramidal disorders such as Parkinson's disease) that is associated with a new onset or progressive memory loss and a decline in at least one of the following cognitive domains (DSM-5):
 - Complex attention
 - Executive function
 - Learning and memory
 - Language
 - Perceptual motor
 - Social cognition
- Neurotoxin exposure with at least one of the following:
 - Demonstrated serum levels of neurotoxins
 - Individual with one or more of the following:
 - Documented prenatal alcohol, drug, or toxin exposure
 - History of radiation therapy or chemotherapy
- Seizure disorder including individuals with epilepsy
- Stroke
- [Traumatic Brain Injury \(TBI\)](#)
- The individual is being considered for a medical or surgical procedure that may affect brain function (e.g., epilepsy surgery, resection of brain tumors or arteriovenous malformations, deep brain stimulation, stem cell or organ transplants)

The following are unproven and not medically necessary due to insufficient evidence of efficacy:

- Baseline neuropsychological testing in asymptomatic individuals at risk for sport-related concussions
- Computerized cognitive testing such as Cognivue®, Mindstreams® Cognitive Health Assessment, BrainCare™ and QbTest
- Computerized neuropsychological testing when used as a stand-alone test for evaluating concussions
- Neuropsychological testing for the following diagnoses alone without other proven conditions as noted above:
 - Headaches including migraine headache
 - History of myocardial infarction
 - Intermittent explosive disorder
- Neuropsychological testing that is comprised exclusively of self-administered or self-scored inventories, or as screening tests of cognitive function or neurological disease whether paper-and-pencil or computerized (e.g., AIMS, Folstein Mini-Mental Status Examination)
- Neuropsychological testing that is used as a routine screening tool
- Neuropsychological testing that is administered for educational or vocational purposes that do not alter or direct medical or health management
- Repeat neuropsychological testing that is not required for medical decision-making
- The individual is neurologically, cognitively, or psychologically unable to participate in a meaningful way in the neuropsychological testing process
- The individual has been diagnosed previously with brain dysfunction, such as Alzheimer's disease, and there is no expectation that neuropsychological testing would impact the individual's medical, functional, or behavioral management

Definitions

Traumatic Brain Injury (TBI): TBI is defined as a bump, blow, or jolt to the head or a penetrating head injury that disrupts the normal function of the brain (Centers for Disease Control and Prevention, 2021).

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
96116	Neurobehavioral status exam (clinical assessment of thinking, reasoning, and judgment, [e.g., acquired knowledge, attention, language, memory, planning, and problem solving, and visual spatial abilities]), by physician or other qualified health care professional, both face-to-face time with the patient and time interpreting test results and preparing the report; first hour
96121	Neurobehavioral status exam (clinical assessment of thinking, reasoning, and judgment, [e.g., acquired knowledge, attention, language, memory, planning and problem solving, and visual spatial abilities]), by physician or other qualified health care professional, both face-to-face time with the patient and time interpreting test results and preparing the report; each additional hour (List separately in addition to code for primary procedure)
96132	Neuropsychological testing evaluation services by physician or other qualified health care professional, including integration of patient data, interpretation of standardized test results and clinical data, clinical decision making, treatment planning and report, and interactive feedback to the patient, family member(s) or caregiver(s), when performed; first hour
96133	Neuropsychological testing evaluation services by physician or other qualified health care professional, including integration of patient data, interpretation of standardized test results and clinical data, clinical decision making, treatment planning and report, and interactive feedback to the patient, family member(s) or caregiver(s), when performed; each additional hour (List separately in addition to code for primary procedure)
96136	Psychological or neuropsychological test administration and scoring by physician or other qualified health care professional, two or more tests, any method; first 30 minutes
96137	Psychological or neuropsychological test administration and scoring by physician or other qualified health care professional, two or more tests, any method; each additional 30 minutes (List separately in addition to code for primary procedure)
96138	Psychological or neuropsychological test administration and scoring by technician, two or more tests, any method; first 30 minutes
96139	Psychological or neuropsychological test administration and scoring by technician, two or more tests, any method; each additional 30 minutes (List separately in addition to code for primary procedure)
96146	Psychological or neuropsychological test administration, with single automated, standardized instrument via electronic platform, with automated result only

CPT® is a registered trademark of the American Medical Association

Description of Services

Neuropsychological testing is a set of formal procedures utilizing diagnostic tests specifically focused on identifying the presence of brain damage, injury or dysfunction and any associated functional deficits. Measurement of deficits cannot be based on single test results and should always be assessed in the context of the medical and neurological examination. Neuropsychological testing is customarily associated with neurological diagnoses rather than behavioral health diagnoses.

Neuropsychological tests are administered in a variety of contexts including paper-and-pencil, computers, and visual aids. Following an initial clinical interview with a neuropsychologist, tests are strategically selected to identify specific deficits and preserved abilities. Standardized tests are then administered by a trained technician or neuropsychologist. Some tests offer multiple forms making them useful for repeated administration to the same patient, thereby minimizing practice effects. In light of the numerous procedures available for assessment of different neurocognitive functions, test selection is based on familiarity of the examiner with certain tests, availability of appropriate normative data, ability of the patient to participate in testing, and validity of particular procedures for the specific function being measured.

Neuropsychological tests include but are not limited to the following: Boston Diagnostic Aphasia Examination (BDAE), Conners' Continuous Performance Test (CCPT), Controlled Oral Word Association Test (COWAT), Delis-Kaplan Test Battery, Freedom from Distractibility Index (FFDI) from the Wechsler Intelligence Scales, Gordon Diagnostic System (GDS), Halstead-Reitan Neuropsychological Battery, Repeatable Battery for Assessment of Neuropsychological Status (RBANS), Rey Auditory Verbal Learning Test (RAVLT), Rey-Osterreith Complex Figure Test, Stroop Color and Word Test, Test of Variables of Attention (TOVA), Trail Making Tests, Wechsler Adult Intelligence Scale-Revised (WAIS-III/IV), Wide Range Assessment of Memory and Learning (WRAML), and Wisconsin Card Sorting Test (WCST). At times, neurocognitive measures are supplemented by emotional functioning and personality testing and include but are not limited to the following: Minnesota Multiphasic Personality Inventory-2 (MMPI-2)/Minnesota Multiphasic Personality Inventory-A (MMPI-A), Personality Assessment Inventory (PAI), Geriatric Rating Scale, Beck Depression Inventory (BDI), Beck Anxiety Inventory (BAI), and Rorschach Inkblot Method.

Computerized testing for dementia and cognitive impairment including the Mindstreams® Cognitive Health Assessment (NeuroTrax® Corp.) uses computer-based assessments in an attempt to identify cognitive impairment in the elderly. The software programs give individuals various stimuli or puzzles to solve using a mouse or a keypad. The Mindstreams system automatically generates a report that details the individual's performance in the standard cognitive domains, or areas, e.g., memory, attention, executive function, visual spatial perception, verbal skills, motor planning, and information processing. According to NeuroTrax, BrainCare™ is the current version of the original MindStreams product. Cognivue (Cerebral Assessment Systems, Inc.) is another computerized cognitive test that is intended for early detection of dementia signs. Individuals take the 10-minute test using the Cognivue mobile computer workstation to assess visuomotor coordination, perceptual processing, and memory. Cognivue is intended to help identify patients who may be in the early stages of dementia and should undergo further evaluation. The QbTest is an online computerized test that measures activity, attention, and impulsivity for assessment of attention-deficit hyperactivity disorder (ADHD).

Computerized neuropsychological tests have been proposed to be used as part of the overall medical management of concussion to monitor recovery. Most computer-based cognitive assessment tools are designed to detect the speed and accuracy of attention, memory, and thinking ability. Currently available computerized tests include ImPACT (Immediate Post-Concussion Assessment and Cognitive Testing, ImPACT Applications, Inc.), ANAM (Automated Neuropsychological Assessment Metrics, the United States Army Medical Department), CogState Sport (Axon Sports, Ltd.), and HeadMinder (Headminder, Inc.). These tests are being investigated for baseline testing of asymptomatic persons and managing concussions once they occur.

Neuropsychological testing is within the scope of the provider's professional training and licensure when the provider is any of the following:

- A doctoral-level psychologist who is licensed to practice independently and demonstrates sufficient training and experience
- A credentialed psychiatrist who meets the following requirements:
 - Recognized certification in neurology through the American Board of Psychiatry and Neurology;
 - Accreditation in behavioral neurology and neuropsychiatry through the American Neuropsychiatric Association;
 - State medical licensure specifically allowing for the provision of neuropsychological testing service(s);
 - Evidence of professional training and expertise in the specific tests and/or assessment measures for which authorization is requested;
 - Physician and supervised psychometrician(s) adhere to the prevailing national professional and ethical standards regarding test administration, scoring, and interpretation
- A board-certified neurologist

Refer to the following Optum Supplemental Clinical Criteria for more information:

- Psychological/ Neuropsychological Testing Guidelines (to access this guideline, go to: [Optum Provider Express>Clinical Resources>Guidelines/Policies/Manuals>Supplemental Clinical Criteria](#)). Accessed May 15, 2023.

Clinical Evidence

Attention Deficit Hyperactivity Disorder (ADHD)

Becke et al. (2023) administered a comprehensive neuropsychological test battery in an analogue study that included 57 adults with ADHD, and 211 university students who were divided into two groups with 60 students in the control group, and 151 students in the simulator group to evaluate individual test's utility in detecting noncredible performance. Participants in the simulator group were then divided to receive one of three sets of instructions: naive simulators received general instructions to feign ADHD and no additional information, symptom-coached simulators were given the DSM diagnostic criteria of ADHD, and fully coached simulators received information on both the neuropsychological assessment of ADHD and its diagnostic criteria. Analysis by the authors of the test results demonstrated that the Simulation Group showed a higher median number of test results falling into the suspect range based on the newly derived cut-off scores than the ADHD Group and Control Group. The authors reported that all of the tests ensured at least 90% specificity in the ADHD Group but that sensitivity differed significantly between tests, ranging from 0% to 64.9%. They also found that tests focusing on selective attention, vigilance, and inhibition were most useful in detecting the instructed simulation of adult ADHD, while tests focusing on figural fluency and task switching lacked sensitivity. Limitations of the study include the single center and simulation study design, the lack of heterogeneity in the pool used to select the control and simulation groups and the risks associated with the embedded validity indicators in some of the tests and from the risk of overfitting. The authors concluded .

Pagán et al. (2023) performed a systematic review to assess the diagnostic utility of the Conner's Continuous Performance Test (CCPT) for diagnosing ADHD in adults. Their review and analysis included 35 published studies with sample sizes ranging between 24 and 413 participants that assessed symptoms for both childhood and adults. The authors stated that there was moderate reliability, subpar discriminant and ecological validity, and mixed sensitivity and specificity for the CCPT. They concluded that their review gave support to previous critiques of the CCPT's diagnostic and utility as a treatment measure and stated that clinicians should assess information from multiple sources when diagnosing ADHD in an adult patient. Limitations of the study include the exclusion of adults with comorbidities from most of the included studies, the heterogeneity in study designs of the studies reviewed including how ADHD was assessed between studies, the instruments being used, the diagnostic measures and outcomes, and the lack of control groups in many of the included studies.

Bechtel et al. (2012) evaluated whether boys with epilepsy-related ADHD and developmental ADHD share a common behavioral, pharmaco-responsive, and neurofunctional pathophysiology. Seventeen boys with diagnosed combined epilepsy/ADHD, 15 boys with developmental ADHD, and 15 healthy controls (aged 8-14 years) performed on working memory tasks (N-back) while brain activation was recorded using functional magnetic resonance imaging. On a behavioral level, boys with epilepsy-related ADHD as well as those with developmental ADHD performed similarly poorly on tasks with high cognitive load when compared to healthy controls. On the functional level, both patient groups showed similar reductions of activation in all relevant parts of the functional network of working memory when compared to controls. The study data showed strong similarities between epilepsy-related and developmental ADHD on the behavioral, pharmaco-responsive, and neural level, favoring the view that ADHD with and without epilepsy shares a common underlying neurobehavioral pathophysiology.

Dementia, Possible Dementia, Memory Loss/Impairment and Mild Cognitive Impairment (MCI)

For memory impairment or dementia screening, a single test of global measures of function or a measure of cognitive function is usually administered along with a test of behavioral or emotional symptoms. In addition to brief screening tests, for some patients, comprehensive neuropsychological testing may be indicated to confirm a diagnosis, evaluate effects of treatment, and assist in designing rehabilitative or intervention strategies for the patient. Standardized test batteries are too long for most patients with dementia; specialized dementia batteries or an individualized test battery is usually more appropriate.

A definitive diagnosis of Alzheimer's disease is based on the presence of memory deficits along with deficits in at least one other aspect of cognition, and in some cases is made on neuropsychological test results alone (Talwalker, 1996). Impairment in primary (short-term) memory alone is not a useful diagnostic marker for Alzheimer's disease in the early stages. Tests of delayed recall (long-term memory) and retrieval of facts of common knowledge have been shown to be the most useful measures to distinguish normal aging and early Alzheimer's disease. Dementia due to Alzheimer's disease can be distinguished

from dementia due to vascular disease by differences in pattern of memory impairment and the progressive nature of Alzheimer's disease (Costa et al., 2017). Careful interpretation of test results, taken in conjunction with medical findings, allows differentiation of Alzheimer's disease from normal memory loss due to aging, and from vascular dementia.

In a test validation study on the cross-cultural dementia (CCD) screening test for diagnosing Alzheimer's disease and Parkinson's disease, Delgado-Álvarez et al. (2023) recruited 150 participants from a single outpatient center and divided them into three groups with 30 participants with Alzheimer's disease with mild dementia (AD-D), 30 participants with Alzheimer's disease in mild cognitive impairment (AD-MCI), 30 participants with mild cognitive impairment associated with Parkinson's disease (PD-MCI) and 60 participants in the healthy control (HC) group (50% for comparisons with AD, 50% for comparisons with PD-MCI) with no significant differences in age, education, and sex. A comprehensive neuropsychological test battery and the CCD screening test were completed for each participant. The authors reported that intergroup differences were found according to the cognitive profile of each clinical condition and that the CCD test described differences in executive functions and speed scores comparing AD-MCI and PD-MCI. They also noted correlations between standardized neuropsychological tests and CCD measures which they stated support the convergent validity of the CCD test. The authors concluded that the CCD test showed good discrimination properties and cut-off scores for dementia and that the CCD test would be useful as a novel cognitive tool in the assessment of patients with cognitive impairment in different neurological conditions. Limitations of the study included the single-center design, the lack of a group of participants with Parkinson's disease that were cognitively preserved, the generally low level of education of the participants, and the lack of evaluation of cognitive reserve.

The Agency for Healthcare Research and Quality (AHRQ) wanted to identify which individual cognitive tests or combinations of cognitive tests are most accurate for clinically diagnosing clinical Alzheimer's-type dementia (CATD). AHRQ noted that there were no evidence-based guidelines about the merits of either brief cognitive testing or comprehensive neuropsychological testing in this patient population and that access to comprehensive neuropsychological testing is limited in many clinical settings. Fink et al (2020) completed a Comparative Effectiveness Review for AHRQ in which they analyzed 56 studies on the accuracy of brief cognitive tests for CATD and found that multiple brief cognitive tests were highly sensitive and specific for distinguishing CATD from normal cognition, but less so for distinguishing mild CATD from normal cognition or CATD from mild cognitive impairment (MCI).

In a systematic review and meta-analysis, Belleville et al. (2017) determined the extent to which cognitive measures can predict progression from mild cognitive impairment (MCI) to Alzheimer's type dementia (AD), assessed the predictive accuracy of different cognitive domain categories, and determined whether accuracy varies as a function of age and length of follow-up. The authors systematically reviewed and meta-analyzed data from longitudinal studies reporting sensitivity and specificity values for neuropsychological tests to identify individuals with MCI who will develop AD. Twenty-eight studies met the eligibility criteria (2365 participants) and reported predictive values from 61 neuropsychological tests with a 31-month mean follow-up. Values were pooled to provide combined accuracy for 14 cognitive domains. Many domains showed very good predictive accuracy with high sensitivity and specificity values. Verbal memory measures and many language tests yielded very high predictive accuracy. Other domains (e.g., executive functions, visual memory) showed better specificity than sensitivity. Predictive accuracy was highest when combining memory measures with a small set of other domains or when relying on broad cognitive batteries. The authors concluded that neuropsychological assessment can strongly contribute to predicting dementia while individuals are still in the MCI phase. According to the authors, cognitive tests are excellent at predicting MCI individuals who will progress to dementia and should be a critical component of any toolkit intended to identify AD at the pre-dementia stage.

Pedersen et al. (2017) examined the incidence, progression, and reversion of mild cognitive impairment in patients with Parkinson disease (PD-MCI) over 5 years. A population-based cohort of patients with incident PD underwent repeated neuropsychological testing of attention, executive function, memory, and visuospatial abilities at baseline (n = 178), 1 year (n = 175), 3 years (n = 163), and 5 years (n = 150). Patients were classified as PD-MCI and diagnosed with dementia according to published criteria. Thirty-six patients (20.2%) fulfilled criteria for PD-MCI at baseline. Among those with normal cognition at baseline (n = 142), the cumulative incidence of PD-MCI was 9.9% after 1 year, 23.2% after 3 years, and 28.9% after 5 years of follow-up. Overall, 39.1% of patients with baseline or incident PD-MCI progressed to dementia during the 5-year study period. The conversion rate to dementia was 59.1% in patients with persistent PD-MCI at 1-year vs 7.2% in those with normal cognition during the first year. A total of 27.8% of patients with baseline PD-MCI and 24.2% of those with incident PD-MCI had reverted to normal cognition at study end, but the reversion rate decreased to 9.4% in those with persistent PD-MCI at 2 consecutive visits. Compared with cognitively normal patients, PD-MCI reverters within the first 3 years of follow-up were at increased risk of

subsequently developing dementia. The authors concluded that early PD-MCI, regardless of persistence or reversion to normal cognition, has prognostic value for predicting dementia in patients with PD.

Weissberger et al. (2017) conducted a systematic review and meta-analysis measuring the sensitivity and specificity for individuals with Mild Cognitive Impairment (MCI) versus Alzheimer's Dementia (AD); AD versus healthy controls (HC), and MCI versus HC utilizing neuropsychological assessments. Memory measures were divided into four categories, immediate, delayed, associative learning and other. Immediate memory testing is a recall of information directly following presentation of stimuli. Delayed memory used a distraction task or minutes of delay before recall. Associative learning tasks required individuals to put together stimulus pairs such as word pair and/or object and location. The other category was recognition memory combined with immediate recall score with yes/no recognition score or interference scores and other miscellaneous tests.

AD versus healthy control (HC) studies (n = 47) revealed generally high sensitivity and specificity ($\geq 80\%$ for AD comparisons) for measures of immediate (sensitivity = 87%, specificity = 88%) and delayed memory (sensitivity = 89%, specificity = 89%), especially those involving word-list recall. Examination of MCI versus HC studies (n = 38) revealed generally lower diagnostic accuracy for both immediate (sensitivity = 72%, specificity = 81%) and delayed memory (sensitivity = 75%, specificity = 81%). Measures that differentiated AD from other conditions (n = 10 studies) yielded mixed results, with high sensitivity in the context of low or variable specificity. The authors concluded memory measures have high diagnostic accuracy for identification of AD, but require further development for the identification of MCI. Further research is needed focusing on specific types of MCI and Alzheimer's disease.

Developmental Disorders

In general, empirical data, rather than evidence from prospective studies with long-term follow-up, support the use of neuropsychological testing for developmental disorders in infants and children.

Trickett et al. (2022) conducted a parallel mediation analysis with cross-sectional data from 152 extremely premature (EP; < 27 weeks of gestation) children and 120 term-born controls who were assessed at age 11 to identify specific cognitive mechanisms that are associated with poor academic attainment in children born preterm for the development of interventional strategies. Mathematics and reading attainment was evaluated to assess the following mediators: verbal working memory, visuospatial working memory, verbal processing speed, attention, and visuospatial processing. The authors reported that children born EP had significantly lower mean composite mathematics and reading scores than controls equating to a deficit of -1.1 SD in reading and -1.4 SD in mathematics, after adjusting for sex and socio-economic status. When children with severe neurodevelopmental disability were excluded, the difference in means, adjusted for sex and socio-economic status, remained significant for reading (-0.4 SD) and mathematics (-1.0 SD). Lower scores were also reported by the authors for verbal working memory (-0.5 SD), visuospatial working memory (-0.7 SD), attention (-0.6 SD), visuospatial processing (-1.1 SD) and verbal processing speed (-0.6 SD) although the magnitude of difference in all five neuropsychological skills were decreased but still significant when children with severe disability were excluded. The authors concluded that children born before 27 weeks of gestation had substantially poorer attainment in reading and mathematics compared to children born at term and that their study identified that a combination of neuropsychological skills including verbal working memory, visuospatial working memory and visuospatial processing may be especially important to target in interventions to improve mathematics and reading outcomes for EP children with average to moderately low IQ. Limitations of the study include the inability of some EP children with severe neurodevelopmental disabilities to complete the entire battery of tests, the recruitment of controls from mainstream schools as this may not be reflective of the general population, the limited time available to assess each child within school, and the use of cross-sectional data for mediation models.

Johnson et al. (2016) explored comorbidity in intellectual disability (ID) and learning disabilities (LD) in children born extremely preterm (EP; < 26 + 0 weeks gestation). A UK national cohort of 161 EP children and 153 term-born controls without neurosensory impairments was assessed at 11 years of age (the EPICure Study). Neuropsychological abilities commonly affected by EP birth were assessed using the NEPSY Developmental Neuropsychological Test. Overall, 75 (47%) EP children and 7 (4.6%) controls had ID or LD. Comorbidity in ID/LD was more common among EP children than controls (24% vs. 0%). EP children with comorbid ID/LD had significantly poorer neuropsychological abilities and curriculum-based attainment than EP children with an isolated disability or no disabilities. LD was associated with a 3 times increased risk for special educational needs (SEN). However, EP children with ID alone had poorer neuropsychological abilities and curriculum-based attainment than children with no disabilities, yet there was no increase in SEN provision among this group. The authors concluded that EP children are at high risk for comorbid intellectual and learning disabilities. According to the authors, education professionals

should be aware of the complex nature of EP children's' difficulties and the need for multi-domain assessments to guide intervention.

Hartman et al. (2010) examined the motor skills and executive functions in school-age children with borderline and mild intellectual disabilities (ID). Sixty-one children aged between 7 and 12 years diagnosed with borderline ID (33 boys and 28 girls; 71 < IQ < 79) and 36 age peers with mild ID (24 boys and 12 girls; 54 < IQ < 70) were assessed. Their abilities were compared with those of 97 age- and gender-matched typically developing children. Qualitative motor skills, i.e., locomotor ability and object control, were evaluated with the Test of Gross Motor Development (TGMD-2). Executive functioning (EF), in terms of planning ability, strategic decision-making and problem solving, was gauged with the Tower of London (TOL) task. Compared with the reference group, the full ID cohort scored significantly lower on all assessments. According to the investigators, the study results support the notion that besides being impaired in qualitative motor skills, intellectually challenged children are also impaired in higher-order executive functions. The authors conclude that deficits in the two domains are interrelated, so early interventions boosting their motor and cognitive development are recommended.

Traumatic Brain Injury

Longitudinal and case controlled studies along with numerous case reports support the use of neuropsychological tests to assess the severity of injury and the prognosis for patients with closed head trauma, to monitor progression, and to provide measures of outcome for determining degree of recovery (Filipčíková et al., 2022; Hanks et al., 2016; Carozzi et al., 2015).

Other Disorders

Neuropsychological testing may have a role in the clinical management of the following medical disorders:

- Brain lesions including abscesses, tumors, and arteriovenous malformations in the brain (Söderström et al., 2022; Pranckeviciene et al., 2017; Meskal et al., 2016; Walsh et al., 2016; Cochereau et al., 2016)
- Demyelinating disease including multiple sclerosis (Delgado-Álvarez et al., 2022; Fuchs et al., 2022; Tekin et al., 2022; Böttrich et al., 2020; Wojcik et al., 2019; von Bismarck et al., 2018; Ruet and Brochet, 2018; Vollmer et al., 2016)
- Encephalopathy (Sigurdardottir et al., 2022; Rayes et al., 2019; Moore et al., 2017; Burton et al., 2017)
- Epilepsy and seizure disorders (Silberg et al., 2020; Parra-Díaz and García-Casares, 2017; Grau-López et al., 2017; Wilson et al., 2015; Filippini et al., 2016; Patrikelis et al., 2016)
- Neurotoxin exposure (Nascimento et al., 2016)
- Stroke (Zuo et al., 2022; Lo Buono et al., 2018; Tan et al., 2017; Zweifel-Zehnder et al., 2015; Chen et al., 2015)

Computerized Neuropsychological Testing for Concussion

The evidence is insufficient to establish the validity and reliability of computerized tests to evaluate concussions. Prospective controlled trials are needed to demonstrate the clinical utility of these tests to detect impairment following concussion.

Ivins et al. (2022) completed an initial psychometric analysis of the Brain Gauge (BG) personal computer-based test battery to evaluate its potential use for evaluating patients with acute mild traumatic brain injury (mTBI). The study participants were 73 military service members (SM) who were assessed within 7 days of their injury at military medical treatment facilities (emergency department (30.1%), primary care (41.1%) or a TBI specialty clinic (28.8%) and 100 healthy service members as a control group. Prior to completing the BG, participants were administered a demographic and military questionnaire, the Neurobehavioral Symptom Inventory (NSI), a PTSD Checklist (PCL-5), and a Patient Health Questionnaire (PHQ-9). The authors reported that SMs with mTBI had statistically significant worse performance on both BG Reaction Time (RT) tests and the Sequential Amplitude Discrimination test as well as having a significantly lower whole-battery composite. The authors stated that, while particular subtests of BG are sensitive to the effects of acute mTBI, there was questionable clinical utility of these scores and that the mTBI group performed worse on some tests, than the control group. The authors noted that the base rate analysis revealed that a minority of those with mTBI had multiple scores at or near potentially clinically meaningful performance thresholds, contradicting the very high diagnostic accuracy statistics published by BG's developers which raised concerns about the use of an aggregate score from the BG test battery. Limitations of the study include the small sample size, the delay of up to 7 days post injury which may have influenced results due to possible cognitive recovery that had occurred and the inclusion of only SMs in the study. The authors concluded that their study did find that SMs with acute mTBI on average performed worse than healthy control SMs on the BG Cortical Metric Symptom Score, the BG RT tests and the Sequential Amplitude Discrimination test but that the results also demonstrated that overall, BG does not distinguish mTBI cases from controls at a clinically meaningful rate, and not nearly at the rates previously reported in the literature.

In a prospective longitudinal observational cohort study, Takagi et al. (2019) examined whether cognitive functioning (measured by CogSport) has prognostic value for predicting rapid versus slow recovery. Data were collected at 1-4, 14-, and 90-days post-injury. Eligible children were aged ≥ 5 and < 18 years presenting to the Emergency Department having sustained a concussion within 48 hours. Concussion was defined according to the Zurich/Berlin Consensus Statement on Concussion in Sport. Dependent variables were reaction times and error rates on the CogSport Brief Battery. In total, 220 cases were analyzed: 98 in a rapid recovery group (asymptomatic at 14 days post-injury, mean age 11.5 [3.2], 73.5% male) and 122 in a slow recovery group (symptomatic at 14 days post-injury, mean age 12.0 [3.1], 69.7% male). Longitudinal GEE analyses modeled the trajectories of both mean log₁₀-transformed reaction time and error rates between groups over time (1-4, 14 and 90 days). Both group main and interaction (time by group) terms for all models were non-significant ($p > .05$). The authors concluded that cognitive functioning, measured by CogSport and assessed within 1-4 days of concussion, does not predict prolonged recovery in a pediatric sample. Further, there were no significant group differences at any time point. The authors stated that considering the widespread use and promotion of Computerized neuropsychological tests (CNTs), it is important that clinicians understand the significant limitations of the CogSport battery.

Ivins et al. (2019) assessed the agreement between the following four brief computerized neurocognitive assessment tools by comparing rates of low scores: Automated Neuropsychological Assessment Metrics (ANAM); CogState, also known as CogSport or Axon Sports; Central Nervous System Vital Signs (CNSVS); and Immediate Post-concussion Assessment and Cognitive Test (ImPACT). Four hundred and six US Army service members (SMs) with and without acute mild traumatic brain injury completed two randomly assigned CNTs with order of administration also randomly assigned. A base rate analysis was performed for each CNT to determine the proportions of SMs in the control and mild traumatic brain injury (mTBI) groups who had various numbers of scores that were 1.0 +, 1.5 +, and 2.0 + standard deviations below the normative mean. These results were used to identify a hierarchy of low score levels ranging from poorest to least poor performance. The agreement was compared between every low score level from each CNT pair administered to the SMs. More SMs in the mTBI group had low scores on all CNTs than SMs in the control group. As performance worsened, the association with mTBI became stronger for all CNTs. Most if not all SMs who performed at the worst level on any given CNT also had low scores on the other CNTs they completed but not necessarily at an equally low level. The authors indicated that the psychometric comparability and clinical utility of these CNTs are not well understood and until such studies are done it will not be possible to make any judgments about which CNT, if any, is superior to the others. The authors state that until more evidence emerges, these CNTs should be used cautiously and only as one source of information from among many other types of clinical assessments. None of them should be used as a definitive or standalone diagnostic tool. An important limitation of this study is that there were relatively small numbers of SMs in each CNT pair who performed at the poorest levels so the results of this study should be treated as preliminary. Another limitation is that the data is from military service members and these findings may not be generalizable to other populations such as high school and college athletes.

Cole et al. (2018) investigated the validity of four computerized neurocognitive assessment tools (NCATs): the ANAM4, CNS-VS, CogState, and ImPACT. Two NCATs were randomly assigned and a battery of traditional neuropsychological (NP) tests administered to 272 healthy control active-duty service members and to 231 service members within 7 days of an mTBI. Analyses included correlations between NCAT and the NP test scores to investigate convergent and discriminant validity, and regression analyses to identify the unique variance in NCAT and NP scores attributed to group status. Effect sizes (Cohen's f^2) were calculated to guide interpretation of data. Only 37 (0.6%) of the 5,655 correlations calculated between NCATs and NP tests are large. The majority of correlations are small, with no clear patterns suggestive of convergent or discriminant validity between the NCATs and NP tests. Though there are statistically significant group differences across most NCAT and NP test scores, the unique variance accounted for by group status is minimal with effect sizes indicating small to no meaningful effect. The authors concluded that although the results are not overly promising for the validity of the four NCATs investigated, traditional methods of investigating psychometric properties may not be appropriate for computerized tests.

Broglio et al. (2018) evaluated the test-retest reliability of commonly implemented and emerging concussion assessment tools across a large nationally representative sample of student-athletes. The assessments were divided into mandatory (Level A measures) and optional emerging concussion measures (Level B measures). Level A measures included Immediate Post-Concussion Assessment and Cognitive Test (ImPACT), Automated Neuropsychological Assessment Metrics, and the Cogstate Computerized Cognitive Assessment Tool (CCAT, formerly named Axon). Participants ($n = 4874$) from the Concussion Assessment, Research, and Education Consortium completed annual baseline assessments on two or three occasions. Each assessment included measures of self-reported concussion symptoms, motor control, brief and extended neurocognitive function, reaction time, oculomotor/oculovestibular function, and quality of life. The authors concluded that this investigation noted less than optimal reliability for most common and emerging concussion assessment tools. None of the assessment tools

met or exceeded the accepted threshold for clinical utility. According to the authors, the use of these tools is still necessitated by the absence of a gold standard diagnostic measure, with the ultimate goal of developing more refined and sound tools for clinical use.

The Centers for Disease Control and Prevention (CDC) National Center for Injury Prevention and Control Board of Scientific Counselors, a federal advisory committee, established the Pediatric Mild Traumatic Brain Injury Guideline Workgroup and developed a guideline based on a previous systematic review of the literature (Lumba-Brown et al., 2018) to obtain and assess evidence toward developing clinical recommendations for health care professionals related to the diagnosis, prognosis, and management/treatment of pediatric mild traumatic brain injury (mTBI). The CDC guideline included the recommendations on the diagnosis, prognosis, and management/treatment of pediatric mTBI that were assigned a level of obligation (i.e., must, should, or may) based on confidence in the evidence. Regarding computerized cognitive testing, the CDC stated that health care professionals may use validated, age-appropriate computerized cognitive testing in the acute period of injury as a component of the diagnosis of mTBI (moderate; level C).

In a consensus statement, the 5th International Conference on Concussion in Sport states that the use of neuropsychological testing (NP) contributes significant information in concussion assessment. Brief computerized cognitive evaluation tools are a commonly utilized component of these assessments worldwide given the logistical limitation in accessing trained neuropsychologists. However, it should be noted that these are not substitutes for complete NP assessment. For children, it is recommended that age-specific validated symptom-rating scales be used in sport-related concussion (SRC) assessment, and further research is required to establish the role and utility of computerized NP testing in this age group. The consensus statement suggests that baseline testing may be useful but is not necessary for interpreting post-injury scores (McCrory et al., 2017).

Farnsworth et al. (2017) analyzed reliability data for computerized neurocognitive tests (CNTs) using meta-analysis and examined moderating factors that may influence reliability. Studies were included in the meta-analysis if they met all of the following criteria: used a test-retest design, involved at least 1 CNT, provided sufficient statistical data to allow for effect-size calculation, and were published in English. Two independent reviewers investigated each article to assess inclusion criteria. Eighteen studies involving 2674 participants were retained. Intraclass correlation coefficients were extracted to calculate effect sizes and determine overall reliability. Moderator analyses were conducted to evaluate the effects of the length of the test-retest interval, intraclass correlation coefficient model selection, participant demographics, and study design on reliability. Heterogeneity was evaluated using the Cochran Q statistic. The proportion of acceptable outcomes was greatest for the Axon Sports CogState Test (75%) and lowest for the Immediate Post-Concussion Assessment and Cognitive Testing (ImPACT) (25%). Moderator analyses indicated that the type of intraclass correlation coefficient model used significantly influenced effect-size estimates, accounting for 17% of the variation in reliability. The authors concluded that the Axon Sports CogState Test, which has a higher proportion of acceptable outcomes and shorter test duration relative to other CNTs, may be a reliable option; however, future studies are needed to compare the diagnostic accuracy of these instruments (The Nakayama et al. (2014) study which was previously cited in this policy, is included in the Farnsworth et al. (2017) meta-analysis).

Gaudet and Weyandt (2017) conducted a systematic review of existing research investigating Immediate Post-Concussion and Cognitive Testing (ImPACT) and the prevalence of invalid baseline results including the effectiveness of ImPACT's embedded invalidity indicators in detecting suspect effort. Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines were followed in order to systematically structure a search across four databases and analysis of studies that presented data related to the prevalence of invalid performance and/or the effectiveness of ImPACT's embedded invalidity indicators. A total of 17 studies included prevalence rates of invalid performances or examined the effectiveness of ImPACT's embedded invalidity indicators. Of the 17 studies, 12 included prevalence rates of invalid baseline results; and across this group of studies (after removing an outlier), the weighted prevalence rate of invalid baseline results was 6%. Four of the 17 studies examined the effectiveness of ImPACT's embedded invalidity indicators. ImPACT's embedded invalidity indicators correctly identified suboptimal effort in approximately 80% of individuals instructed to perform poorly and avoid detection ('coached') or instructed to perform poorly ('naïve'). According to the authors, these findings raise a number of issues pertaining to the use of ImPACT. Invalid performance incidence may increase with large group versus individual administration, use in nonclinical settings, and among those with Attention Deficit-Hyperactivity Disorder or learning disability. Additionally, the older desktop version of ImPACT appears to be associated with a higher rate of invalid performances than the online version. Although ImPACT's embedded invalidity indicators detect invalid performance at a rate of 6% on average, known group validity studies suggest that these measures miss invalid performance approximately 20% of the time when individuals purposefully underperform.

Nelson et al. (2017) evaluated the reliability and validity of three computerized neurocognitive assessment tools (Automated Neuropsychological Assessment Metrics [ANAM], Defense Automated Neurobehavioral Assessment [DANA], and Immediate Post-Concussion Assessment and Cognitive Testing [ImPACT]) for assessing mild traumatic brain injury (mTBI). The study included mTBI (n = 94) and matched trauma control (n = 80) subjects recruited from a level I trauma center emergency department (ED) completed symptom and neurocognitive assessments within 72 hours of injury and at 15- and 45-days post-injury. Concussion symptoms were also assessed via phone at 8 days post-injury. Computerized neurocognitive assessment tools (CNTs) did not differentiate between groups at any time point. Roughly a quarter of stability coefficients were over .70 across measures and test-retest intervals in controls. The authors concluded that the CNTs evaluated, developed, and widely used to assess sport-related concussion, did not yield significant differences between patients with mTBI versus other injuries. Symptom scores better differentiated groups than CNTs, with effect sizes weaker than those reported in sport-related concussion studies. According to the authors, nonspecific injury factors, and other characteristics common in ED settings, likely affect CNT performance across trauma patients as a whole and thereby diminish the validity of CNTs for assessing mTBI in this patient population. The authors indicated that this investigation had several limitations. First, subjects were evaluated in a laboratory setting within 72 hr of injury; thus, it is possible that stronger group differences in clinical assessment measures would have been found had subjects been assessed more acutely (such as within the ED). Second, the study design (i.e., assignment of two of three CNTs to each subject) and presence of loss to follow-up (16% at 45 days post-injury) contributed to smaller sample sizes (< 50) for some CNT measures and at some time points.

In a systematic review, Alsalaheen et al. (2016) assessed the literature on the reliability of the Immediate Post-Concussion Assessment and Cognitive Testing (ImPACT). Ten studies that met the inclusion criteria were included in the review. With the exception of processing speed, all composite scores consistently exhibited poor to moderate reliability. When considering 2 time points, participants who were misclassified as experiencing a "reliable change" in any score ranged between 5% and 26% for verbal memory, 2.2% and 19.6% for visual memory, 4% and 24% for processing speed, and 4% and 23.2% for reaction time. The authors concluded that the majority of ImPACT composite scores did not consistently demonstrate good reliability. According to the authors, clinicians should be cautious when ImPACT is used as a criterion for medical clearance to return to play after concussion. Because of its widespread use in concussion-related clinical research, researchers must exercise due diligence when utilizing ImPACT to evaluate outcomes after concussion or to validate other outcome measures. (Cited in Farnsworth et al., 2017).

Nelson et al. (2016) evaluated the reliability and validity of three computerized neurocognitive tests (CNTs): ANAM, Axon Sports/Cogstate Sport, and ImPACT-in a common sample. High school and collegiate athletes completed two CNTs each at baseline. Concussed (n = 165) and matched non-injured control (n = 166) subjects repeated testing within 24 hours and at 8-, 15-, and 45-days post-injury. Group differences in performance were mostly moderate to large at 24 hours and small by day 8. The sensitivity of reliable change indices (RCIs) was best at 24 hours (67.8%, 60.3%, and 47.6% with one or more significant RCIs for ImPACT, Axon, and ANAM, respectively) but diminished to near the false positive rates thereafter. Across time, the CNTs' sensitivities were highest in those athletes who became asymptomatic within 1 day before neurocognitive testing but was similar to the tests' false positive rates when including athletes who became asymptomatic several days earlier. Test-retest reliability was similar among these three CNTs and below optimal standards for clinical use on many subtests. The authors indicated that their findings suggest that the clinical utility of CNTs in the context of SRC management is maximal very soon (within 24 hours) after injury or after symptom resolution and quite limited at later time points (day 8 and beyond). According to the authors, the rapid clinical recovery course from concussion and modest stability probably jointly contribute to limited signal detection capabilities of neurocognitive tests outside a brief post-injury window.

Kontos et al. (2014) performed a meta-analysis assessing the effects of sport-related concussion as measured by computerized neurocognitive tests (NCT) 1-week post injury. Thirty-seven studies involving 3960 participants between 2000 and 2011 were included. Code substitution, visual memory, processing speed, and memory tasks demonstrated negative effects for concussion. Younger adolescents had lower NCT performance than older adolescents and college aged athletes. ImPACT studies demonstrated a negative effect for concussion as did those involving contact sports. The authors found that computerized neurocognitive testing results suggest athletes suffer impairments within one week of a concussion. Several factors such as age, type of neurocognitive test, and test administrator may lead to more pronounced impairments. The authors indicated that no single tool can or should be used to measure the effect of concussion. Instead, clinicians and researchers should adopt a comprehensive approach to assessing this injury.

Echemendia et al. (2013) critically reviewed the literature from the past 12 years regarding key issues in sports-related neuropsychological assessment of concussion. Based on the review of the literature, the authors concluded that traditional and

computerized neuropsychological tests are useful in the evaluation and management of concussion. A brief cognitive evaluation tools cannot substitute for formal neuropsychological assessment. According to the authors, there is insufficient evidence to recommend the widespread routine use of baseline neuropsychological testing.

Hang et al. (2015) determined if computerized neurocognitive testing (Immediate Post-Concussion Assessment and Cognitive Testing [ImPACT]) in the emergency department (ED) can be used as a prognostic tool to detect young athletes at risk of having protracted concussive symptoms. This was a prospective cohort study of athletes aged 11 to 18 years who presented to an ED less than 24 hours after sustaining a sports-related concussion. ImPACT was administered in the ED, and performance was categorized as "poor" if the athlete had 3 (of 4) or greater low domain scores. Participants completed the Post-Concussion Symptom Scale (PCSS) in the ED and by phone at 1 and 2 weeks after injury. Athletes were symptomatic if their PCSS score was more than 6 in males and more than 8 in females. One hundred nine patients were enrolled; 60% and 36% remained symptomatic at 1 and 2 weeks after injury, respectively. "Poor" ImPACT performance was not particularly useful in predicting athletes with protracted symptoms. In bivariate analysis, a higher ED PCSS score was associated with protracted symptoms. The authors concluded that computerized neurocognitive testing in the ED has limited usefulness in predicting protracted symptoms. Total acute symptom burden may be a useful prognostic tool in the ED evaluation of concussed young athletes, yet further research is necessary.

Baseline Neuropsychological Testing for Concussion

There is insufficient evidence to indicate that the use of baseline neuropsychological testing in athletes or other individuals alters risk from concussion. There is insufficient evidence that baseline tests influence physician decision-making or outcomes of treatment of concussion.

Cosgrave et al. (2023) completed a prospective cohort study with 135 school-aged (15 to 19 years) rugby players from 5 schools to explore whether the Sports Concussion Assessment Tool (SCAT), Cogstate Brief Battery (CBB) and the King-Devick test (K-D test) can be used to monitor concussion status through to full recovery. In this study, all participants completed baseline tests in the preseason where it was found that 61 (45.2%) reported a prior lifetime history of sport related concussion (SRC) and 64 (48%) participants reported symptoms on their baseline SCAT (mean 3.3; range 1-16) with the most common symptoms being fatigue/low energy (31%), neck pain (16%) and irritability (14%). The season consisted of 25 training weeks and 18 games on average across the five teams. During the season, 16 participants experienced 18 SRCs with 9 (56.3%) of the participants having a prior history of SRC. These participants underwent weekly post-concussion testing starting within 1-7 days of injury (mean 3.9 days) with the full battery of tests and an individualized rehabilitation program until recovered. One participant remained symptomatic at 87 days and was referred to a neurologist. Of the remaining 17 concussions, mean severity was 20 days (range 4-42 days). Participants with SRC underwent 52 post-concussion CBB assessments with results consistent with clinically assessed recovery status on 27 (51.9%) occasions. The CBB had a false positive rate of 33% and test specificity was 67%. On 7 (13.5%) occasions participants failed the CBB when clinically they were deemed to be recovered from their concussion. The CBB had a false negative rate of 58% and test sensitivity was 42%. On 18 (34.6%) occasions participants passed the CBB when clinically they were deemed not recovered from their concussion. There were 50 post-concussion K-D tests performed that had results consistent with clinically assessed recovery status on 32 (64%) occasions. The K-D test had a false positive rate of 11% and test specificity was 90%. On 2 (4%) occasions participants failed the K-D when clinically they were deemed to be recovered from their concussion. The K-D test had a false negative rate of 52% and test sensitivity was 48%. On 16 (32%) occasions participants passed the K-D when clinically they had been deemed not recovered from their concussion. The authors reported that the CBB and K-D tests were poorly associated with clinical assessment and produced high false negative rates of 0.58 and 0.52, respectively. The authors concluded that analysis of clinical recovery with CBB and K-D test revealed a relatively poor ability to accurately monitor concussion status compared to clinical assessment. The authors stated that their findings suggest that these tools not be used in isolation for monitoring SRC recovery in adolescents. Limitations of the study include the lack of objective measures of concussion recovery, dependence of concussion detection on school medical staff, the small sample size and concussion incidence and the lack of a follow-up comparison of the tests on the participants who did not sustain an SRC.

Tsushima et al. (2019) identified valid, invalid (identified by five embedded Invalidity Indicators), and sandbagging (identified by three "red flags") results in the ImPACT baseline test scores of 6,346 high school athletes. In addition, the ImPACT post-concussion scores of 266 athletes who sustained a concussion during the school year were evaluated to compare the baseline-to-post concussion changes of valid versus a combined group of invalid and sandbagging scorers. There were 3,299 (51.99%) athletes who had valid baseline scores, 269 (4.24%) had invalid scores, and 3,009 (47.42%) had sandbagging scores. (There were 231 who obtained both invalidity and sandbagging scores.) The overall difference in baseline-to-post concussion changes

between the valid scorers and the combined group of invalid and sandbagging scorers was statistically significant. The authors stated that the high rate of athletes who had invalid and sandbagging scores raised concern that the underperformance of baseline testing occurs more commonly than is probably realized by those who utilize computerized neuropsychological testing with high school athletes. Accordingly, efforts are needed to improve test administration procedures so that maximal attention and effort can be maintained among the test takers. According to the authors, increased caution is called for in employing the baseline-to-post concussion paradigm when return-to-play decisions are made.

Abeare et al. (2018) assessed the prevalence of invalid performance on baseline neurocognitive testing using embedded measures within computerized tests and individually administered neuropsychological measures and examined the influence of incentive status and performance validity on neuropsychological test scores. A total of 83 collegiate football athletes completing their preseason baseline assessment within the University's concussion management program and a control group of 140 non-athlete students were included in the study. The cross-sectional design of the study was based on differential incentive status: motivated to do poorly to return to play more quickly after sustaining a concussion (athletes) versus motivated to do well due to incentivizing performance (students). The main measures of the study included Immediate Post-Concussion and Cognitive Testing (ImPACT), performance validity tests, and measures of cognitive ability. Half of the athletes failed at least 1 embedded validity indicator within ImPACT (51.8%), and the traditional neuropsychological tests (49.4%), with large effects for performance validity on cognitive test scores, incentive status, and the combination of both factors on measures of attention and processing speed. The authors concluded that invalid performance on baseline assessment is common (50%), consistent across instruments (ImPACT or neuropsychological tests) and settings (one-on-one or group administration), increases as a function of incentive status (risk ratios: 1.3-4.0) and results in gross underestimates of the athletes' true ability level, complicating the clinical interpretation of the postinjury evaluation and potentially leading to premature return to play.

In a retrospective, cross-sectional study, Abeare et al. (2018) assessed the prevalence of invalid performance on baseline testing and assessed whether the prevalence varies as a function of age and validity indicator. Participants included 7897 consecutively tested equivalently proportioned male and female athletes aged 10 to 21 years, who completed baseline neurocognitive testing for the purpose of concussion management. Baseline assessment was conducted with the Immediate Post-concussion Assessment and Cognitive Testing (ImPACT). Base rates of failure on published ImPACT validity indicators were compared within and across age groups. Hypotheses were developed after data collection but prior to analyses. Of the 7897 study participants, 4086 (51.7%) were male, mean (SD) age was 14.71 years, 7820 (99.0%) were primarily English speaking, and the mean (SD) educational level was 8.79 years. The base rate of failure ranged from 6.4% to 47.6% across individual indicators. Most of the sample (55.7%) failed at least 1 of 4 validity indicators. The base rate of failure varied considerably across age groups (117 of 140 [83.6%] for those aged 10 years to 14 of 48 [29.2%] for those aged 21 years), representing a risk ratio of 2.86. The authors indicated that the results for base rate of failure were surprisingly high overall and varied widely depending on the specific validity indicator and the age of the examinee. The strong age association, with 3 of 4 participants aged 10 to 12 years failing validity indicators, suggests that the clinical interpretation and utility of baseline testing in this age group is questionable. According to the authors, these findings underscore the need for close scrutiny of performance validity indicators on baseline testing across age groups.

MacDonald and Duerson (2015) examined the test-retest reliability of a computerized neurocognitive test used for baseline assessments in high school athletes over 1 year. Study participants included high school athletes (n = 117) participating in American football or soccer. All study participants completed 2 baseline computerized neurocognitive tests taken 1 year apart at their respective schools. The test measures performance on 4 cognitive tasks: identification speed (Attention), detection speed (Processing Speed), one card learning accuracy (Learning), and one back speed (Working Memory). Reliability was assessed by measuring the intraclass correlation coefficient (ICC) between the repeated measures of the 4 cognitive tasks. Pearson and Spearman correlation coefficients were calculated as a secondary outcome measure. The measure for identification speed performed best and the measure for one card learning accuracy performed worst. All tests had marginal or low reliability. The authors concluded that in a population of high school athletes, computerized neurocognitive testing performed in a community setting demonstrated low to marginal test-retest reliability on baseline assessments 1 year apart. The authors stated that further investigation should focus on (1) improving the reliability of individual tasks tested, (2) controlling for external factors that might affect test performance, and (3) identifying the ideal time interval to repeat baseline testing in high school athletes. According to the authors, this study adds to the evidence that suggests in this population baseline testing may lack sufficient reliability to support clinical decision making.

Computerized Cognitive Testing such as Cognivue, Mindstreams BrainCare, and QbTest

Available clinical trials fail to document a beneficial effect of computerized cognitive testing on long-term clinical outcomes. The evidence is insufficient to establish the validity of computerized cognitive testing compared with traditional tests for the assessment of dementia and cognitive impairment.

In a systematic review and meta-analysis of diagnostic tests for the screening of mild cognitive impairment (MCI) and dementia, Chan et al. (2022) included 90 studies with 22, 567 participants to evaluate diagnostic performance among different types of digital drawing tests and paper-and-pencil drawing tests. Seventy-six of the included studies included participants with MCI or dementia in an outpatient clinic or from the community while the rest of the studies recruited participants in a hospital or long-term care setting. The digital drawing tests included in their review and analysis included the digital clock drawing test (CDT), digital pentagon drawing test, digital Rey-Osterrieth complex figure (ROCF), digital tree drawing test, digital house drawing test, and digital spiral test while the paper-and-pencil drawing tests included the CDT, pentagon drawing test, cube drawing test, and ROCF. Six of the studies used digital CDT and 80 of the studies used paper-and-pencil CDT. The primary outcome of the study was the diagnostic performance of the CDT for the screening of MCI and dementia and the secondary outcome was the diagnostic performance of the other types of drawing tests. The authors reported that the performances of the digital and paper-and-pencil pentagon drawing tests were comparable in the screening of dementia, but that the digital CDT demonstrated better sensitivity and specificity diagnostic performance than paper-and-pencil CDT for MCI. Other types of digital drawing tests showed comparable performance with paper-and-pencil formats. Limitations of this study include the lack of head-to-head comparisons, and that the number of studies to compare diagnostic performance of drawing tests are limited. The authors stated that the benefits of digital drawing tests may be stronger if there were more studies available for this meta-analysis.

Romero-Garcia et al. (2022) completed a single-center, prospective cohort study to assess if cognitive impairments would be apparent in a young and high functioning cohort and that app-based cognitive screening would complement traditional neuropsychological assessments. Their study included 17 patients with diffuse gliomas who completed a neuropsychological battery of tests that took 2-3 hours to complete and the OCS-BRIDGE assessment, an app-based touchscreen assessment that could be completed in 30 minutes. The traditional neuropsychological assessment was administered pre-operatively while the OCS-BRIDGE was administered pre- and post-operatively at the 3- and 12-month follow-ups. The authors reported that the traditional assessment showed that 79% of participants had an impairment in at least one domain, and an average of 2.88 cognitive impairments per participant before surgery, and that, after surgery, all but one participant had at least one impairment with a mean of 4.5 impairments per participants. The OCS-BRIDGE touchscreen assessment showed that 59% of participants had an impairment in at least one domain with a mean of 0.94 impairments per participant before surgery while longitudinal post-operative changes showed that 44% had a reduced number of impairments by their last assessment, 25% had the same, and 31% showed an increased number of impairments. Overall, the traditional neuropsychological tests detected 44 preoperative impairments among the 17 participants in the four combined domains of attention, memory, verbal skills, and non-verbal skills. OCS-BRIDGE detected 13 impairments and 28 possible impairments pre-operatively. The authors recognized that the traditional assessment using multiple items across the difficulty range proved more sensitive than the brief touchscreen assessment; however, they also noted that the capacity of the screening app to capture reaction times enhanced its sensitivity in the area of non-verbal function. The authors concluded that a combined approach, using traditional assessment in those areas where brief screening, may be less sensitive, and OCS-BRIDGE style measures for reaction time and perceptual tasks may be most effective and recommended robust, objective, and accessible assessment across multiple centers. Limitations of the study include the small sample size and single-center design, logistical and technical limitations to the assessments, heterogeneity of tumor location, size and treatment and the potential for practice effects due to reuse of the cognitive assessment tools.

A statistical analysis by Ye et al. (2022) was performed to evaluate BrainCheck, a computerized cognitive testing battery, for its diagnostic accuracy and ability to distinguish the severity of cognitive impairment. A total of 99 participants diagnosed with dementia, mild cognitive impairment (MCI), or normal cognition (NC) completed the BrainCheck battery. Statistical analyses compared participant performances on BrainCheck based on their diagnostic group. BrainCheck battery performance showed differences between the NC, MCI, and dementia groups, achieving 88% or higher sensitivity and specificity (i.e., true positive and true negative rates) for separating dementia from NC, and 77% or higher sensitivity and specificity in separating the MCI group from the NC and dementia groups. Three-group classification found true positive rates of 80% or higher for the NC and dementia groups and true positive rates of 64% or higher for the MCI group. The authors concluded that BrainCheck was able to distinguish between diagnoses of dementia, MCI, and NC, providing a potentially reliable tool for early detection of cognitive impairment. A small sample size makes it difficult to decide whether these conclusions can be generalized to a larger population. Further research with randomized controlled trials is needed to validate these findings.

Chan et al. (2021) performed a systematic review to evaluate the diagnostic performance of digital cognitive tests for mild cognitive impairment (MCI) and dementia in older adults. Literature searches were systematically performed in the OVID databases. Validation studies that reported the diagnostic performance of a digital cognitive test for MCI or dementia were included. The main outcome was the diagnostic performance of the digital test for the detection of MCI or dementia. A total of 56 studies with 46 digital cognitive tests were included in this study. Most of the digital cognitive tests were shown to have comparable diagnostic performances with the paper-and-pencil tests. Twenty-two digital cognitive tests showed a good diagnostic performance for dementia, with a sensitivity and a specificity over 0.80, such as the Computerized Visuo-Spatial Memory test and Self-Administered Tasks Uncovering Risk of Neurodegeneration. Eleven digital cognitive tests showed a good diagnostic performance for MCI such as the Brain Health Assessment. However, all the digital tests only had a few validation studies to verify their performance. The authors concluded that digital cognitive tests showed good performances for MCI and dementia, and that the digital test can collect digital data that is far beyond the traditional ways of cognitive tests. Further research with randomized controlled trials is needed to validate these findings.

An observational study by Rubin et al. (2021) was performed to determine the feasibility of implementing an iPad-based cognitive impairment screening tool, the psychometric properties of the tool, and predictors of cognitive impairment among adults seeking HIV care. A convenience sample of participants completed Brain Baseline Assessment of Cognition and Everyday Functioning (BRACE), which included (1) Trail Making Test Part A, measuring psychomotor speed; (2) Trail Making Test Part B, measuring set-shifting; (3) Stroop Color, measuring processing speed; and (4) the Visual-Spatial Learning Test. Global neuropsychological function was estimated as mean T score performance on the 4 outcomes. Impairment on each test or for the global mean was defined as a T score ≤ 40 . Subgroups of participants repeated the tests 4 weeks or > 6 months after completing the first test to evaluate intrapersonal test-retest reliability and practice effects (improvements in performance due to repeated test exposure). An additional subgroup completed a lengthier cognitive battery concurrently to assess validity. Relevant factors were abstracted from electronic medical records to examine predictors of global neuropsychological function. The study population consisted of 404 people with HIV (age: mean 53.6 years; race: 332/404, 82% Black; 34/404, 8% White, 10/404, 2% American Indian/Alaskan Native; 28/404, 7% other and 230/404, 58% male; 174/404, 42% female) of whom 99% (402/404) were on antiretroviral therapy. Participants completed BRACE in a mean of 12 minutes (SD 3.2), and impairment was demonstrated by 34% (136/404) on Trail Making Test A, 44% (177/404) on Trail Making Test B, 40% (161/404) on Stroop Color, and 17% (67/404) on Visual-Spatial Learning Test. Global impairment was demonstrated by 103 out of 404 (25%). Test-retest reliability for the subset of participants ($n = 26$) repeating the measure at 4 weeks was 0.81 and for the subset of participants ($n = 67$) repeating the measure almost 1 year later (days: median 294, IQR 50) was 0.63. There were no significant practice effects at either time point ($p = .20$ and $p = .68$, respectively). With respect for validity, the correlation between global impairment on the lengthier cognitive battery and BRACE was 0.63 ($n = 61$; $p < .001$), with 84% sensitivity and 94% specificity to impairment on the lengthier cognitive battery. The authors concluded that they were able to successfully implement BRACE and estimate cognitive impairment burden in the context of routine clinic care. BRACE was also shown to have good psychometric properties. This easy-to-use tool in clinical settings may facilitate the care needs of people with HIV as cognitive impairment continues to remain a concern in people with HIV. Further research with randomized controlled trials is needed to validate these findings.

Wilson et al. (2021) performed a systematic review of literature to evaluate the benefits, limitations, and validity of computerized neuropsychological assessment devices (CNADs) in the evaluation of HIV-associated neurocognitive disorder (HAND). Following a comprehensive search, the abstracts of relevant articles were compiled and then reviewed for the use of digital neuropsychological testing in the setting of HIV. The articles that met these criteria were read, and their reference lists further examined to compile a more inclusive review. The review was limited to peer-reviewed English-language journals published within the past 20 years, with no other restrictions, such as sample size or analysis type. Eight CNADs that have undergone validity testing in the setting of HIV were identified and included in the review. The studies included CNADs modeled after traditional testing batteries as well as non-traditional cognitive batteries with advanced technology features including simulated or virtual realities and quick, daily mobile phone assessments, which were reviewed. This review suggests that these computerized neuropsychological assessment devices remain in the early stages of development. The authors concluded that these digital batteries do not have the ability to supplant gold standard neuropsychological tests in screening for HAND. However, many have the potential to become effective clinical screening tools. This review reveals most of these validity studies do not employ large enough sample sizes (fewer than 100) to conclusively determine their ability to detect HAND, creating a degree of uncertainty in external validity. A small sample size makes it difficult to decide whether these conclusions can be generalized to a larger population. The findings of this study need to be validated by well-designed studies. Further investigation is needed before clinical usefulness of this procedure is proven.

In a cohort of 114 patients presenting to an attention deficit hyperactivity disorder (ADHD) outpatient clinic, Brunkhorst-Kanaan et al. (2020) investigated how well a commercially available continuous performance test (CPT) (QbTest) can differentiate between patients with ADHD (n = 94) and patients with a disconfirmed ADHD diagnosis (n = 20). Both groups showed numerous comorbidities, predominantly depression (27.2% in the ADHD group vs. 45% in the non-ADHD group) and substance-use disorders (18.1% vs. 10%, respectively). Patients with ADHD showed significant higher activity (2.07 ± 1.23) than patients without ADHD (1.34 ± 1.27 , $dF = 112$; $p = 0.019$), whereas for the other core parameters, inattention, and impulsivity no differences could be found. Reaction time variability has been discussed as a typical marker for inattention in ADHD. Therefore, the authors investigated how well ex-Gaussian analysis of response time can differentiate between ADHD and other patients, showing, that it does not help to identify patients with ADHD. Even though patients with ADHD showed significantly higher activity, this parameter differed only poorly between patients. The authors concluded that CPTs do not help to identify patients with ADHD in a specialized outpatient clinic. According to the authors, the usability of this test for differentiating between ADHD and other psychiatric disorders is poor and a sophisticated analysis of reaction time did not decisively increase the test accuracy.

Cahn-Hidalgo et al. (2020) determined the cut-off scores for classification of cognitive impairment and assessed Cognivue safety and efficacy in a validation study. Adults (age 55-95 years) at risk for age-related cognitive decline or dementia were invited via posters and email to participate in two cohort studies conducted at various outpatient clinics and assisted- and independent-living facilities. In the cut-off score determination study (n = 92), optimization analyses by positive percent agreement (PPA) and negative percent agreement (NPA), and by accuracy and error bias were conducted. In the clinical validation study (n = 401), regression, rank linear regression, and factor analyses were conducted. Participants in the clinical validation study also completed other neuropsychological tests. For the cut-off score determination study, 92 participants completed St. Louis University Mental Status (SLUMS, reference standard) and Cognivue tests. Analyses showed that SLUMS cut-off scores of < 21 (impairment) and > 26 (no impairment) corresponded to Cognivue scores of 54.5 (NPA = 0.92; PPA = 0.64) and 78.5 (NPA = 0.5; PPA = 0.79), respectively. Therefore, conservatively, Cognivue scores of 55-64 corresponded to impairment, and 74-79 to no impairment. For the clinical validation study, 401 participants completed ≥ 1 testing session, and 358 completed 2 sessions 1-2 wk apart. Cognivue classification scores were validated, demonstrating good agreement with SLUMS scores (weighted κ 0.57; 95%CI: 0.50-0.63). Reliability analyses showed similar scores across repeated testing for Cognivue ($R^2 = 0.81$; $r = 0.90$) and SLUMS ($R^2 = 0.67$; $r = 0.82$). Psychometric validity of Cognivue was demonstrated vs. traditional neuropsychological tests. Scores were most closely correlated with measures of verbal processing, manual dexterity/speed, visual contrast sensitivity, visuospatial/executive function, and speed/sequencing. The investigators concluded that Cognivue scores ≤ 50 avoid misclassification of impairment, and scores ≥ 75 avoid misclassification of unimpairment. According to the investigators, this validation study demonstrates good agreement between Cognivue and SLUMS; superior reliability; and good psychometric validity. A limitation of these studies is the use of a single reference standard, SLUMS. Longitudinal follow-up studies are needed to evaluate the ability of Cognivue to monitor cognitive deterioration over time.

Groppell et al., (2019) determined the accuracy and validity of BrainCheck Memory as a diagnostic aid for age-related cognitive impairment, as compared against physician diagnosis and other commonly used neurocognitive screening tests, including the Saint Louis University Mental Status (SLUMS) exam, the Mini-Mental State Examination (MMSE), and the Montreal Cognitive Assessment (MoCA). A total of 583 volunteers over the age of 49 were tested from various community centers and living facilities. The volunteers were divided into five cohorts: a normative population and four comparison groups for the SLUMS exam, the MMSE, the MoCA, and physician diagnosis. Each comparison group completed their respective assessment and BrainCheck Memory. A total of 398 subjects were included in the normative population. A total of 84 participants were in the SLUMS exam cohort, 51 in the MMSE cohort, 35 in the MoCA cohort, and 18 in the physician cohort. BrainCheck Memory assessments were significantly correlated to the SLUMS exam, with coefficients ranging from .5 to .7. Correlation coefficients for the MMSE and BrainCheck and the MoCA and BrainCheck were also significant. Of the 18 subjects evaluated by a physician, 9 (50%) were healthy, 6 (33%) were moderately impaired, and 3 (17%) were severely impaired. A significant difference was found between the severely and moderately impaired subjects and the healthy subjects ($p = .02$). The investigators found that the BrainCheck Memory composite score showed stronger correlations with the standard assessments as compared to the individual BrainCheck assessments. Receiver operating characteristic (ROC) curve analysis of this composite score found a sensitivity of 81% and a specificity of 94%. The investigators concluded that BrainCheck Memory provides a sensitive and specific metric for age-related cognitive impairment in older adults, with the advantages of a mobile, digital, and easy-to-use test. According to the authors, some participants were unable to complete BrainCheck's entire battery of assessments. While this was accounted for during the analysis, the missing data may have limited statistical power. The

investigators also indicated that due to the study's small sample size, more research is needed to compare and validate BrainCheck against physician diagnosis.

Hollis et al. (2018) evaluated the impact of providing a computerized test of attention and activity (QbTest) report on the speed and accuracy of diagnostic decision-making in children with suspected ADHD. A randomized, parallel, single-blind controlled trial in mental health and community pediatric clinics in England was conducted. Participants were 6-17 years-old and referred for ADHD diagnostic assessment; all underwent assessment-as-usual, plus QbTest. Participants and their clinician were randomized to either receive the QbTest report immediately (QbOpen group), or the report was withheld (QbBlind group). The primary outcome was number of consultations until a diagnostic decision confirming/excluding ADHD within 6-months from baseline. One hundred and thirty-two participants were randomized to QbOpen group (123 analyzed) and 135 to QbBlind group (127 analyzed). Clinicians with access to the QbTest report (QbOpen) were more likely to reach a diagnostic decision about ADHD (hazard ratio 1.44, 95% CI 1.04-2.01). At 6-months, 76% of those with a QbTest report had received a diagnostic decision, compared with 50% without. QbTest reduced appointment length by 15% (time ratio 0.85, 95% CI 0.77-0.93), increased clinicians' confidence in their diagnostic decisions (odds ratio 1.77, 95% CI 1.09-2.89) and doubled the likelihood of excluding ADHD. There was no difference in diagnostic accuracy. The authors concluded that the QbTest may increase the efficiency of ADHD assessment pathway allowing greater patient throughput with clinicians reaching diagnostic decisions faster without compromising diagnostic accuracy. Limitations of the study include that follow-up was limited to a six-month time period. Given that overall, almost one third of participants had still not received a diagnostic decision after six months, it was not possible to determine the impact of QbTest on the eventual diagnosis of those participants still awaiting a diagnostic decision at the end of the study. Additionally, the study was not powered enough to assess possible age effects.

Aslam et al. (2018) conducted a systematic review to determine whether automated computerized tests accurately identify patients with progressive cognitive impairment and, if so, to investigate their role in monitoring disease progression and/or response to treatment. Six electronic databases were searched from January 2005 to August 2015 to identify papers for inclusion. Studies assessing the diagnostic accuracy of automated computerized tests for mild cognitive impairment (MCI) and early dementia against a reference standard were included. Where possible, sensitivity, specificity, positive predictive value, negative predictive value, and likelihood ratios were calculated. The Quality Assessment of Diagnostic Accuracy Studies tool was used to assess risk of bias. Sixteen studies assessing 11 diagnostic tools for MCI and early dementia were included. No studies were eligible for inclusion in the review of tools for monitoring progressive disease and response to treatment. The overall quality of the studies was good. However, the wide range of tests assessed, and the non-standardized reporting of diagnostic accuracy outcomes meant that statistical analysis was not possible. The authors concluded that some tests have shown promising results for identifying MCI and early dementia. However, concerns over small sample sizes, lack of replicability of studies, and lack of evidence available make it difficult to make recommendations on the clinical use of the computerized tests for diagnosing, monitoring progression, and treatment response for MCI and early dementia.

Racine et al. (2016) conducted a study that included 469 late middle-aged participants from the Wisconsin Registry for Alzheimer's Prevention (mean age 63.8 ± 7 years at testing; 67% female; 39% APOE4+) to evaluate whether computerized cognitive assessments, like the CogState battery, are sensitive to preclinical cognitive changes or pathology in people at risk for Alzheimer's disease (AD). The study examined relationships between a CogState abbreviated battery (CAB) of seven tests and demographic characteristics, traditional paper-based neuropsychological tests as well as a composite cognitive impairment index, cognitive impairment status (determined by consensus review), and biomarkers for amyloid and tau (CSF phosphorylated-tau/A β 42 and global PET-PiB burden) and neural injury (CSF neurofilament light protein). CSF and PET-PiB were collected in n = 71 and n = 91 participants, respectively, approximately four years prior to CAB testing. For comparison, three traditional tests of delayed memory in parallel were examined. Similar to studies in older samples, the CAB was less influenced by demographic factors than traditional tests. CAB tests were generally correlated with most paper-based cognitive tests examined and mapped onto the same cognitive domains. Greater composite cognitive impairment index was associated with worse performance on all CAB tests. Cognitively impaired participants performed significantly worse compared to normal controls on all but one CAB test. Poorer One Card Learning test performance was associated with higher levels of CSF phosphorylated-tau/A β 42. The authors concluded that these results support the use of the CogState battery as measures of early cognitive impairment in studies of people at risk for Alzheimer's disease. However, according to the authors, the study also suggests that CogState at a single time point may not substantially improve preclinical AD detection over traditional neuropsychological tests.

Shopin et al. (2013) compared a computerized battery of neuropsychological tests for memory, attention, and executive functions (MindStreams) with the Montreal Cognitive Assessment (MoCA) to detect mild-to-moderate cognitive impairments in poststroke patients. A total of 454 patients with transient ischemic attack (TIA) or stroke enrolled to the TABASCO (Tel Aviv

Brain Acute Stroke Cohort) study, a prospective study which includes consecutive first-ever mild-to-moderate stroke patients, were included. All participants underwent neurological and cognitive evaluations. The patients' mean MoCA and MindStreams scores were lower than normal; however, the TIA group presented significantly better scores using either method. The correlation between the MoCA and the computerized global score was 0.6. A significant correlation was found between the subcategory scores (executive function, memory, and attention). However, the MoCA identified many more subjects with low scores (< 26) compared to the MindStreams (70.6 vs. 15.7%).

Intermittent Explosive Disorder

There is insufficient clinical evidence to demonstrate that the use of neuropsychological testing for individuals with intermittent explosive disorder without associated cognitive disorders can be used effectively for clinical decision making to improve patient management of this condition.

There are no clear underlying medical issues associated with intermittent explosive disorder, nor are there published clinical trials that support the use of neuropsychological testing for this disorder. According to the Diagnostic and Statistical Manual of Mental Disorders (DSM-5), published by the American Psychiatric Association, the following criteria must be met in order for a patient to be diagnosed with intermittent explosive disorder:

- Recurrent behavioral outbursts that represent a failure to control aggressive impulses as manifested by one of the following:
 - Verbal aggression (e.g., temper tantrums, tirades, verbal arguments, or fights) or physical aggression towards property, animals, or other individuals, occurring, on average, twice weekly for a period of three months. The physical aggression does not result in damage or destruction of property and does not result in physical injury to animals or other individuals
 - Three behavioral outbursts involving damage or destruction of property and/or physical assault with physical injury against animals or other individuals occurring within a 12-month period
- The magnitude of aggressiveness expressed during the recurrent outbursts is grossly out of proportion to the provocation or any precipitating psychosocial stressors
- The recurrent aggressive outbursts are not premeditated (i.e., are impulsive) and are not committed to achieve some tangible objective (e.g., money, power, intimidation)
- The recurrent aggressive outbursts cause either marked distress in the individual or impairment in occupational or interpersonal functioning or are associated with financial or legal consequences
- Chronological age is at least 6 years (or equivalent developmental level)
- The recurrent aggressive outbursts are not better explained by another mental disorder and are not attributable to another medical condition or to physiological effects of a substance

Headaches Including Migraine

There is insufficient clinical evidence to demonstrate that the use of neuropsychological testing for individuals with migraine or other headaches without associated cognitive disorders can be used effectively for clinical decision making to improve patient management of this condition.

Lozano-Soto et al. (2023) conducted a case-control study to examine the presence of neuropsychological deficits in chronic migraine (CM) patients during the interictal phase. The study included 39 CM patients recruited from a single outpatient center and 20 age-, sex-, and education-matched healthy controls (HCs). All study participants underwent clinical, neuropsychiatric, and neuropsychological evaluation by a clinical neurologist to evaluate cognitive domains, including sustained attention (SA), information processing speed (IPS), visuospatial episodic memory, working memory (WM), and verbal fluency (VF), as well as depressive and anxiety symptoms. CM patients exhibited higher scores than HCs for all clinical and neuropsychiatric measures, but no differences were found in personality characteristics. The authors reported that more than half of the CM patients (54%) showed mild-to-severe neuropsychological impairment (NI) with 35.9% classified as having mild NI, 12.8% with moderate NI and 5.1% with severe NI. Additional exploratory analysis showed that more than half (54%) of CM patients with mild, moderate, or severe NI took two or more than two medications and that the severity of NI was associated with the number of treatments received. The authors reported that CM patients exhibited variable NI during periods between acute migraine attacks and that the patients demonstrated cognitive impairment in SA, verbal episodic memory, and Stroop-like interference. Limitations of the study included the small sample size, the single center design, and the large variety of the treatments that the patients received. The authors concluded that CM can be accompanied by a variety of cognitive symptoms during the interictal phase and that these cognitive impairments were most likely related to the mechanisms underlying migraine-induced disability.

In another study that investigated the cognitive impairment of migraineurs, Qin et al. (2022) enrolled 117 adult patients with primary headaches, including 87 with migraine, 30 with tension-type headache (TTH) and 30 healthy controls. No significant differences were found in age, sex, or years of education among the three groups. The authors reported that the Montreal Cognitive Assessment (MoCA) total score and the scores of visuospatial and executive functions, language, and delayed recall in the migraine and TTH groups were significantly lower than those in the healthy control group (all $p < 0.05$) while no significant differences were observed in naming, attention, abstraction, and orientation between the patients and healthy controls. Limitations of the study included the questionnaire and scales to assess the study subjects, the risk of recall bias in the evaluation of the subjects' anamnestic description of migraine history, the small sample size, the short-term follow-up period the lack of supplementary examinations and the lack of specific information regarding antimigraine medication use and type. The authors concluded the study confirmed cognitive impairment in patients with migraine and TTH and that the duration of attack had an effect on cognitive function in migraineurs.

A cross-sectional study by Chen et al. (2021) was performed to assess whether patients with migraine without aura (MwoA) during the interictal period have attention impairment and to identify the migraine characteristics related to attention deficits. Forty-four subjects with MwoA (4 males, 40 females) and 20 controls matched for age, gender, and literacy education were included in the study. The attention network test (ANT) and a battery of neuropsychological tests, including the trail-making test (TMT), the digit span test (DST), and the Stroop test, were administered to the participants during the headache-free period. Patients in MwoA were more anxious ($p = 0.007$) and depressed ($p = 0.001$) than healthy subjects. Significant differences between the two groups were detected in the executive network ($p = 0.006$) but not in the alerting and orienting networks of ANT. Mean reaction time of ANT in the MwoA group was significantly longer than that in the control group ($p = 0.028$). Patients showed worse performance on DST-forward ($p < 0.001$), DST-backward ($p < 0.001$), DS Total ($p < 0.001$), TMT-A ($p < 0.001$), TMT-B ($p < 0.001$) and TMT-d ($p = 0.002$). Differences found in executive functions between the two groups were unrelated to gender, age, literacy, anxiety, and depression. Multiple regression analysis revealed no relation between clinical characteristics of headache and scores on the executive function with MwoA. The authors concluded that the study suggested that patients in MwoA present worse performances on the executive control of attention networks during the headache-free period, which appear not be associated with measures of migraine severity. The authors also stated although more studies are needed in this area, the results could be useful to find a specific neuropsychological biomarker for migraine pathophysiology. A small sample size makes it difficult to decide whether these conclusions can be generalized to a larger population. Further research with randomized controlled trials is needed to validate these findings.

Foti et al. (2017) identified 16 studies evaluating the association between migraine and cognitive impairment. The authors found that these studies demonstrated conflicting results. Some studies show a detrimental effect of migraine on cognitive skills and other studies have shown no difference in cognitive skills for patients with migraine headaches.

Dresler et al. (2012) evaluated three neuropsychological tests (Trail Making Test (TMT), Go/Nogo Task and Stroop Task) that were completed by four headache patient samples (chronic CH, episodic CH in the active or inactive period, and migraine patients) and compared to healthy controls. Analyses revealed that patients with chronic and active episodic CH appeared particularly impaired in tests relying more on intact executive functioning (EF) than on basal cognitive processes. Within the CH groups performance decreased linearly with increasing severity. The authors stated that impaired EF could also result from medication and sleep disturbances due to active CH. The authors went on to say that because decreased performance was also present outside the attacks it may hint at generally altered brain function but does not necessarily reflect clinically relevant behavior.

History of Myocardial Infarction

There is insufficient clinical evidence to demonstrate that the use of neuropsychological testing for individuals with a history of myocardial infarction without associated cognitive disorders can be used effectively for clinical decision making to improve patient management of this condition.

Studies on the relationship between myocardial infarction and cognitive functioning have demonstrated conflicting results. Some studies show a detrimental effect of myocardial infarction on cognitive skills (Gallagher et al., 2023; Sauv e et al., 2009; Almeida et al., 2008). Other studies have shown no difference in cognitive skills for patients with myocardial infarctions (Ahto et al., 1999, Grubb et al., 2000).

In a systematic review, Cameron et al. (2016) evaluated the diagnostic accuracy of cognitive screening instruments in screening for mild cognitive impairment (MCI) in heart failure (HF) patients. Inclusion criteria for the review were as follows: primary

studies examining cognitive impairment in HF, administration of a cognitive screening instrument and neuropsychological test battery, and cognitive impairment indicated by performance on neuropsychological tests 1.5 SDs less than that of normative data. The precision, accuracy, and receiver operating characteristic curves of the Mini Mental State Examination were computed. From 593 citations identified, 8 publications met inclusion criteria. Risk of bias included selective HF patient samples, and no study examined the diagnostic test accuracy of the cognitive screening instruments. The Mini Mental State Examination had low sensitivity (26%) and high specificity (95%) with a score of 28 or less as the optimal threshold for MCI screening. The authors concluded that screening for cognitive impairment in HF is recommended; however, future studies need to establish the diagnostic accuracy of screening instruments of MCI in this population.

Self-Administered or Self-Scored Neuropsychological Testing

Oliva et al. (2022) performed a validation study of the NAIHA Neuro Cognitive Test (NNCT), a computerized, self-administered neuropsychological screening test designed for elderly people with and without cognitive impairment via digitized cognitive assessments. The study included 147 adults over 65 years of age. The authors reported that the validity of the NNCT was demonstrated by correlating outcomes from the Mini Mental State Exam (MMSE), the Clock Drawing Test (CDT) and the Cambridge Cognitive Examination-Revised (CAMCOG-R) test as all subscales of the NNCT test correlated significantly and positively with some of these tests. The authors reported that the NNCT also discriminated correctly to assign the participants into the three groups, Healthy Older Adults (HOA; $n = 70$), mild cognitive impairment (MCI; $n = 44$) and Alzheimer's dementia (AD; $n = 33$) and that the test can be used for screening and for diagnostic support. Limitations of the study include the small sample size, the single center design, and the inability of some participants to complete all phases of the study due to poor vision and limited ability to use tablet devices.

A cross-sectional study by Paterson et al. (2022) was performed to validate the online Brain Health Assessment (BHA) for detection of amnesic mild cognitive impairment (aMCI) compared to gold-standard neuropsychological assessment, the Montreal Cognitive Assessment (MoCA). Using a cross-sectional design, community-dwelling older adults completed a neuropsychological assessment, were diagnosed as normal cognition (NC) or aMCI and completed the BHA and MoCA. Both logistic regression (LR) and penalized logistic regression (PLR) analyses determined BHA and demographic variables predicting aMCI; MoCA variables were similarly modeled. Diagnostic accuracy was compared using area under the receiver operating characteristic curve (ROC AUC) analyses. Ninety-one participants met inclusion criteria (51 aMCI, 40 NC). PLR modeling for the BHA indicated Face- Name Association, Spatial Working Memory, and age-predicted aMCI (ROC AUC = 0.76; 95% confidence interval [CI]: 0.66–0.86). Optimal cut-points resulted in 21% classified as aMCI (positive), 23% negative, and 56% inconclusive. For the MoCA, digits, abstraction, delayed recall, orientation, and age predicted aMCI (ROC AUC = 0.71; 95% CI: 0.61–0.82). Optimal cut-points resulted in 22% classified positive, 8% negative, and 70% inconclusive (LR results presented within). The BHA model classified fewer participants into the inconclusive category and more as negative for aMCI, compared to the MoCA model (Stuart–Maxwell $p = .004$). The authors concluded that self-administered BHA provides similar detection of aMCI as a clinician-administered screener (MoCA), with fewer participants classified inconclusively. The BHA has the potential to save practitioners time and decrease unnecessary referrals for a comprehensive assessment to determine the presence of aMCI. Further research with randomized controlled trials (RCTs) is needed to validate these findings.

A cross-sectional study by Vyshedskiy et al. (2022) was performed to assess test scores and the correlation between Boston Cognitive Assessment (BOCA) and Montreal Cognitive Assessment (MoCA) test scores. BOCA is a self-administered 10-minute at-home test intended for longitudinal cognitive monitoring, and MoCA, a gold standard pen-and-paper test of global cognition. BOCA uses randomly selected non-repeating tasks to minimize practice effects. BOCA evaluates eight cognitive domains: 1) Memory/Immediate Recall, 2) Combinatorial Language Comprehension/Prefrontal Synthesis, 3) Visuospatial Reasoning/Mental rotation, 4) Executive function/Clock Test, 5) Attention, 6) Mental math, 7) Orientation, and 8) Memory/Delayed Recall. A total of 100 patients were included in the study. BOCA was administered to patients with cognitive impairment ($n = 50$) and age- and education-matched controls ($n = 50$). Test scores were significantly different between patients and controls ($p < 0.001$) suggesting good discriminative ability. The Cronbach's alpha was 0.87 implying good internal consistency. BOCA demonstrated strong correlation with MoCA ($R = 0.90$, $p < .0001$). The study revealed strong ($R = 0.94$, $p < 0.001$) test-retest reliability of the total BOCA score one week after participants' initial administration. The practice effect tested by daily BOCA administration over 10 days was insignificant ($\beta = 0.03$, $p = 0.68$). The effect of the screen size tested by BOCA administration on a large computer screen and re-administration of the BOCA to the same participant on a smartphone was insignificant ($\beta = 0.82$, $p = 0.17$; positive β indicates greater score on a smartphone). The authors concluded that BOCA has the potential to reduce the cost and improve the quality of longitudinal cognitive tracking essential for testing novel interventions designed to reduce or reverse cognitive aging. The authors also state that additionally, the test can be used to assess the effect

of anesthesia, long-term effect of cancer drugs, COVID fog, and other conditions known to affect cognition. Further research with randomized controlled trials is needed to validate these findings.

A randomized clinical trial was completed by Mahncke et al. (2021) to evaluate the efficacy of self-administered computerized cognitive training. A multisite randomized double-blind clinical trial of a behavioral intervention with an active control was conducted from September 2013 to February 2017 including assessments at baseline, post-training, and after a 3-month follow-up period. The goal of this study was to evaluate the efficacy of a self-administered computerized plasticity-based cognitive training programs in primarily military/veteran participants with a history of mild traumatic brain injury (mTBI) and cognitive impairment. Participants self-administered cognitive training (experimental and active control) programs at home, remotely supervised by a healthcare coach, with an intended training schedule of 5 days per week, 1 hour per day, for 13 weeks. Participants (149 contacted, 83 intent-to-treat) were confirmed to have a history of mTBI (mean of 7.2 years post-injury) through medical history/clinician interview and persistent cognitive impairment through neuropsychological testing and/or quantitative participant reported measure. The experimental intervention was a brain plasticity-based computerized cognitive training program targeting speed/accuracy of information processing, and the active control was composed of computer games. The primary cognitive function measure was a composite of nine standardized neuropsychological assessments, and the primary directly observed functional measure a timed instrumental activities of daily living assessment. Secondary outcome measures included participant-reported assessments of cognitive and mental health. The treatment group showed an improvement in the composite cognitive measure larger than that of the active control group at both the post-training [+ 6.9 points, confidence interval (CI) + 1.0 to + 12.7, $p = 0.025$, $d = 0.555$] and the follow-up visit (+ 7.4 points, CI + 0.6 to + 14.3, $p = 0.039$, $d = 0.591$). Both large and small cognitive function improvements were seen twice as frequently in the treatment group than in the active control group. No between-group effects were seen on other measures, including the directly observed functional and symptom measures. Statistically equivalent improvements in both groups were seen in depressive and cognitive symptoms. Further investigation is needed before clinical usefulness of this procedure is proven.

A cohort study by Scharre et al. (2021) was performed to compare longitudinal Self-Administered Gerocognitive Examination (SAGE) test scores to non-self-administered Mini-Mental State Examination (MMSE) scores in 5 different diagnostic subgroups. A cohort study evaluating annual rates of change was performed on 665 consecutive patients from Ohio State University Memory Disorders Clinic. Patients with at least two visits 6 months apart evaluated with SAGE and MMSE and classified according to standard clinical criteria as subjective cognitive decline (SCD), mild cognitive impairment (MCI), or Alzheimer's disease (AD) dementia were included. The pattern of change in SAGE scores was compared to MMSE. One way and repeated measures ANOVA and linear regression models were used. Four hundred twenty-four individuals (40 SCD, 94 MCI non-converters to dementia, 70 MCI converters to dementia (49 to AD dementia and 21 to non-AD dementia), 220 AD dementia) met inclusion criteria. SAGE and MMSE scores declined respectively at annual rates of 1.91 points/year ($p < 0.0001$) and 1.68 points/year ($p < 0.0001$) for MCI converters to AD dementia, and 1.82 points/year ($p < 0.0001$) and 2.38 points/year ($p < 0.0001$) for AD dementia subjects. SAGE and MMSE scores remained stable for SCD and MCI non-converters. Statistical decline from baseline scores in SAGE occurred at least 6 months earlier than MMSE for MCI converters to AD dementia (14.4 vs. 20.4 months), MCI converters to non-AD dementia (14.4 vs. 32.9 months), and AD dementia individuals (8.3 vs. 14.4 months). The authors concluded that SAGE detects MCI conversion to dementia at least 6 months sooner than MMSE. Being self-administered, SAGE also addresses a critical need of removing some barriers in performing cognitive assessments. Limitations of this study include potential referral and sampling biases. Repetitively administering SAGE and identifying stability or decline may provide clinicians with an objective cognitive biomarker impacting evaluation and management choices. Further research with randomized controlled trials is needed to validate these findings.

Clinical Practice Guidelines

American Academy of Child and Adolescent Psychiatry (AACAP)

Practice parameters from the American Academy of Child and Adolescent Psychiatry (Volkmar et al., 2014) state that neuropsychological correlates of autism spectrum disorder include impairments in executive functioning (e.g., simultaneously engaging in multiple tasks) (Ozonoff et al., 1991), weak central coherence (integrating information into meaningful wholes) (Happé and Frith 2006), and deficits in theory-of-mind tasks (taking the perspective of another person) (Baron-Cohen et al., 1985).

American Academy of Clinical Neuropsychology (AACN) and National Academy of Neuropsychology (NAN)

A joint position paper of the AACN and NAN sets forth their position on appropriate standards and conventions for computerized neuropsychological assessment devices (CNADs). The authors state that CNADs are subject to, and should meet, the same standards for the development and use of educational, psychological, and neuropsychological tests (American Psychological Association, 1999) as are applied to examiner-administered tests. The authors also state that those employing CNADs have the education, training, and experience necessary to interpret their results in a manner that will best meet the needs of the patients they serve (Bauer et al., 2012).

American Academy of Neurology (AAN)

In an evidence-based guideline update for the evaluation and management of concussion in sports, the AAN states that it is likely that neuropsychological testing of memory performance, reaction time, and speed of cognitive processing, regardless of whether administered by paper-and-pencil or computerized method, is useful in identifying the presence of concussion (sensitivity 71%- 88% of athletes with concussion). This is based on evidence from 1 Class II study and multiple Class III studies. The AAN also states that both types of testing (paper-and-pencil or computerized) generally require a neuropsychologist for accurate interpretation, although the tests may be administered by a non-neuropsychologist. According to AAN, there is insufficient evidence to support conclusions about the use of neuropsychological testing in identifying concussion in preadolescent age groups. The AAN goes on to say that inexperienced licensed health care providers (LHCPs) should be instructed in the proper administration of standardized validated sideline assessment tools. This instruction should emphasize that these tools are only an adjunct to the evaluation of the athlete with suspected concussion and cannot be used alone to diagnose concussion (Level B- probably effective). The AAN further states that LHCPs caring for athletes might utilize individual baseline scores on concussion assessment tools, especially in younger athletes, those with prior concussions, or those with preexisting learning disabilities/attention deficit/hyperactivity disorder, as doing so fosters better interpretation of postinjury scores (Level C- Possibly effective) (Giza et al., 2013, reaffirmed 2022).

A practice guideline update for disorders of consciousness (DoC) developed by the American Academy of Neurology; the American Congress of Rehabilitation Medicine; and the National Institute on Disability, Independent Living, and Rehabilitation Research indicates that clinicians should use standardized neurobehavioral assessment measures that have been shown to be valid and reliable to improve diagnostic accuracy for the purpose intended in patients with DoC (Giacino et al. 2018).

In a practice guideline update summary for mild cognitive impairment (MCI), the AAN recommends that when performing a Medicare Annual Wellness Visit, clinicians should not rely on historical report of subjective memory concerns alone when assessing for cognitive impairment (Level B). Various assessment instruments have acceptable diagnostic accuracy for detecting MCI, with no instrument being superior to another. The guideline states that because brief cognitive assessment instruments are usually calibrated to maximize sensitivity rather than specificity, patients who test positive for MCI should then have further assessment (e.g., more in-depth cognitive testing, such as neuropsychological testing with interpretation based on appropriate normative data) to formally assess for this diagnosis (Petersen et al., 2018; reaffirmed 2021).

A practice parameter for the screening and diagnosis of autism developed by the American Academy of Neurology and the Child Neurology Society indicates that neuropsychological, behavioral, and academic assessments should be performed as needed, in addition to the cognitive assessment, to include social skills and relationships, educational functioning, problematic behaviors, learning style, motivation and reinforcement, sensory functioning, and self-regulation for the diagnosis of autism (Filipek et al., 2000; Reaffirmed on October 18, 2003, July 28, 2006, July 10, 2010, and August 9, 2014).

American Academy of Pediatrics (AAP)

A joint statement for learning disabilities, dyslexia, and vision from the American Academy of Pediatrics, Section on Ophthalmology, Council on Children with Disabilities; American Academy of Ophthalmology; American Association for Pediatric Ophthalmology and Strabismus; and the American Association of Certified Orthoptists states that children who exhibit signs of learning disabilities should be referred for educational, psychological, neuropsychological, and/or medical diagnostic assessments (AAP, 2009; Reaffirmed 2014).

In 2018, the AAP updated the clinical report guidance for sport-related concussion (SRC) in children and adolescents. The authors of the report indicate that there are numerous studies evaluating the reliability of various computerized neurocognitive tests (CNTs) platforms; however, studies conducted independently of the developers of the tests have questioned the overall

reliability of testing from year to year. The reliability of pencil and paper testing has also been questioned. The authors indicate that ideally, neurocognitive testing is performed and interpreted by a neuropsychologist. However, given the large number of athletes participating in sports and the relative scarcity of and limited access to neuropsychologists, a widespread CNT program would not be practical or possible. If a non-neuropsychologist is using CNTs, collaboration with a neuropsychologist to aid in test administration and interpretation may be beneficial. CNTs or baseline testing is not specifically addressed in the conclusion or recommendation sections of the report (Halstead et al., 2018).

American Heart Association and the American Stroke Association

In a guideline for Healthcare Professionals from the American Heart Association and the American Stroke Association, Winstein et al. (2016) provided a synopsis of best clinical practices in the rehabilitative care of adults recovering from stroke. According to the guideline, a formal neuropsychological examination (including assessment of language, neglect, praxis, memory, emotional responses, and specific cognitive syndromes) may be helpful after the detection of cognitive impairment with a screening instrument. Neuropsychological protocols must be sensitive to a wide range of abilities, especially the assessment of executive and attentional functions. These guidelines state that screening for cognitive deficits is recommended for all stroke patients before discharge home (class I, level B evidence). The guidelines also indicate that when screening reveals cognitive deficits, a more detailed neuropsychological evaluation to identify areas of cognitive strength and weakness may be beneficial (class I/a, level C evidence).

American Medical Society for Sports Medicine

The American Medical Society for Sports Medicine position statement on concussion in sport states that baseline testing may be useful in some cases but is not necessary, required or an accepted standard of care for the appropriate management of sport-related concussion (Harmon et al., 2019).

American Psychiatric Association

In its guidelines on the treatment of Alzheimer's disease and other dementias, the American Psychiatric Association states that neuropsychological testing may be helpful in deciding whether a patient with subtle or atypical symptoms actually has dementia. Neuropsychological testing is particularly useful in the evaluation of individuals who present with mild cognitive impairment, which requires evidence of memory and/or other cognitive difficulties in the presence of intact functioning, and in the evaluation of individuals with the onset of dementia early in life. Testing may help to characterize the extent of cognitive impairment, to distinguish among the types of dementias, and to establish baseline cognitive function. The American Psychiatric Association also states that a variety of research definitions for mild cognitive impairment are in place, but there is no consensus on the optimal definition. The most widely accepted definition (American Psychiatric Association, 2007) requires the following:

- Subjective cognitive complaints
- Evidence of objective deficits in cognitive function based on age- and education-adjusted norms on standardized neuropsychological tests
- Intact daily functioning
- Evidence of cognitive decline from a prior level, and
- Evidence of not meeting the criteria for dementia

American Psychological Association (APA)

The American Psychological Association published a Psychological and Neuropsychological Testing Billing and Coding Guide (American Psychological Association, 2019). The guide states that neuropsychological testing is considered medically necessary where initial assessment or assessment over time is needed to:

- Measure cognitive or behavioral deficits related to known or suspected CNS impairment, trauma, or neuropsychiatric disorders, including when the information will be useful in determining a diagnosis, prognosis, or informing treatment planning;
- Evaluate primary symptoms of impaired attention and concentration that can occur in many neurological and psychiatric conditions;
- Determine the potential impact of substances that may cause cognitive impairment (e.g., radiation, chemotherapy, prescribed or illicit drugs, toxins) or result in measurable improvement in cognitive function, including when this information is used to determine treatment planning;

- Conduct pre-surgical or treatment-related measurement of cognitive function to determine whether it is safe to proceed with a medical or surgical procedure that may impact brain function (e.g., deep brain stimulation, resection of brain tumors or arteriovenous malformations, epilepsy surgery, stem cell or organ transplant) or significantly alter a patient's functional status;
- Design, administer, and/or monitor outcomes of cognitive rehabilitation procedures, such as compensatory memory training for brain-injured patients;
- Measure cognitive or functional deficits in children and adolescents based on an inability to develop expected knowledge, skills or abilities as required to adapt to cognitive, social, emotional, or physical demands;
- Evaluate primary symptoms of impaired attention and concentration that can occur in many neurological and psychiatric conditions.

The American Psychological Association Guide (2019) also indicates that neuropsychological testing is not considered reasonable and necessary when:

- Administered for educational or vocational purposes that do not inform medical or health management
- Comprised exclusively of self-administered or self-scored inventories, or as screening tests of cognitive function or neurological disease (whether paper-and-pencil or computerized; e.g., AIMS, Folstein Mini-Mental Status Examination)
- The patient is neurologically, cognitively, or psychologically unable to participate in a meaningful way in the testing process
- Used as a routine screening tool given to the individual or to general populations
- Repeat testing is not required for medical decision-making
- Administered when the patient is currently under the influence or impaired by alcohol, drugs (prescription or illicit), or other substances
- The patient has been diagnosed previously with brain dysfunction, such as Alzheimer's disease, and there is no expectation that the testing would impact the patient's medical, functional, or behavioral management

The American Psychological Association published updated guidelines for the evaluation of dementia and age-related cognitive change. The guidelines include the following information regarding neuropsychological testing for this condition (American Psychological Association, 2012; updated 2021):

- Neuropsychological evaluation and cognitive testing remain among the most effective differential diagnostic methods in discriminating pathophysiological dementia from age-related cognitive decline, cognitive difficulties that are depression-related, and other related disorders. Even after reliable biological markers have been discovered, neuropsychological evaluation and cognitive testing will still be necessary to determine the onset of dementia, the functional expression of the disease process, the rate of decline, the functional capacities of the individual, and hopefully, response to therapies
- Comprehensive neuropsychological evaluations for dementia and cognitive change include tests of multiple cognitive domains, typically including memory, attention, perceptual and motor skills, language, visuospatial abilities, reasoning, and executive functions. Measures of mood and personality may be relevant in many cases. Psychologists are encouraged to refer to current compendia resources and the clinical research literature in selecting assessment instruments
- Psychologists are aware that standardized psychological and neuropsychological tests are important tools in the assessment of dementia and age-related cognitive change
- Technology assisted assessments (e.g., computer administered cognitive batteries, tele-health visits) are rapidly advancing but appropriate psychometric properties and normative data are nascent. These technologies may have significant advantages for older persons with limited mobility or health-care access but may also disadvantage older persons with limited experience and expertise interacting with technology

National Academy of Neuropsychology (NAN)

The National Academy of Neuropsychology developed an education paper to provide information to clinicians, healthcare administrators, and policy developers about the purpose, strengths, and limitations of computerized cognitive screening tests versus comprehensive neuropsychological evaluations. Screening tests are generally brief and narrow in scope, they can be administered during a routine clinical visit, and they can be helpful for identifying individuals in need of more comprehensive assessment. Some screening tests can also be helpful for monitoring treatment outcomes. Comprehensive neuropsychological assessments are multidimensional in nature and used for purposes such as identifying primary and secondary diagnoses, determining the nature and severity of a person's cognitive difficulties, determining functional limitations, and planning treatment and rehabilitation. Cognitive screening tests are expected to play an increasingly important role in identifying individuals with cognitive impairment and in determining which individuals should be referred for further neuropsychological

assessment. However, limitations of existing cognitive screening tests are present and cognitive screening tests should not be used as a replacement for comprehensive neuropsychological testing (Roebuck-Spencer et al, 2017).

In a policy for the evaluation of childhood learning disorders, the NAN states that when comprehensive information about a child's brain-related strengths and weaknesses is necessary to understand potential sources of the problem and implications for functioning, a neuropsychological evaluation is most often the best choice (Silver et al., 2006).

In a position paper on the diagnosis and management of sports-related concussion, the NAN states that neuropsychological evaluation is recommended for the diagnosis, treatment, and management of sports-related concussion at all levels of play (Moser et al., 2007).

National Institute for Health and Care Excellence (NICE)

In their guideline for the management of Multiple Sclerosis, NICE (2022) advises that the practitioner should be aware that the symptoms of MS can include cognitive problems, including memory problems. The guideline recommends that cognition should be assessed as part of the person's comprehensive review and that the assessment should be tailored to the person's needs, which may include a clinic interview or brief formal assessment, or consideration of a referral for a full neuropsychological assessment if needed.

Dementia assessment, management and support guidelines published by NICE (2018) state that following initial assessment and diagnosis of suspected dementia, patients are to be referred to a specialist once reversible causes of cognitive decline have been ruled out. Following standard, validated criteria use and assessment, neuropsychological testing may be considered if it is unclear:

- Whether the patient has cognitive impairment, or
- Whether their cognitive impairment is caused by dementia, or
- What the correct subtype diagnosis is

US Preventive Services Task Force (USPSTF)

A recommendation statement published by the US Preventive Services Task Force (2020) on screening for cognitive impairment in older adults, including neuropsychological testing, concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for cognitive impairment.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

In March 2021, the FDA cleared the ANAM Test system for Computerized Cognitive Assessment Aid for concussion. The ANAM system is an assessment aid in the management of concussion. The device consists of a software program that administers a battery of neurocognitive tests to an individual to assess their cognitive status. The device may be used with an off-the-shelf computer or a novel device. Refer to the following websites for more information:

- <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpd/classification.cfm?id=3918>
- https://www.accessdata.fda.gov/cdrh_docs/pdf20/K201376.pdf

(Accessed May 17, 2023)

In June 2015, the FDA cleared Cognivue through the de novo classification pathway. The de novo pathway is used for low- to moderate-risk medical devices that are not equivalent to an already legally marketed device. FDA identifies Cognivue as a "Computerized Cognitive Assessment Aid." According to the FDA, this test is indicated as an adjunctive tool for evaluating perceptual and memory function in individuals aged 55 to 95 years old. Refer to the following website for more information: https://www.accessdata.fda.gov/cdrh_docs/pdf13/DEN130033.pdf. (Accessed May 17, 2023)

On August 22, 2016, the FDA began to allow the marketing of two computerized neurocognitive tests for assessing individuals immediately following a suspected brain injury or concussion: ImPACT and ImPACT Pediatric (ImPACT Applications). Both tests were reviewed via the agency's de novo classification process, a pathway to market for certain "first-of-a-kind" and low- to moderate-risk medical devices. ImPACT and ImPACT Pediatric are computerized cognitive assessment aids intended for use in conjunction with standard medical evaluation for signs and symptoms of a head injury. ImPACT is designed to assess people

12 to 59 years of age, while ImPACT Pediatric is designed for children aged 5 to 11 years. The FDA states that these tests should not be used to “rule out a concussion or determine whether an injured player should return to a game.” Refer to the following websites for more information:

- http://www.accessdata.fda.gov/cdrh_docs/pdf15/DEN150037.pdf
- <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm517526.htm>

(Accessed May 17, 2023)

References

Abeare C, Messa I, Whitfield C, et al. Performance Validity in collegiate football athletes at baseline neurocognitive testing. *J Head Trauma Rehabil.* 2018 Nov 28.

Abeare CA, Messa I, Zuccato BG, et al. Prevalence of invalid performance on baseline testing for sport-related concussion by age and validity indicator. *JAMA Neurol.* 2018 Jun 1;75(6):697-703.

Ahto M, Isoaho R, Puolijoki H, et al. Cognitive impairment among elderly coronary heart disease patients. *Gerontology.* 1999;45(2):87-95.

Almeida OP, Garrido GJ, Beer C, et al. Coronary heart disease is associated with regional grey matter volume loss: implications for cognitive function and behavior. *Intern Med J.* 2008 Jul;38(7):599-606.

Alsalaheen B, Stockdale K, Pechumer D, et al. Measurement error in the Immediate Postconcussion Assessment and Cognitive Testing (ImPACT): systematic review. *J Head Trauma Rehabil.* 2016 Jul-Aug;31(4):242-51.

American Academy of Pediatrics, Section on Ophthalmology, Council on Children with Disabilities; American Academy of Ophthalmology; American Association for Pediatric Ophthalmology and Strabismus; American Association of Certified Orthoptists. Joint statement—Learning disabilities, dyslexia, and vision. *Pediatrics.* 2009 Aug;124(2):837-44. Reaffirmed 2014. Available at: <http://pediatrics.aappublications.org/content/124/2/837.long>. Accessed May 17, 2023.

American Psychiatric Association. Practice Guideline for the Treatment of Patients with Alzheimer's Disease and Other Dementias. Second Edition. October 2007. Available at: <https://www.psychiatry.org/psychiatrists/practice/clinical-practice-guidelines>. Accessed May 17, 2023.

American Psychological Association. APA guidelines for the evaluation of dementia and age-related cognitive change. *Am Psychol.* 2021, Feb. Available at: <https://www.apa.org/practice/guidelines/guidelines-dementia-age-related-cognitive-change.pdf>. Accessed May 17, 2023.

American Psychological Association. Psychological and Neuropsychological Testing Billing and Coding Guide. 2019. Available at: <https://www.apaservices.org/practice/reimbursement/health-codes/testing/billing-coding>. Accessed May 17, 2023.

Aslam RW, Bates V, Dundar Y, et al. A systematic review of the diagnostic accuracy of automated tests for cognitive impairment. *Int J Geriatr Psychiatry.* 2018 Apr;33(4):561-575.

Baron-Cohen S, Leslie AM, Frith U. Does the autistic child have a theory of mind? *Cognition.* 1985;21:37-46.

Bauer RM, Iverson GL, Cernich AN, et al. Computerized neuropsychological assessment devices: joint position paper of the American Academy of Clinical Neuropsychology and the National Academy of Neuropsychology. *Clin Neuropsychol.* 2012;26(2):177-96.

Bechtel N, Kobel M, Penner IK, et al. Attention-deficit/hyperactivity disorder in childhood epilepsy: a neuropsychological and functional imaging study. *Epilepsia.* 2012 Feb;53(2):325-33.

Becke M, Tucha L, Butzbach M, et al. Feigning Adult ADHD on a Comprehensive Neuropsychological Test Battery: An Analogue Study. *Int J Environ Res Public Health.* 2023 Feb 24;20(5):4070.

Belleville S, Fouquet C, Hudon C, et al.; Consortium for the early identification of Alzheimer's disease-Quebec. Neuropsychological measures that predict progression from mild cognitive impairment to Alzheimer's type dementia in older adults: a systematic review and meta-analysis. *Neuropsychol Rev.* 2017 Dec;27(4):328-353.

Böttrich N, Mückschel M, Dillenseger A, et al. On the Reliability of Examining Dual-Tasking Abilities Using a Novel E-Health Device—A Proof of Concept Study in Multiple Sclerosis. *J Clin Med.* 2020 Oct 25;9(11):3423.

Broglio SP, Katz BP, Zhao S, et al.; CARE Consortium Investigators. Test-retest reliability and interpretation of common concussion assessment tools: Findings from the NCAA-DoD CARE Consortium. *Sports Med.* 2018 May;48(5):1255-1268.

Brunkhorst-Kanaan N, Verdenhalven M, Kittel-Schneider S, et al. The quantified behavioral test-a confirmatory test in the diagnostic process of adult ADHD? *Front Psychiatry*. 2020 Mar 20;11:216.

Burton KL, Williams TA, Catchpole SE, et al. Long-term neuropsychological outcomes of childhood onset acute disseminated encephalomyelitis (ADEM): a Meta-Analysis. *Neuropsychol Rev*. 2017 Mar 31.

Cahn-Hidalgo D, Estes PW, Benabou R. Validity, reliability, and psychometric properties of a computerized, cognitive assessment test (Cognivue®). *World J Psychiatry*. 2020 Jan 19;10(1):1-11.

Cameron J, Kure CE, Pressler SJ, et al. Diagnostic accuracy of cognitive screening instruments in heart failure: a systematic review. *J Cardiovasc Nurs*. 2016 Sep-Oct;31(5):412-24.

Carlozzi NE, Kirsch NL, Kisala PA, et al. An examination of the Wechsler Adult Intelligence Scales, Fourth Edition (WAIS-IV) in individuals with complicated mild, moderate, and severe traumatic brain injury (TBI). *Clin Neuropsychol*. 2015;29(1):21-37.

Centers for Disease Control and Prevention. Traumatic Brain Injury & Concussion. May 13, 2021. Available at: <http://www.cdc.gov/TraumaticBrainInjury/index.html>. Accessed May 17, 2023.

Chan JYC, Bat BKK, Wong A, et al. Evaluation of digital drawing tests and paper-and-pencil drawing tests for the screening of mild cognitive impairment and dementia: A systematic review and meta-analysis of diagnostic studies. *Neuropsychol Rev*. 2022 Sep;32(3):566-576.

Chan JYC, Yau STY, Kwok TCY, et al. Diagnostic performance of digital cognitive tests for the identification of MCI and dementia: a systematic review. *Ageing Res Rev*. 2021 Dec;72:101506.

Chen C, Dong X, Gu P, et al. Attention impairment during the interictal state in migraineurs without aura: a cross-sectional study. *J Pain Res*. 2021 Oct 6;14:3073-3083.

Chen X, Wong A, Ye R, et al. Validation of NINDS-CSN neuropsychological battery for vascular cognitive impairment in Chinese stroke patients. *BMC Neurol*. 2015 Mar 3;15:20.

Cochereau J, Herbet G, Duffau H. Patients with incidental WHO grade II glioma frequently suffer from neuropsychological disturbances. *Acta Neurochir (Wien)*. 2016 Feb;158(2):305-12.

Cole WR, Arrieux JP, Ivins BJ, et al. A comparison of four computerized neurocognitive assessment tools to a traditional neuropsychological test battery in service members with and without mild traumatic brain injury. *Arch Clin Neuropsychol*. 2018 Feb 1;33(1):102-119.

Cosgrave C, Fuller C, Kung S, et al. A comparison of clinical assessment with common diagnostic tools for monitoring concussion recovery in adolescent rugby union players. *Phys Ther Sport*. 2023 Apr 6;61:165-171.

Costa A, Bak T, Caffarra P, et al. The need for harmonization and innovation of neuropsychological assessment in neurodegenerative dementias in Europe: consensus document of the Joint Program for Neurodegenerative Diseases Working Group. *Alzheimer's Res Ther*. 2017 Apr 17;9(1):27.

Delgado-Álvarez A, Delgado-Alonso C, Goudsmit M, et al. Validation of a brief cross-cultural cognitive screening test in Multiple Sclerosis. *Mult Scler Relat Disord*. 2022 Nov;67:104091.

Dresler T, Lürding R, Paelecke-Habermann Y, et al. Cluster headache and neuropsychological functioning. *Cephalalgia*. 2012 Aug;32(11):813-21.

DSM-5 Diagnostic and Statistical Manual of Mental Disorders. American Psychiatric Association. Fifth edition. Washington DC: American Psychiatric Association, 2013.

Echemendia RJ, Iverson GL, McCrea M, et al. Advances in neuropsychological assessment of sport-related concussion. *Br J Sports Med*. 2013 Apr;47(5):294-8.

Farnsworth JL 2nd, Dargo L, Ragan BG, et al. Reliability of computerized neurocognitive tests for concussion assessment: a meta-analysis. *J Athl Train*. 2017 Sep;52(9):826-833.

Filipek PA, Accardo PJ, Ashwal S, et al. Practice parameter: screening and diagnosis of autism: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Child Neurology Society. *Neurology* 2000 Aug 22;55(4):468-79. Reaffirmed on October 18, 2003, July 28, 2006, July 10, 2010, and August 9, 2014.

Filipčíková M, Quang H, Cassel A, et al. Exploring neuropsychological underpinnings of poor communication after traumatic brain injury: The role of apathy, disinhibition, and social cognition. *Int J Lang Commun Disord*. 2022 Dec 21.

Filippini M, Ardu E, Stefanelli S, et al. Neuropsychological profile in new-onset benign epilepsy with centrotemporal spikes (BECTS): Focusing on executive functions. *Epilepsy Behav.* 2016 Jan;54:71-9.

Fink HA, Hemmy LS, Linskens EJ, et al. Diagnosis and treatment of clinical Alzheimer's-type dementia: A systematic review. Comparative Effectiveness Review No. 223. (Prepared by the Minnesota Evidence-based Practice Center under Contract No. 290- 2015-00008-I.) AHRQ Publication No. 20-EHC003. Rockville, MD: Agency for Healthcare Research and Quality; April 2020.

Foti M, Lo Buono V, Corallo F, et al. Neuropsychological assessment in migraine patients: a descriptive review on cognitive implications. *Neurol Sci.* 2017 Apr;38(4):553-562.

Fuchs TA, Gillies J, Jaworski MG 3rd, et al. Repeated forms, testing intervals, and SDMT performance in a large multiple sclerosis dataset. *Mult Scler Relat Disord.* 2022 Dec;68:104375.

Gallagher R, Ouyang ML, Tofler G, et al. Sensitivity and specificity of 5 min cognitive screening tests in patients with acute coronary syndrome. *Eur J Cardiovasc Nurs.* 2023 Mar 1;22(2):166-174.

Gaudet CE, Weyandt LL. Immediate Post-Concussion and Cognitive Testing (ImPACT): a systematic review of the prevalence and assessment of invalid performance. *Clin Neuropsychol.* 2017 Jan;31(1):43-58.

Giacino JT, Katz DI, Schiff ND, et al. Practice guideline update recommendations summary: Disorders of consciousness: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology; the American Congress of Rehabilitation Medicine; and the National Institute on Disability, Independent Living, and Rehabilitation Research. *Neurology.* 2018 Sep 4;91(10):450-460.

Giza CC, Kutcher JS, Ashwal S, et al. Summary of evidence-based guideline update: Evaluation and management of concussion in sports: Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology.* 2013 Mar 18. Reaffirmed April 30, 2022.

Grau-López L, Jiménez M, Ciurans J, et al. Importance of neuropsychological and clinical features to predict seizure control in medically treated patients with mesial temporal epilepsy and hippocampal sclerosis. *Epilepsy Behav.* 2017 Feb 24;69:121-125.

Grubb NR, Simpson C, Fox KA. Memory function in patients with stable, moderate to severe cardiac failure. *Am Heart J.* 2000 Jul;140(1):E1-5.

Groppell S, Soto-Ruiz KM, Flores B, et al. A rapid, mobile neurocognitive screening test to aid in identifying cognitive impairment and dementia (BrainCheck): Cohort Study. *JMIR Aging.* 2019 Mar 21;2(1):e12615.

Halstead ME, Walter KD, Moffatt K; Council on Sports Medicine and Fitness. Sport-related concussion in children and adolescents. *Pediatrics.* 2018 Dec;142(6).

Hang B, Babcock L, Hornung R, et al. Can computerized neuropsychological testing in the emergency department predict recovery for young athletes with concussions? *Pediatr Emerg Care.* 2015 Oct;31(10):688-93.

Hanks RA, Jackson AM, Crisanti LK. Predictive validity of a brief outpatient neuropsychological battery in individuals 1-25 years post traumatic brain injury. *Clin Neuropsychol.* 2016 Oct;30(7):1074-86.

Happe F, Frith U. The weak coherence account: detail-focused style in autism spectrum disorders. *J Autism Dev Disord.* 2006;36:5-25.

Harmon KG, Clugston JR, Dec K, et al. American Medical Society for Sports Medicine Position Statement on Concussion in Sport. *Clin J Sport Med.* 2019 Mar;29(2):87-100.

Hartman E, Houwen S, Scherder E, et al. On the relationship between motor performance and executive functioning in children with intellectual disabilities. *J Intellect Disabil Res.* 2010 May;54(5):468-77.

Hollis C, Hall CL, Guo B, et al.; the AQUA Trial Group. The impact of a computerized test of attention and activity (QbTest) on diagnostic decision-making in children and young people with suspected attention deficit hyperactivity disorder: single-blind randomized controlled trial. *J Child Psychol Psychiatry.* 2018 Dec;59(12):1298-1308.

Ivins BJ, Arrieux JP, Cole WR. An initial psychometric analysis of the Brain Gauge tactile-based test battery and its potential for clinical use assessing patients with acute mild traumatic brain injury. *Arch Clin Neuropsychol.* 2022 Oct 19;37(7):1564-1578.

Ivins BJ, Arrieux JP, Schwab KA, et al. Using rates of low scores to assess agreement between brief computerized neuropsychological assessment batteries: A clinically based approach for psychometric comparisons. *Arch Clin Neuropsychol.* 2019 Feb 23. Pii: acz004.

Johnson S, Strauss V, Gilmore C, et al. Learning disabilities among extremely preterm children without neurosensory impairment: Comorbidity, neuropsychological profiles, and scholastic outcomes. *Early Hum Dev.* 2016 Dec;103:69-75.

Kontos AP, Braithwaite R, Dakan S, et al. Computerized neurocognitive testing within 1 week of sport-related concussion: meta-analytic review and analysis of moderating factors. *J Int Neuropsychol Soc.* 2014 Mar;20(3):324-32.

Lo Buono V, Bonanno L, Palmeri R, et al. Relation among psychopathological symptoms, neuropsychological domains, and functional disability in subacute poststroke rehabilitation. *J Stroke Cerebrovasc Dis.* 2018 Feb 5. Pii: S1052-3057(17)30707-3.

Lozano-Soto E, Cruz-Gómez ÁJ, Rashid-López R, et al. Neuropsychological and neuropsychiatric features of chronic migraine patients during the interictal phase. *J Clin Med.* 2023 Jan 9;12(2):523.

Lumba-Brown A, Yeates KO, Sarmiento K, et al. Centers for Disease Control and Prevention Guideline on the Diagnosis and Management of Mild Traumatic Brain Injury Among Children. *JAMA Pediatr.* 2018 Nov 1;172(11):e182853.

Mahncke HW, DeGutis J, Levin H, et al. A randomized clinical trial of plasticity-based cognitive training in mild traumatic brain injury. *Brain.* 2021 Aug 17;144(7):1994-2008.

MacDonald J, Duerson D. Reliability of a Computerized Neurocognitive Test in Baseline Concussion Testing of High School Athletes. *Clin J Sport Med.* 2015 Jul;25(4):367-72.

McCrory P, Meeuwisse W, Dvorak J, et al. Consensus statement on concussion in sport-the 5(th) international conference on concussion in sport held in Berlin, October 2016. *Br J Sports Med.* 2017 Apr 26.

Meskal I, Gehring K, Rutten GJ, et al. Cognitive functioning in meningioma patients: a systematic review. *J Neurooncol.* 2016 Jun;128(2):195-205.

Moore RC, Paolillo EW, Heaton A, et al. Clinical utility of the UCSD Performance-Based Skills Assessment-Brief (UPSA-B) in adults living with HIV: Associations with neuropsychological impairment and patient-reported everyday functioning difficulties. *PLoS One.* 2017 Aug 24;12(8):e0183614.

Moser RS, Iverson GL, Echemendia RJ, et al; NAN Policy and Planning Committee. Neuropsychological evaluation in the diagnosis and management of sports-related concussion. *Arch Clin Neuropsychol.* 2007 Nov;22(8):909-16.

Nakayama Y, Covassin T, Schatz P, et al. Examination of the Test-Retest Reliability of a Computerized Neurocognitive Test Battery. *Am J Sports Med.* 2014 Jun 6;42(8):2000-2005.

Nascimento S, Baierle M, Göethel G, et al. Associations among environmental exposure to manganese, neuropsychological performance, oxidative damage, and kidney biomarkers in children. *Environ Res.* 2016 May;147:32-43.

National Institute for Health and Care Excellence (NICE). Dementia: assessment, management and support for people living with dementia and their carers. Available at: <https://www.nice.org.uk/guidance/ng97>. Accessed May 18, 2023.

National Institute for Health and Care Excellence (NICE). Multiple sclerosis in adults: management. Available at: <https://www.nice.org.uk/guidance/ng220/resources/multiple-sclerosis-in-adults-management-pdf-66143828948677>. Accessed May 18, 2023.

Nelson LD, Furger RE, Gikas P, et al. Prospective, head-to-head study of three computerized neurocognitive assessment tools part 2: utility for assessment of mild traumatic brain injury in emergency department patients. *J Int Neuropsychol Soc.* 2017 Apr;23(4):293-303.

Nelson LD, LaRoche AA, Pfaller AY, et al. Prospective, head-to-head study of three computerized neurocognitive assessment tools (CNTs): reliability and validity for the assessment of sport-related concussion. *J Int Neuropsychol Soc.* 2016 Jan;22(1):24-37.

Oliva I, Losa J. Validation of the Computerized Cognitive Assessment Test: NNCT. *Int J Environ Res Public Health.* 2022 Aug 23;19(17):10495.

Ozonoff S, Pennington BF, Rogers SJ. Executive function deficits in high functioning autistic individuals: relationship to theory of mind. *J Child Psychol Psychiatry.* 1991;32:1081-1105.

Pagán AF, Huizar YP, Schmidt AT. Conner's Continuous Performance Test and adult ADHD: A systematic literature review. *J Atten Disord.* 2023 Feb;27(3):231-249.

Parra-Díaz P, García-Casares N. Memory assessment in patients with temporal lobe epilepsy to predict memory impairment after surgery: A systematic review. *Neurologia.* 2017 Apr 19. Pii: S0213-4853(17)30150-0.

Paterson TSE, Sivajohan B, Gardner S, et al. Accuracy of a self-administered online cognitive assessment in detecting amnesic mild cognitive impairment. *J Gerontol B Psychol Sci Soc Sci*. 2022 Feb 3;77(2):341-350.

Patrikelis P, Gatzonis S, Siatouni A, et al. Preoperative neuropsychological presentation of patients with refractory frontal lobe epilepsy. *Acta Neurochir (Wien)*. 2016 Jun;158(6):1139-50.

Pedersen KF, Larsen JP, Tysnes OB, et al., Natural course of mild cognitive impairment in Parkinson disease: A 5-year population-based study. *Neurology*. 2017 Feb 21;88(8):767-774.

Petersen RC, Lopez O, Armstrong MJ, et al. Practice guideline update summary: Mild cognitive impairment: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018 Jan 16;90(3):126-135. Reaffirmed January 30, 2021.

Pranckeviciene A, Deltuva VP, Tamasauskas A, et al. Association between psychological distress, subjective cognitive complaints and objective neuropsychological functioning in brain tumor patients. *Clin Neurol Neurosurg*. 2017 Dec;163:18-23.

Qin T, Chen C. Cognitive Dysfunction in Migraineurs. *Medicina (Kaunas)*. 2022 Jun 29;58(7):870.

Racine AM, Clark LR, Berman SE, et al. Associations between performance on an abbreviated cogstate battery, other measures of cognitive function, and biomarkers in people at risk for Alzheimer's disease. *J Alzheimers Dis*. 2016 Oct 18;54(4):1395-1408.

Rayes HA, Tani C, Kwan A, et al. What is the prevalence of cognitive impairment in lupus and which instruments are used to measure it? A systematic review and meta-analysis. *Semin Arthritis Rheum*. 2018 Oct;48(2):240-255.

Roebuck-Spencer TM, Glen T, Puente AE, et al. Cognitive screening tests versus comprehensive neuropsychological test batteries: A National Academy of Neuropsychology education paper. *Arch Clin Neuropsychol*. 2017 Mar 10:1-8.

Romero-Garcia R, Owen M, McDonald A, et al. Assessment of neuropsychological function in brain tumor treatment: a comparison of traditional neuropsychological assessment with app-based cognitive screening. *Acta Neurochir (Wien)*. 2022 Aug;164(8):2021-2034.

Rubin LH, Severson J, Marcotte TD, et al. Tablet-based cognitive impairment screening for adults with HIV seeking clinical care: observational study. *JMIR Ment Health*. 2021 Sep 9;8(9):e25660.

Ruet A, Brochet B. Cognitive assessment in patients with multiple sclerosis: From neuropsychological batteries to ecological tools. *Ann Phys Rehabil Med*. 2018 Feb 17. pii: S1877-0657(18)30012-5.

Sauvé MJ, Lewis WR, Blankenbiller M, et al. Cognitive impairments in chronic heart failure: a case-controlled study. *J Card Fail*. 2009 Feb;15(1):1-10.

Scharre DW, Chang SI, Nagaraja HN, et al. Self-administered gerocognitive examination: longitudinal cohort testing for the early detection of dementia conversion. *Alzheimers Res Ther*. 2021 Dec 6;13(1):192.

Shopin L, Shenhar-Tsarfaty S, Ben Assayag E, et al. Cognitive assessment in proximity to acute ischemic stroke/transient ischemic attack: comparison of the Montreal cognitive assessment test and mindstreams computerized cognitive assessment battery. *Dement Geriatr Cogn Disord*. 2013;36(1-2):36-42.

Sigurdardottir AH, Knudtzen FC, Nymark A, Bang M. Fatigue and cognitive impairment in neuroborreliosis patients posttreatment-A neuropsychological retrospective cohort study. *Brain Behav*. 2022 Sep;12(9):e2719.

Silberg T, Ahoniska-Assa J, Bord A, et al. In the eye of the beholder: Using a multiple-informant approach to examine the mediating effect of cognitive functioning on emotional and behavioral problems in children with an active epilepsy. *Seizure*. 2020 Nov;82:31-38.

Silver CH, Blackburn LB, Arffa S, et al. The importance of neuropsychological assessment for the evaluation of childhood learning disorders NAN Policy and Planning Committee. *Arch Clin Neuropsychol*. 2006 Oct;21(7):741-4.

Söderström H, Brocki K, Kleberg JL, et al. Neurocognitive functions before and after radiotherapy in pediatric brain tumor survivors. *Pediatr Neurol*. 2022 Aug;133:21-29.

Takagi M, Hearps SJC, Babl FE, et al. Does a computerized neuropsychological test predict prolonged recovery in concussed children presenting to the ED? *Child Neuropsychol*. 2020 Jan;26(1):54-68.

Talwalker S, Overall JE, Srirama MK, et al. Cardinal features of cognitive dysfunction in Alzheimer's disease: a factor-analytic study of the Alzheimer's Disease Assessment Scale. *J Geriatr Psychiatr Neurol*. 1996;9:39-46.

Tan HH, Xu J, Teoh HL, et al. Decline in changing Montreal Cognitive Assessment (MoCA) scores is associated with post-stroke cognitive decline determined by a formal neuropsychological evaluation. *PLoS One*. 2017 Mar 27;12(3):e0173291.

Tekin S, Bir LS, Oncel C, et al. Evaluation of cognitive dysfunction by the clock drawing test in multiple sclerosis and clinically isolated syndrome patients: Correlation with other neuropsychological tests. *Neurosciences (Riyadh)*. 2022 Oct;27(4):251-256.

Trickett J, Bernardi M, Fahy A, et al. Neuropsychological abilities underpinning academic attainment in children born extremely preterm. *Child Neuropsychol*. 2022 Aug;28(6):746-767.

Tsushima WT, Yamamoto MH, Ahn HJ, et al. Invalid baseline testing with ImPACT: Does sandbagging occur with high school athletes? *Appl Neuropsychol Child*. 2019 Aug 13:1-10.

US Preventive Services Task Force, Owens DK, Davidson KW, et al. Screening for cognitive impairment in older adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2020 Feb 25;323(8):757-763.

Volkmar F, Siegel M, Woodbury-Smith M, et al. American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI). Practice parameter for the assessment and treatment of children and adolescents with autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry*. 2014 Feb;53(2):237-57.

Vollmer T, Huynh L, Kelley C, et al. Relationship between brain volume loss and cognitive outcomes among patients with multiple sclerosis: a systematic literature review. *Neurol Sci*. 2016 Feb;37(2):165-79.

von Bismarck O, Dankowski T, Ambrosius B, et al. Treatment choices and neuropsychological symptoms of a large cohort of early MS. *Neurol Neuroimmunol Neuroinflamm*. 2018 Mar 1;5(3):e446.

Vyshedskiy A, Netson R, Fridberg E, et al. Boston cognitive assessment (BOCA) - a comprehensive self-administered smartphone- and computer-based at-home test for longitudinal tracking of cognitive performance. *BMC Neurol*. 2022 Mar 15;22(1):92.

Walsh KS, Noll RB, Annett RD, et al. Standard of Care for Neuropsychological Monitoring in Pediatric Neuro-Oncology: Lessons From the Children's Oncology Group (COG). *Pediatr Blood Cancer*. 2016 Feb;63(2):191-5.

Weissberger GH, Strong JV, Stefanidis KB, et al. Diagnostic Accuracy of Memory Measures in Alzheimer's Dementia and Mild Cognitive Impairment: a Systematic Review and Meta-Analysis. *Neuropsychol Rev*. 2017 Dec;27(4):354-388.

Wilson S, Milanini B, Javandel S, et al. Validity of digital assessments in screening for HIV-related cognitive impairment: a review. *Curr HIV/AIDS Rep*. 2021 Dec;18(6):581-592.

Wilson SJ, Baxendale S, Barr W, et al. Indications and expectations for neuropsychological assessment in routine epilepsy care: Report of the ILAE Neuropsychology Task Force, Diagnostic Methods Commission, 2013-2017. *Epilepsia*. 2015 May;56(5):674-81.

Winstein CJ, Stein J, Arena R, et al.; American Heart Association Stroke Council, Council on Cardiovascular and Stroke Nursing, Council on Clinical Cardiology, and Council on Quality of Care and Outcomes Research. Guidelines for Adult Stroke Rehabilitation and Recovery: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2016 Jun;47(6):e98-e169.

Wojcik CM, Beier M, Costello K, et al; National MS Society Cognition Work Team. Computerized neuropsychological assessment devices in multiple sclerosis: A systematic review. *Mult Scler*. 2019 Dec;25(14):1848-1869.

Ye S, Sun K, Huynh D, et al. A computerized cognitive test battery for detection of dementia and mild cognitive impairment: instrument validation study. *JMIR Aging*. 2022 Apr 15;5(2):e36825.

Zuo L, Dong Y, Liao X, et al. Risk factors for decline in Montreal Cognitive Assessment (MoCA) scores in patients with acute transient ischemic attack and minor stroke. *J Clin Hypertens (Greenwich)*. 2022 Jul;24(7):851-857.

Zweifel-Zehnder AE, Stienen MN, Chicherio C, et al.; Swiss SOS study group. Call for uniform neuropsychological assessment after aneurysmal subarachnoid hemorrhage: Swiss recommendations. *Acta Neurochir (Wien)*. 2015 Sep;157(9):1449-58.

Policy History/Revision Information

Date	Summary of Changes
10/01/2023	<p data-bbox="337 1738 641 1768">Supporting Information</p> <ul data-bbox="337 1774 1396 1864" style="list-style-type: none"> <li data-bbox="337 1774 1396 1801">• Updated <i>Clinical Evidence</i> and <i>References</i> sections to reflect the most current information <li data-bbox="337 1801 1396 1829">• Removed <i>Benefit Considerations</i> section <li data-bbox="337 1829 1396 1864">• Archived previous policy version CS083TN.N

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state, or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state, or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state, or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state, or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.