

Tzield™ (Teplizumab-Mzwv)

Policy Number: 2023D00117A
Effective Date: February 1, 2023

[Instructions for Use](#)

Table of Contents	Page
Coverage Rationale	1
Applicable Codes	2
Background	2
Benefit Considerations	2
Clinical Evidence	3
U.S. Food and Drug Administration	3
References	3
Policy History/Revision Information	3
Instructions for Use	3

Community Plan Policy
• Tzield™ (Teplizumab-Mzwv)

Coverage Rationale

See [Benefit Considerations](#)

Tzield has been added to the Review at Launch program. Some members may not be eligible for coverage of this medication at this time. Refer to the Medical Benefit Drug Policy titled [Review at Launch for New to Market Medications](#) for additional details.

Tzield, administered as a one-time 14-day course of therapy, is proven to delay the onset of stage 3 type 1 diabetes in patients that meet the following criteria:

- Diagnosis of stage 2 type 1 diabetes confirmed by all of following:
 - At least two positive pancreatic islet autoantibodies
 - Dysglycemia without overt hyperglycemia using an oral glucose tolerance test (OGTT)
 - Clinical history of patient does not suggest type 2 diabetes
 and
- Dosing is in accordance with the United States Food and Drug Administration approved labeling; and
- Patient has not been previously treated with Tzield; and
- Authorization will be issued for no more than 14 doses

Tzield is medically necessary when all of the following criteria are met:

- Diagnosis of stage 2 type 1 diabetes confirmed by all of the following:
 - At least two of the following pancreatic islet autoantibodies:
 - Glutamic acid decarboxylase 65 (GAD) autoantibodies
 - Insulin autoantibody (IAA)
 - Insulinoma-associated antigen 2 autoantibody (IA-2A)
 - Zinc transporter 8 autoantibody (ZnT8A)
 - Islet cell autoantibody (ICA)
 and
 - Dysglycemia without overt hyperglycemia on an oral glucose tolerance test (OGTT) defined by one of the following:
 - Fasting blood glucose > 110mg/dL and < 126 mg/dL; or

- 2-hour post-prandial plasma glucose level \geq 140 mg/dL and $<$ 200 mg/dL; or
 - 30-, 60-, or 90-minute post-prandial glucose level \geq 200 mg/dL
- and
- Clinical history of patient does not suggest type 2 diabetes
- and
- Prescribed by an endocrinologist; and
 - Dosing is in accordance with the United States Food and Drug Administration approved labeling; and
 - Patient has not been previously treated with Tzield; and
 - Authorization will be issued for no more than 14 doses

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
C9399	Unclassified drugs or biologicals
J3490	Unclassified drugs
J3590	Unclassified biologics

Diagnosis Code	Description
E10.8	Type 1 diabetes mellitus with unspecified complications
E10.9	Type 1 diabetes mellitus without complications

Background

Type 1 diabetes (T1D) is a chronic autoimmune disease that leads to destruction of insulin-producing beta cells and dependence on exogenous insulin for survival. Approximately 1 to 1.5 million Americans have TD1, which is one of the most common diseases of childhood. Although it can appear at any age, TD1 is usually diagnosed in children and young adults. A person is at higher risk for T1D if they have a parent, brother, or sister with T1D, although most patients do not have a family history. Type 1 diabetes progresses through asymptomatic stages before the development of overt hyperglycemia. These stages are characterized by the appearance of autoantibodies (stage 1) and then dysglycemia (stage 2). In stage 2, metabolic responses to a glucose load are impaired but the level of glycosylated hemoglobin remains normal. Insulin therapy and glucose monitoring are currently the standard of care for treating the clinical stage, Stage 3 T1D.

Teplizumab-mzvw is a CD3-directed monoclonal antibody which binds CD3 on the surface of T lymphocytes. Teplizumab-mzvw may deactivate the T lymphocytes that attack pancreatic insulin-producing beta cells, while increasing the proportion of regulatory T lymphocytes that help moderate the immune response.

Benefit Considerations

Some Certificates of Coverage allow for coverage of experimental/investigational/unproven treatments for life-threatening illnesses when certain conditions are met. The member specific benefit plan document must be consulted to make coverage decisions for this service. Some states mandate benefit coverage for off-label use of medications for some diagnoses or under some circumstances when certain conditions are met. Where such mandates apply, they supersede language in the benefit document or in the medical or drug policy. Benefit coverage for an otherwise unproven service for the treatment of serious rare diseases may occur when certain conditions are met. Refer to the Policy and Procedure addressing the treatment of serious rare diseases.

Clinical Evidence

Proven

Teplizumab-mzwv is indicated to delay the onset of Stage 3 type 1 diabetes in adults and pediatric patients 8 years of age and older with Stage 2 type diabetes.

The efficacy of teplizumab-mzwv was established in a randomized double-blind, event-driven, placebo-controlled study in 76 patients, 8 to 49 years of age with Stage 2 T1D. Patients were randomized to receive teplizumab-mzwv or placebo once daily by intravenous (IV) infusion for 14 days. The primary efficacy endpoint was the time from randomization to development of Stage 3 T1D was diagnosed in 20 (45%) of the teplizumab-treated patients and in 23 (72%) of the placebo-treated patients. A Cox proportional hazards model, stratified by age and oral glucose tolerance test status at randomization, demonstrated that the median time from randomization to Stage 3 T1D diagnosis was 50 months in the teplizumab-mzwv group and 25 months in the placebo group, for a difference of 25 months. With a median follow-up time of 51 months, therapy with Tziel resulted in a statistically significant delay in the development of Stage 3 T1D (hazard ratio 0.41, 95% CI: 0.22, 0.78; p = 0.0066).

The most common adverse reactions (> 10%) with teplizumab-mzwv use were lymphopenia, rash, leukopenia, and headache.

Lymphocyte count decreased to a nadir on day 5 (total decrease, 72.3%; interquartile range, 82.1 to 68.4; P < 0.001) and resolved by day 45 in all participants except one; in that participant, the lymphocyte counts returned to the normal range on day 105. A spontaneously resolving rash occurred in 16 (36%) of participants who received teplizumab-mzwv.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Teplizumab-mzwv is indicated to delay the onset of Stage 3 type 1 diabetes in adults and pediatric patients 8 years of age and older with Stage 2 type diabetes.

References

1. Tziel™ [package insert]. Red Bank, NJ: Provention Bio Inc.; November 2022.
2. Herold KC, Bundy BN, Long SA, et al. An Anti-CD3 Antibody, Teplizumab, in Relatives at Risk for Type 1 Diabetes [published correction appears in *N Engl J Med*. 2020 Feb 6;382(6):586]. *N Engl J Med*. 2019;381(7):603-613. doi:10.1056/NEJMoa1902226.

Policy History/Revision Information

Date	Summary of Changes
02/01/2023	<ul style="list-style-type: none">• New Medical Benefit Drug Policy

Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates.

UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

This Medical Benefit Drug Policy may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence ([Medicare IOM Pub. No. 100-16, Ch. 4, §90.5](#)).

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.