

Epidural Steroid Injections for Spinal Pain

Policy Number: CS039.U
Effective Date: August 1, 2022

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	2
Applicable Codes	2
Description of Services	4
Clinical Evidence	4
U.S. Food and Drug Administration	10
References	10
Policy History/Revision Information	12
Instructions for Use	13

Related Community Plan Policies
<ul style="list-style-type: none"> Ablative Treatment for Spinal Pain Facet Joint Injections for Spinal Pain Occipital Nerve Injections and Ablation (Including Occipital Neuralgia and Headache)
Commercial Policy
<ul style="list-style-type: none"> Epidural Steroid Injections for Spinal Pain
Medicare Advantage Coverage Summary
<ul style="list-style-type: none"> Pain Management and Rehabilitation

Application

This Medical Policy does not apply to the states listed below; refer to the state-specific policy/guideline, if noted:

State	Policy/Guideline
Indiana	None
Kentucky	None
Louisiana	Epidural Steroid Injections (for Louisiana Only)
Mississippi	Epidural Steroid Injections for Spinal Pain (for Mississippi Only)
Nebraska	Epidural Steroid Injections for Spinal Pain (for Nebraska Only)
New Jersey	Epidural Steroid Injections for Spinal Pain (for New Jersey Only)
North Carolina	Epidural Steroid Injections for Spinal Pain (for North Carolina Only)
Pennsylvania	Epidural Steroid Injections for Spinal Pain (for Pennsylvania Only)
Tennessee	Epidural Steroid Injections for Spinal Pain (for Tennessee Only)

Coverage Rationale

Epidural Steroid Injections (ESI) are proven and medically necessary when the following criteria are met:

- The injection is intended for the short term management of acute or subacute radicular pain and;
- The radicular pain is unresponsive to Conservative Treatment:
 - Pharmacotherapy such as NSAIDs or acetaminophen ≥ 3 weeks or;
 - Activity modification ≥ 4 weeks (including but not limited to heavy lifting, bending, spinal torsion activities) or;
 - PT or home exercise ≥ 4 weeks

The following are unproven and not medically necessary due to insufficient evidence of efficacy:

- The use of ultrasound guidance for ESIs
- ESI for all other indications of the spine not included above

Epidural Steroid Injection Limitations

- A maximum of three (3) ESI sessions (per region, regardless of level, location, or side) per year
 - A session is defined as one date of service in which ESI(s) is performed
 - A region is defined by either the region of the cervical, thoracic or lumbosacral
 - A year is defined as the 12-month period starting from the date of service of the first approved injection
- Repeat ESIs may be provided only if:
 - The initial injection resulted in $\geq 50\%$ pain relief achieved for 3 or more months
 - The initial injection resulted in a functional improvement as measured by validated measurement tools, such as The Oswestry Disability Index
 - Repeat injections do not exceed 3 per year, per region

Definitions

Acute Low Back Pain: Low back pain present for up to six weeks. The early acute phase is defined as less than two weeks and the late acute phase is defined as two to six weeks, secondary to the potential for delayed-recovery or risk phases for the development of chronic low back pain. Low back pain can occur on a recurring basis. If there has been complete recovery between episodes, it is considered acute recurrent. (Goertz et al. 2012)

Conservative Therapy: Consists of an appropriate combination of medication (for example, NSAIDs, analgesics, etc.) in addition to physical therapy, spinal manipulation therapy, cognitive behavioral therapy (CBT) or other interventions based on the individual's specific presentation, physical findings and imaging results. (AHRQ 2013; Qassem 2017; Summers 2013)

Epidural Steroid Injections (ESI): Is a nonsurgical treatment for managing radiculopathy caused by disc herniation or degenerative changes in the vertebrae. Steroids are injected directly into the epidural space of the spine. The goal of ESI is to relieve pain, improve function, and improve quality of life. (Patel 2021)

Non-Radicular Back Pain: Pain which does not radiate along a dermatome (sensory distribution of a single root). Appropriate imaging does not reveal signs of spinal nerve root compression and there is no evidence of spinal nerve root compression seen on clinical exam. (Lenahan, 2018)

Oswestry Disability Index (also known as the Oswestry Low Back Pain Disability Questionnaire): A tool that researchers and disability evaluators use to measure a patient's permanent functional disability. The test is considered the 'gold standard' of low back functional outcome tools. (Fairbank 2000)

Radicular Back Pain: Pain which radiates from the spine into the extremity along the course of the spinal nerve root. The pain should follow the pattern of a dermatome associated with the irritated nerve root identified. (Lenahan, 2018)

Radiculopathy: Radiculopathy is characterized by pain which radiates from the spine to extend outward to cause symptoms away from the source of the spinal nerve root irritation. (Lenahan, 2018)

Sub-Acute Low Back Pain: Low back pain with duration of greater than six weeks after injury but no longer than 12 weeks after onset of symptoms. (Goertz et al., 2012)

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
62320	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance
62321	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (i.e., fluoroscopy or CT)
62322	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance
62323	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); with imaging guidance (i.e., fluoroscopy or CT)
64479	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); cervical or thoracic, single level
64480	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); cervical or thoracic, each additional level (List separately in addition to code for primary procedure)
64483	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); lumbar or sacral, single level
64484	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); lumbar or sacral, each additional level (List separately in addition to code for primary procedure)

CPT® is a registered trademark of the American Medical Association

Diagnosis Code	Description
All Regions	
M47.25	Other spondylosis with radiculopathy, thoracolumbar region
M51.15	Intervertebral disc disorders with radiculopathy, thoracolumbar region
M96.1	Postlaminectomy syndrome, not elsewhere classified
Cervical/Thoracic	
G54.2	Cervical root disorders, not elsewhere classified
G54.3	Thoracic root disorders, not elsewhere classified
M47.21	Other spondylosis with radiculopathy, occipito-atlanto-axial region
M47.22	Other spondylosis with radiculopathy, cervical region
M47.23	Other spondylosis with radiculopathy, cervicothoracic region
M47.24	Other spondylosis with radiculopathy, thoracic region
M50.10	Cervical disc disorder with radiculopathy, unspecified cervical region
M50.11	Cervical disc disorder with radiculopathy, high cervical region
M50.121	Cervical disc disorder at C4-C5 level with radiculopathy
M50.122	Cervical disc disorder at C5-C6 level with radiculopathy
M50.123	Cervical disc disorder at C6-C7 level with radiculopathy
M50.13	Cervical disc disorder with radiculopathy, cervicothoracic region
M51.14	Intervertebral disc disorders with radiculopathy, thoracic region
M54.11	Radiculopathy, occipito-atlanto-axial region
M54.12	Radiculopathy, cervical region
M54.13	Radiculopathy, cervicothoracic region

Diagnosis Code	Description
Cervical/Thoracic	
M54.14	Radiculopathy, thoracic region
M54.15	Radiculopathy, thoracolumbar region
S24.2XXA	Injury of nerve root of thoracic spine, initial encounter
Lumbar/Sacral	
G54.4	Lumbosacral root disorders, not elsewhere classified
M47.26	Other spondylosis with radiculopathy, lumbar region
M47.27	Other spondylosis with radiculopathy, lumbosacral region
M47.28	Other spondylosis with radiculopathy, sacral and sacrococcygeal region
M48.062	Spinal stenosis, lumbar region with neurogenic claudication
M51.16	Intervertebral disc disorders with radiculopathy, lumbar region
M51.17	Intervertebral disc disorders with radiculopathy, lumbosacral region
M54.16	Radiculopathy, lumbar region
M54.17	Radiculopathy, lumbosacral region
M54.18	Radiculopathy, sacral and sacrococcygeal region
M54.30	Sciatica, unspecified side
M54.31	Sciatica, right side
M54.32	Sciatica, left side
M54.40	Lumbago with sciatica, unspecified side
M54.41	Lumbago with sciatica, right side
M54.42	Lumbago with sciatica, left side
S34.21XA	Injury of nerve root of lumbar spine, initial encounter
S34.22XA	Injury of nerve root of sacral spine, initial encounter

Description of Services

Spine pain, in particular, pain in the lower back is a common concern, affecting up to 90% of Americans at some point in their lifetime. The majority of episodes are mild and self-limiting, and up to 50% of affected persons will have more than one episode. It is a symptom of a variety of different conditions, including injury, spinal stenosis, disc herniation or degenerative changes in the vertebrae. Epidural steroid injections (ESIs) may be used as a non-surgical modality to treat low back and neck pain, and involve the injection of a solution containing corticosteroids and/or anesthetic into the epidural space. The ESI can be performed via Interlaminar (ILES), transforaminal (TFESI), or caudal approaches (caudal ESI).

Epidural steroid injections generally require local anesthetic only. However, for some patients, Moderate/Conscious Sedation, non-intravenous sedation, and Monitored Anesthesia Care (MAC) may be necessary. These sedation procedures are generally safe when administered by trained, certified providers with appropriate monitoring, but are not without risk. Examples of procedures that typically do not require moderate sedation or an anesthesia care team include but are not limited to epidural steroid injections; epidural blood patch; trigger point injections; shoulder, hip, sacroiliac, facet, and knee joint injections; medial branch nerve blocks; and peripheral nerve blocks (American Society of Anesthesiologists, 2021).

Clinical Evidence

Ultrasound Guidance

There is limited evidence in the peer-reviewed literature demonstrating the overall health benefit of the use of ultrasonic guidance during spinal injections over the use of fluoroscopy or CT-guidance.

Ultrasound-guided spine injection therapy is a comparatively new technique in the management of axial and radicular pain from degenerative lumbar spinal conditions and may be a reasonable alternative to conventional methods of injection guidance. In 2020, Tay et al. completed a retrospective clinical audit of 42 patients who underwent ultrasound-guided lumbar spinal injection at a single institution for chronic axial and radicular pain in an acute public hospital sports medicine center between June 1, 2018 and June 1, 2019. 27 patients (64.3%) receiving facet joint injections and 18 patients (42.9%) receiving nerve root injections. The majority (90.5%) of patients experienced an improvement of > 30% in pain intensity at 3 months post-injection, using the Numerical Rating Scale pain score ($p < 0.001$); with 40 patients (95.2%) reporting a reduction in Oswestry Disability Index score ($p < 0.001$). No complications were reported. It was concluded that the experience of this institution confirms the safety, feasibility, and effectiveness of ultrasound-guided lumbar spinal injection for the treatment of axial and radicular pain. The authors also note that ultrasound-guided spinal injection remains technically challenging and requires a steep learning phase, as well as careful patient selection, and that the study was not designed to directly compare outcomes for ultrasound-guided injection against the conventional standard of care. A larger dataset is required to confirm the efficacy of ultrasound-guided spine injection and the rate of adverse events, and a prospective study would be useful to determine clinical factors predicting success. This study is also limited by lack of comparison group and a small number of participants standard of care. A larger dataset is required to confirm the efficacy of ultrasound-guided spine injection and the rate of adverse events, and a prospective study would be useful to determine clinical factors predicting success. This study is also limited by lack of comparison group and a small number of participants.

Epidural Steroid Injections

Overall, the volume of evidence for the use of therapeutic epidural injections in the treatment of acute and chronic back pain is large. Clinical studies have shown that epidural steroid injections have provided short-term improvement and may be considered in the treatment of selected patients with radicular pain as part of an active therapy program. There is, however, insufficient evidence to demonstrate that epidural steroid injections are effective in the treatment of back pain in the absence of radicular symptoms.

In a 2021 Hayes Evolving Evidence Review regarding epidural steroid injections (ESI) for the treatment of thoracic spine pain, it was concluded that thoracic disc herniation is rare, and patients may present with thoracic axial pain, but no radicular pain. The clinical evidence is limited, and the results of one randomized controlled trial suggests that ESI, either anesthetic alone or anesthetic plus corticosteroid, for chronic thoracic pain in patients who primarily had disc-associated pain provides clinical benefits at up to 2 years.

Helm et al. (2021) conducted a systematic review and meta-analysis of the efficacy and safety of transforaminal epidural steroid injections for 4 indications: radicular pain from spinal stenosis and failed back surgery syndrome; and for axial low back pain. The available literature on transforaminal injections was reviewed and the level of evidence was classified on a 5-point scale based on the quality of evidence developed by the U.S. Preventive Services Task Force (USPSTF) and modified by the American Society of Interventional Pain Physicians (ASIPP). Data sources included relevant literature from 1966 to April 2020, and manual searches of the bibliographies of known primary and review articles. Pain relief and functional improvement were the primary outcome measures. A minimum of 6 months pain relief follow-up was required. Eighteen randomized controlled trials met the inclusion criteria. Eleven randomized controlled trials dealt with various aspects of transforaminal injections for radicular pain due to disc herniation and show Level 1 evidence supporting the use of transforaminal injections for this condition. A meta-analysis showed that at both 3 and 6 months, there was highly statistically significant improvement in both pain and function with both particulate and nonparticulate steroids. For radicular pain from central stenosis, there is one moderate quality study with Level IV evidence. For radicular pain caused by failed back surgery syndrome, there is one moderate quality study with Level IV evidence. For radicular pain from foraminal stenosis and for axial pain, there is Level V evidence, opinion-based/consensus, supporting the use of transforaminal injections. The authors concluded that Level I evidence indicates transforaminal injections are generally safe but have been associated with major neurological complications related to spinal cord infarction. Due to concern over the role of particulate steroids, multiple other injectates have been evaluated, including nonparticulate steroids, tumor necrosis factor alpha (TNF- α) inhibitors, and local anesthetics without steroids, and none have been proven superior. This review is limited by the paucity of literature for some indications.

Verheijen et al. (2021) conducted a systematic review and meta-analysis comparing epidural steroid injections (ESIs) with placebo injections in sciatica patients. The review included a total of 17 out of 732 reports: epidural placebo ($n = 13$), non-epidural placebo ($n = 2$), and both placebo groups ($n = 2$). The primary outcome measures were pooled using a random-effects model for 6-week, 3-month, and 6-month follow-up. Secondary outcomes were described qualitatively. Results showed that ESI was superior compared to epidural placebo at 6 weeks [-8.6 (-13.4; -3.9)] and 3 months [-5.2 (-10.1; -0.2)] for leg pain and at

6 weeks for functional status [-4.1 (-6.5; -1.6)], though the minimally clinical important difference (MCID) was not met. There was no difference in ESI and placebo for back pain, except for non-epidural placebo at 3 months [6.9 (1.3; 12.5)]. Proportions of treatment success were not different. ESI reduced analgesic intake in some studies and complication rates were low. Of the 17 trials, five were considered low risk of bias, two raised some concerns, and 10 studies were considered high-risk. One serious adverse event was documented (retroperitoneal hematoma after ESI) and several minor complications related to needle placement and corticosteroid were noted. Limitations of the review include a low quality of evidence and limited number of comparison studies. The authors concluded that ESIs compared to placebo is considered safe and effective treatment for short-term pain management, however, at three and six months, no proven additional value of ESI compared to placebo was noted.

In a 2020 meta-analysis of randomized controlled trials, Yang et al. compared the clinical effectiveness of epidural steroid injections (ESI) versus conservative treatments for patients with lumbosacral radicular pain. A search was conducted on relevant studies published between 2000 and January 10, 2019 and randomized controlled trials directly comparing the efficacy of ESI with conservative treatment were selected. Primary outcomes included pain relief, functional improvement using the Oswestry Disability Index, or successful events. 6 randomized controlled trials (249 patients with ESI and 241 patients with conservative treatment) were identified and included in this meta-analysis. The results showed that ESI was beneficial for pain relief at short-term (1-3 months) and intermediate-term (3-6 months) when compared with conservative treatment, but this effect was not maintained at long-term (6 months to one year) follow-up. In terms of functional improvement, the overall outcome of meta-analysis showed that ESI did not have any advantage over conservative treatment at short-term and intermediate-term follow-up. Successful event rates were significantly higher in patients who received ESI than in patients who received conservative treatment. There were no statistically significant differences in functional improvement after ESI and conservative treatment at short-term and intermediate-term follow-up. The authors concluded that the use of ESI is more effective for alleviating lumbosacral radicular pain than conservative treatments in terms of short-term and intermediate term. Patients also reported more successful outcomes after receiving ESI when compared to conservative treatment. However, this effect was not maintained at long-term follow-up. The limitations of this meta-analysis resulted from the variation in types of interventions and small sample size.

A 2019 Hayes Health Technology Assessment regarding epidural steroid injections for cervical radiculopathy concluded that the evidence did not demonstrate any beneficial effect of ESIs on pain or disability associated with cervical radiculopathy compared with epidural injection of anesthetic alone. Although complications reported in the reviewed studies were generally mild and transient, serious AEs have occurred, including paraplegia, meningitis, and epidural abscess. Differences, often subtle, in the injection route, region, steroid, anesthetic, and patient pathology result in a vast array of procedural options for ESI, and such variability makes interpretation of existing ESI data difficult.

Smith et al. (2019) published the results of a systematic review of 19 studies assessing the efficacy of lumbar transforaminal steroid injection for radicular pain due to lumbar disc herniation. Placebo controlled RCTs, pragmatic studies, and observational studies were included in the analysis. Utilizing a threshold of $\geq 50\%$ reduction in pain, treatment success rates across studies were 63% (Range: 58 to 68%) at 1-month, 74% (Range: 68-80%) at 3-months, 64% (59-69%) at 6-months, and 64% (57-71%) at 1year. The authors concluded that there is strong evidence that lumbar transforaminal injection of steroids is an effective treatment for radicular pain due to disc herniation.

In a systematic review, Manchikanti et al. (2015) reported on the long-term efficacy of cervical intralaminar and transforaminal epidural injections, focusing on cervical disc herniation, spinal stenosis, and discogenic pain. Based on seven randomized controlled trials of different types of injections, none of which included comparison to a placebo group or to non-invasive treatment, the authors concluded that the evidence demonstrated Level II evidence for efficacy of cervical interlaminar epidural injections, in spite of the scant available clinical literature specific to conditions other than disc herniation. The findings are limited by lack of relevant comparison group that would allow to estimate the benefit of cervical epidural injection as compared to other treatment approaches.

A randomized, double-blind controlled trial was conducted by Manchikanti et al. (2014). The objective of this trial was to assess the effect on pain relief and functional improvement using thoracic interlaminar epidural injections in patients with chronic mid back pain and/or upper back pain. Two groups of 55 patients each were randomized to receive injections with local anesthetic alone, or injections with local anesthetic plus steroids. After two years both groups of patients saw significant improvement (71% using anesthetic alone, and 80% using anesthetic plus steroids). The authors concluded that chronic thoracic pain (not

originating in the facet joint) could be managed with both types of thoracic epidural injections. This study was limited by lack of a placebo group.

Rosas et al. (2010) performed a retrospective case series to evaluate fluoroscopically directed thoracic transforaminal epidural injections. One hundred and ninety-eight foraminal nerve blocks and foraminal epidural injections to the thoracic spine were performed between 1997 and 2007. This new technique was reviewed to evaluate improved safety, as this approach should decrease the chance of inadvertently injuring surrounding structures. There were no major complications when this new technique was properly performed. The authors concluded that this new technique of performing thoracic transforaminal epidural injections under fluoroscopy allows the ability to gauge needle depth, thereby decreasing potential injury to surrounding structures, including the pleura, dura mater, and vasculature. The findings are limited by lack of outcome data other than safety data and lack of comparison group.

Manchikanti et al. (2010) conducted a double-blind randomized controlled trial of interlaminar epidural steroid injections, with and without steroids, in managing chronic pain of lumbar disc herniation or radiculitis. Seventy patients were equally randomized to receive either a local anesthetic only (group I) or a local anesthetic mixed with a steroid (group II). Outcomes were measured at baseline, 3, 6, and 12 months post-treatment with the Numeric Rating Scale (NRS), the Oswestry Disability Index 2.0 (ODI), employment status, and opioid intake. Significant pain relief ($\geq 50\%$) was seen at 12 months in 74% of patients in group I and 86% in group II, and 69% and 83% in ODI scores respectively. Patients in group II also had more improvement in functional status at 12 months (83% vs. 69%) and required less opioid intake.

Cyteval et al. (2006) prospectively followed 229 patients with lumbar radiculopathy (herniated disc and degenerative lesions) at 2 weeks and 1 year after percutaneous periradicular (transforaminal) steroid infiltration. The aim of the study was to find predictive factors of efficacy of the steroid injection procedure. ESIs were performed under fluoroscopic guidance, and periradicular flow was confirmed with contrast medium. Short- and long-term pain relief was demonstrated. The only predictive factor of pain relief was symptom duration before the procedure. The authors concluded that periradicular (transforaminal) infiltration was a simple, safe, and effective (short- and long-term relief) nonsurgical procedure with an improved benefit when performed early in the course of the illness. The primary limitation of the study was the lack of a control group.

A retrospective case series by Botwin et al. (2006) assessed thoracic interlaminar epidural steroid injections, done under fluoroscopy, for the incidence of adverse effects or complications. The study included 21 patients who received the injections over a five-year period who were experiencing thoracic radicular pain from herniated nucleus pulposus or thoracic spondylosis. The authors concluded that there were no major complications, and there was no difference in the complication rate between the two diagnoses. The findings are limited by lack of outcome data other than safety data and lack of comparison group.

Complications associated with epidural injections include steroid side effects, dural puncture, transient increased pain, transient paresthesias, aseptic and/or bacterial meningitis, neurological dysfunction or damage, epidural abscess, intracranial air, allergic reaction, epidural hematoma, persistent dural leak, nausea, headache, paraplegia, tetraplegia, seizure, stroke, and death. (Everett, 2004)

Clinical Practice Guidelines

American College of Occupational and Environmental Medicine (ACOEM)

In the 2021 guidelines for invasive treatments for low back disorders, the ACOEM state the following regarding epidural steroid injections (ESI):

- Recommended (I), Moderate Confidence for select circumstances as an option for treatment of acute or subacute radicular pain syndromes, typically after treatment with NSAID and waiting at least 3 weeks
- Moderately Not Recommended (B), Moderate Confidence for treatment of spinal stenosis
- Not Recommended, Evidence (C), High Confidence for treatment of acute, subacute, or chronic low back pain in the absence of significant radicular symptoms

Agency for Healthcare Research and Quality (AHRQ) Technology Assessment Program

The 2015 AHRQ comparative effectiveness study on injection therapies for low back pain (LBP) concluded that ESIs for radiculopathy were associated with immediate improvements in pain and might be associated with immediate improvements in function, but benefits were small and not sustained, and there was no effect on long-term risk of surgery. Evidence did not suggest that effectiveness varies based on injection technique, corticosteroid, dose, or comparator. Limited evidence

suggested that epidural corticosteroid injections are not effective for spinal stenosis or nonradicular back pain. (Chou et al. 2015)

American Society of Anesthesiologists (ASA)

As of 2010, the ASA had not issued a statement specifically on the use of epidural steroids for the management of low back pain and/or sciatica. However, the ASA Task Force on Pain Management issued more general practice guidelines for chronic pain management. The 2010 ASA guidelines recommended that: epidural steroid injections with or without local anesthetics may be used as part of a multimodal treatment regimen to provide pain relief in selected patients with radicular pain or radiculopathy. Transforaminal epidural injections should be performed with appropriate image guidance to confirm correct needle position and spread of contrast before injecting a therapeutic substance.

American Academy of Neurology (AAN)

In 2007, and reaffirmed in 2010, the Therapeutics and Technology Assessment Subcommittee of the AAN released an assessment addressing the use of epidural steroid injections (ESIs) to treat radicular lumbosacral pain:

- Epidural steroid injections may result in some improvement in radicular lumbosacral pain when determined between 2 and 6 weeks following the injection, compared to control treatment (Level C, Class I to III evidence). The average magnitude of effect is small, and the generalizability of the observation is limited by the small number of studies, limited to highly selected patient populations, the few techniques and doses studied, and variable comparison treatments.
- In general, epidural steroid injections for radicular lumbosacral pain have shown no impact on average impairment of function, on need for surgery, or on long-term pain relief beyond 3 months. Their routine use for these indications is not recommended (Level B, Class I to III evidence).
- Data on use of epidural steroid injections to treat cervical radicular pain are inadequate to make any recommendation (Level U).

American Society of Interventional Pain Physicians (ASIPP)

The ASIPP published updated evidence-based guidelines regarding epidural interventional techniques in the management of chronic spinal pain in 2021 (Manchikanti et al.). The ASIPP maintains a comprehensive guideline for epidural steroid injections including indications, limitations and therapy frequencies. Specifically, these guidelines make the following conclusion or recommendations, among others:

- Disc herniation: based on relevant, high-quality fluoroscopically-guided epidural injections, with or without steroids, and results of previous systematic reviews, the evidence is Level I for caudal epidural injections, lumbar interlaminar epidural injections, lumbar transforaminal epidural injections, and cervical interlaminar epidural injections with strong recommendation for long-term effectiveness.
- For thoracic disc herniation, based on one relevant, high-quality RCT of thoracic epidural with fluoroscopic guidance, with or without steroids, the evidence is Level II with moderate to strong recommendation for long-term effectiveness.
- Spinal stenosis: the evidence based on one high-quality RCT in each category the evidence is Level III to II for fluoroscopically guided caudal epidural injections with moderate to strong recommendation and Level II for fluoroscopically guided lumbar and cervical interlaminar epidural injections with moderate to strong recommendation for long-term effectiveness.
- The evidence for lumbar transforaminal epidural injections is Level IV to III with moderate recommendation with fluoroscopically guided lumbar transforaminal epidural injections for long-term improvement.
- Axial discogenic pain: the evidence for axial discogenic pain without facet joint pain or sacroiliac joint pain in the lumbar and cervical spine with fluoroscopically guided caudal, lumbar and cervical interlaminar epidural injections, based on one relevant high quality RCT in each category is Level II with moderate to strong recommendation for long-term improvement, with or without steroids.
- Post-surgery syndrome: the evidence for lumbar and cervical post-surgery syndrome based on one relevant, high-quality RCT with fluoroscopic guidance for caudal and cervical interlaminar epidural injections, with or without steroids, is Level II with moderate to strong recommendation for long-term improvement.

The authors also observe that in “the therapeutic phase (after the diagnostic phase is completed), the suggested frequency of interventional techniques should be 2½ to 3 months or longer between each injection, provided that > 50% relief is obtained for 2½ to 3 months, not exceeding 4 per year, per region.”

American Association of Neurological Surgeons and the Congress of Neurological Surgeons

In a 2014 joint guideline update, the American Association of Neurological Surgeons and the Congress of Neurological Surgeons states that the published medical literature continues to fail to support the use of lumbar epidural injections for long-term relief of chronic back pain without radiculopathy, and that there is limited support for their use for short-term relief in selected patients with chronic back pain (Watters, 2014).

North American Spine Society (NASS)

In 2020, NASS revised its coverage policy recommendations for epidural steroid injections and selective spinal nerve blocks. They stated that the rationale for coverage is based on high-level evidence and what most practitioners would consider to be accepted practice patterns. Multiple randomized-controlled trials (RCTs) have demonstrated that lumbar epidural steroid injections (LESIs) are effective in the treatment of lumbar radiculitis caused by disc herniation. There is sufficient literature to suggest that a trial of ESIs for radicular pain caused by conditions other than disc herniation is appropriate prior to considering surgical intervention.

In their 2020 Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis & Treatment of Low Back Pain, the NASS states there is insufficient evidence to make a recommendation for or against the use of caudal epidural steroid injections and interlaminar epidural steroid injections in patients with low back pain.

In 2013 NASS published a Review and Recommendation Statement entitled “Lumbar Transforaminal Epidural Steroid Injections”. A grade A recommendation (defined as good evidence) was given for the effectiveness of ESI at treating radicular pain related to lumbar disc herniation for at least 1 month in more than 50% of individuals. The review graded the evidence as insufficient for a recommendation to treat lumbar radicular pain in the presence of stenosis. There was insufficient evidence to provide an evidence-based recommendation on the maximum number of lumbar ESIs that are appropriate in any given timeframe or the amount of pain/functional improvement needed to justify repeat injections.

In the 2010 Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis & Treatment of Cervical Radiculopathy from Degenerative Disorders, NASS states that transforaminal epidural steroid injections using fluoroscopic or CT guidance may be considered when developing a medical/interventional treatment plan for patients with cervical radiculopathy from degenerative disorders, and that consideration should be given to the potential complications before performing this procedure.

In 2011, NASS revised its Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis with the following recommendation: while there is evidence that nonfluoroscopically guided interlaminar and single radiographically guided transforaminal ESIs can result in short-term symptom relief in patients with neurogenic claudication or radiculopathy, there is conflicting evidence concerning long-term efficacy. The guidelines also note that there is some evidence that a multiple injection regimen of radiographically guided transforaminal ESIs or caudal injections can produce long-term relief of pain in patients with radiculopathy or neurogenic intermittent claudication from lumbar spinal stenosis. However, the evidence is of relatively poor quality, and therefore no strong recommendation in support of this therapy was made.

World Federation of Neurosurgical Societies (WFNS)

In 2020, the WFNS published the Spine Committee Recommendations on Conservative Treatment and Percutaneous Pain Relief in Patients with Lumbar Spinal Stenosis (Fornari et al. 2020). The following recommendations were made:

Conservative Treatment or Follow-Up for Lumbar Spinal Stenosis (LSS)

- In nonsevere clinical conditions a conservative approach based on at least 3 weeks of therapeutic exercise may be the first therapeutic choice
- Medical/interventional treatment should be preferred to surgical treatment in patients with spinal stenosis with mild symptoms
- Physical therapy should consist of multimodal approaches
- If conservative treatment is chosen, surgery should be considered in cases in which the clinical condition does not change in 3 months
- There are some cases in which immediate surgical treatment can be indicated

Percutaneous Pain Relief Techniques for LSS

- The literature supports short- to intermediate-term benefits of epidural injections for symptomatic treatment of LSS
- Inclusion of steroids does not confer a benefit compared with local anesthetic alone in epidural injections for symptomatic treatment of LSS
- For patients with symptomatic relief lasting < 3 months after epidural injections, proceeding with further injections is not recommended

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Epidural Steroid Injection is a procedure and, therefore, not subject to FDA regulation. However, any medical devices, drugs, biologics, or tests used as a part of this procedure may be subject to FDA regulation. Injectable corticosteroids include methylprednisolone, hydrocortisone, triamcinolone, betamethasone, and dexamethasone, and are approved by the FDA, however, the effectiveness and safety of the drugs for Epidural Steroid Injection have not been established, and FDA has not approved corticosteroids for such use. Additional information may be obtained from the U.S. Food and Drug Administration - Center for Drug Evaluation and Research (CDER) at: <https://www.fda.gov/about-fda/fda-organization/center-drug-evaluation-and-research-cder> (Accessed January 10, 2022)

In April 2014, the U.S. Food and Drug Administration (FDA) warned, that injection of corticosteroids into the epidural space of the spine may result in rare but serious adverse events, including loss of vision, stroke, paralysis, and death. They noted the effectiveness and safety of epidural administration of corticosteroids have not been established, and the FDA has not approved corticosteroids for this use. FDA is requiring the addition of a warning to the drug labels of injectable corticosteroids to describe these risks. The FDA recommends that individuals should discuss the benefits and risks of epidural corticosteroid injections with their health care professionals, along with the benefits and risks associated with other possible treatments. Further information can be found at: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-requires-label-changes-warn-rare-serious-neurologic-problems-after> (Accessed January 10, 2022)

References

- American Academy of Neurology (AAN). Summary of Evidence-based Guideline for Clinicians: Use of Epidural Steroid Injections to Treat Radicular Lumbosacral Pain. 2007. Reaffirmed in 2010.
- American College of Occupational and Environmental Medicine Guidelines. Invasive Treatments for Low Back Disorders. 2021.
- American Society of Anesthesiologists (ASA). Committee on Pain Medicine. Statement on Anesthetic Care During Interventional Pain Procedures for Adults. October 2005; Amended October 2021. <https://www.asahq.org/standards-and-guidelines/statement-on-anesthetic-care-during-interventional-pain-procedures-for-adults>. Accessed March 1, 2022.
- American Society of Anesthesiologists (ASA). Practice Guidelines for Chronic Pain Management: An Updated Report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. Anesthesiology 2010.
- Benyamin RM, et al. The effectiveness of lumbar interlaminar epidural injections in managing chronic low back and lower extremity pain. Pain Physician 2012.
- Botwin KP, Baskin M, Rao S. Adverse effects of fluoroscopically guided interlaminar thoracic epidural steroid injections. *Am J Phys Med Rehabil*. 2006;85(1):14-23. doi:10.1097/01.phm.0000184475.44853.82.
- Chou R, Hashimoto R, Friedly J, et al. Pain Management Injection Therapies for Low Back Pain. Rockville (MD): Agency for Healthcare Research and Quality 2015.
- Cyteval C, Fescquet N, Thomas E, et al. Predictive factors of efficacy of periradicular corticosteroid injections for lumbar radiculopathy. *AJNR Am J Neuroradiol*. 2006.
- Everett CR, Baskin MN, Novoseletsky D, et al. Flushing as a side effect following lumbar transforaminal epidural steroid injection. Pain Physician. 2004.
- Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine* 2000 Nov 15;25(22):2940-52.

Fornari M, Robertson SC, Pereira P, et al. Conservative treatment and percutaneous pain relief techniques in patients with lumbar spinal stenosis: WFNS Spine Committee Recommendations. *World Neurosurg* X. 2020 Jun 23;7:100079.

Goertz M, Thorson D, Bonsell J, et al. Institute for Clinical Systems Improvement. Adult Acute and Subacute Low Back Pain. Updated November 2012.

Hayes, Inc. Health Technology Assessment. Epidural Steroid Injections For Cervical Radiculopathy. Landsdale, PA: February 28, 2019.

Hayes, Inc. Evolving Evidence Review. Epidural Steroid Injections for the Treatment of Thoracic Spine Pain. Lansdale, PA: July 23, 2021.

Helm li S, Harmon PC, Noe C, et al. Transforaminal Epidural Steroid Injections: A Systematic Review and Meta-Analysis of Efficacy and Safety. *Pain Physician*. 2021 Jan;24(S1): S209-S232.

Lenahan et al. Current Guidelines for Management of Low Back Pain. *Clinical Advisor*. 2018.

Manchikanti L, Boswell MV, Singh V, et al. Comprehensive evidence-based guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician*. 2009.

Manchikanti L, Cash KA, Pampati V, et al. Transforaminal epidural injections in chronic lumbar disc herniation: a randomized, double-blind, active-control trial. *Pain Physician*. 2014.

Manchikanti L, Cash KA, McManus CD, Pampati V, Benyamin RM. Thoracic interlaminar epidural injections in managing chronic thoracic pain: a randomized, double-blind, controlled trial with a 2-year follow-up. *Pain Physician*. 2014;17(3):E327-E338.

Manchikanti L, Nampiaparampil DE, Candido KD, et al. Do cervical epidural injections provide long-term relief in neck and upper extremity pain? A systematic review. *Pain Physician*. 2015;18(1):39-60.

Manchikanti L, Knezevic NN, Navani A, et al. Epidural Interventions in the Management of Chronic Spinal Pain: American Society of Interventional Pain Physicians (ASIPP) Comprehensive Evidence-Based Guidelines. *Pain Physician*. 2021 Jan;24(S1): S27-S208.

Manchikanti L, Singh V, Falco FJ, et al. Evaluation of the effectiveness of lumbar interlaminar epidural injections in managing chronic pain of lumbar disc herniation or radiculitis: a randomized, double-blind, controlled trial. *Pain Physician*. 2010.

Mattie R, Schneider BJ, Smith C. Frequency of Epidural Steroid Injections. *Pain Med*. 2020 May 1;21(5):1078-1079.

North American Spine Society (NASS). Evidence Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Low Back Pain. 2020.

North American Spine Society (NASS). Evidence- Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis (Revised 2011).

North American Spine Society (NASS). Evidence- based Clinical Guidelines for Multidisciplinary Spine Care. Diagnosis and Treatment of Cervical Radiculopathy from Degenerative Disorders. 2010.

North American Spine Society (NASS). Lumbar transforaminal epidural steroid injections: Review and recommendation statement. 2013. <https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/LTFESIRReviewRecStatement.pdf>. (Accessed January 10, 2022).

Patel K, Chopra P, Upadhyayula S. Epidural Steroid Injections. [Updated 2021 Jul 19]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470189/>.

Qassem, Amir, et al. Noninvasive treatments for acute, subacute and chronic low back pain: A clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*. April 2017.

Rosas, HG, Gilula, LA. Performing thoracic transforaminal injections: A new technique. *Radiology*, Vol 254, No. 2, January 7, 2010.

Smith CC, McCormick ZL, Mattie R et al. The effectiveness of lumbar transforaminal injection of steroid for the treatment of radicular pain. *Pain Med*, July, 2019.

Staehler, R. Epidural Steroid Injection Pain Relief Success Rates [Internet]. Deerfield, IL: Veritas Health. Spine-health; 2020 Oct 22. Available from: https://www.spine-health.com/treatment/injections/epidural-steroid-injection-pain-relief-success-rates#vh_footnotes.

Staehler, R. Lumbar Epidural Steroid Injections for Low Back Pain and Sciatica [Internet]. Deerfield, IL: Veritas Health. Spine-health, 2020 Oct 22. Available from: <https://www.spine-health.com/treatment/injections/lumbar-epidural-steroid-injections-low-back-pain-and-sciatica>.

Summers, Jeffrey. International Spine Intervention Society Recommendations for treatment of Cervical and Lumbar Spine Pain. 2013.

Tay M, Sian SCSH, Eow CZ, et al. Ultrasound-guided lumbar spine injection for axial and radicular pain: A Single Institution Early Experience. Asian Spine J. 2020 Sep 3.

Verheijen EJA, Bonke CA, Amorij EMJ, et al. Epidural steroid compared to placebo injection in sciatica: a systematic review and meta-analysis. Eur Spine J. 2021 Nov;30(11):3255-3264.

Watters WC 3rd, Resnick DK, Eck JC, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 13: injection therapies, low-back pain, and lumbar fusion. J Neurosurg Spine. 2014 Jul;21(1):79-90.

Yang S, Kim W, Kong HH, et al. Epidural steroid injection versus conservative treatment for patients with lumbosacral radicular pain: A meta-analysis of randomized controlled trials. Medicine (Baltimore). 2020 Jul 24;99(30).

Policy History/Revision Information

Date	Summary of Changes
08/01/2022	<p>Coverage Rationale</p> <ul style="list-style-type: none"> ● Replaced language indicating “Epidural Steroid Injections (ESI) are proven and medically necessary <i>for treating</i> radicular pain <i>caused by spinal stenosis, disc herniation, degenerative changes in the vertebrae, or</i> for the short-term management of <i>spine</i> pain” with “Epidural Steroid Injections (ESI) are proven and medically necessary <i>when the injection is intended</i> for the short term management of <i>acute or subacute</i> radicular pain” ● Revised coverage criteria: <ul style="list-style-type: none"> ○ Removed criterion requiring the pain is associated with symptoms of nerve root irritation and/or spine pain due to disc extrusions and/or contained herniations ○ Replaced criterion requiring “the pain is unresponsive to Conservative Treatment, <i>including but not limited to</i> pharmacotherapy, exercise, or physical therapy” with “the <i>radicular</i> pain is unresponsive to Conservative Treatment: pharmacotherapy <i>such as NSAIDS or acetaminophen</i> ≥ 3 weeks; activity modification ≥ 4 weeks (<i>including but not limited to heavy lifting, bending, spinal torsion activities</i>); or physical therapy (PT) <i>or home exercise</i> ≥ 4 weeks” <p>Epidural Steroid Injection Limitations</p> <ul style="list-style-type: none"> ● Added language to indicate repeat ESIs may be provided only if: <ul style="list-style-type: none"> ○ The initial injection resulted in ≥ 50% pain relief achieved for 3 or more months ○ The initial injection resulted in a functional improvement as measured by validated measurement tools, such as the <i>Oswestry Disability Index</i> ○ Repeat injections do not exceed 3 per year, per region ● Replaced language indicating: <ul style="list-style-type: none"> ○ “A maximum of three (3) ESI sessions (per region, regardless of level, location, or side) [may be provided] <i>in a year when criteria (indications for coverage) are met for each injection</i>” with “a maximum of three (3) ESI sessions (per region, regardless of level, location, or side) [may be provided] <i>per year</i>” ○ “A region is defined by either the region of the cervical or thoracic spine or the region of the lumbar or sacral spine” with “a region is defined by the region of the cervical, thoracic, or lumbosacral spine” <p>Definitions</p> <ul style="list-style-type: none"> ● Added definition of “Oswestry Disability Index (also known as the Oswestry Low Back Pain Disability Questionnaire)” ● Updated definition of “Epidural Steroid Injections (ESIs)”

Date	Summary of Changes
	<p data-bbox="337 138 639 170">Supporting Information</p> <ul data-bbox="337 176 1455 268" style="list-style-type: none"> <li data-bbox="337 176 1455 237">• Updated <i>Description of Services</i>, <i>Clinical Evidence</i>, <i>FDA</i>, and <i>References</i> sections to reflect the most current information <li data-bbox="337 243 854 268">• Archived previous policy version CS039.T

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state, or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state, or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state, or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state, or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.