

Molecular Oncology Testing for Cancer Diagnosis, Prognosis, and Treatment Decisions (for Indiana Only)

Policy Number: CS152IN.03
Effective Date: October 1, 2021

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	3
Applicable Codes	3
Description of Services	7
Clinical Evidence	8
U.S. Food and Drug Administration	29
References	30
Policy History/Revision Information	37
Instructions for Use	37

Related Policy

- [Chemosensitivity and Chemoresistance Assays in Cancer \(for Indiana Only\)](#)

Application

This Medical Policy only applies to the state of Indiana.

Coverage Rationale

Breast Cancer

The use of Oncotype Dx Breast and EndoPredict gene assay are proven and medically necessary in certain circumstances. For medical necessity clinical criteria, refer to the [Indiana Genetic Testing Provider Reference Module](#) and [Indiana Health Coverage Programs \(IHCP\) February 11, 2020 bulletin](#).

Thyroid Cancer

Molecular profiling of thyroid nodules (e.g., Afirma GSC, ThyroSeq V3, ThyGeNEXT/ThyraMIR, or the gene and gene fusion panel *BRAF*, *RAS*, *HRAS*, *NRAS*, *RET/PTC1*, *RET/PTC3*, *PAX8/PPARγ*) is proven and medically necessary when all the following criteria are met:

- Follicular pathology on fine needle aspiration is indeterminate; and
- The results of the test will be used for making decisions about further surgery

Molecular profiling of thyroid nodules or thyroid cancers is unproven and not medically necessary for all other indications due to insufficient evidence of efficacy.

Use of more than one molecular profile test in an individual with a thyroid nodule is unproven and not medically necessary due to insufficient evidence of efficacy.

Hematological Cancer

Molecular profiling using Chromosomal Microarray analysis (e.g., Oncoscan, Reveal SNP-Oncology, CGH or SNP array) is proven and medically necessary for individuals with acute leukemia.

Use of a Next Generation Sequencing profile test to assess minimal residual disease (e.g., ClonoSeq) is proven and medically necessary for individuals with multiple myeloma when the following criteria are met:

- Individual had an allogenic or autologous bone marrow transplant for multiple myeloma; and
- Within 3 months of completing a treatment; and
- Has no evidence of progression

All other multigene, gene expression or microarray molecular profiling for hematological malignancies is unproven and not medically necessary due to insufficient evidence of efficacy.

This includes, but is not limited to the following:

- Assessment of minimal residual disease by Next Generation Sequencing for acute myeloid leukemia
- Use of multi-gene Next Generation Sequencing gene panels for predicting prognosis

Lung Cancer

Multigene molecular profiling of non-small cell lung cancer is proven and medically necessary when all of the following criteria are met:

- The panel selected has no more than 50 genes; and
- No prior molecular profiling has been performed on the same tumor; and
- Individual and treating physician have had a discussion prior to testing regarding the potential results of the test and determined to use the results to guide therapy

Liquid biopsy (circulating tumor cell free DNA) molecular profiling tests for non-small cell lung cancer are proven and medically necessary when the following criteria is met:

- The test selected has no more than 50 genes; and
- No prior molecular profiling has been performed on the same tumor; and
- The individual is not medically fit for invasive biopsy; or
- Non-small cell lung cancer has been pathologically confirmed, but there is insufficient material available for molecular testing; and
- Individual and treating physician have had a discussion prior to testing regarding the potential results of the test and determined to use the results to guide therapy

Liquid biopsy (circulating tumor cell free DNA or circulating tumor cells) for any other tumor genetic analysis or tumor screening (e.g., Guardant, Colosentry, epi ProColon, OncoCEE CTC) is unproven and not medically necessary due to insufficient evidence of efficacy.

Due to insufficient evidence of efficacy, molecular profiling using gene expression profiling, Chromosome Microarray multi-gene cancer panels are unproven and not medically necessary for all other indications, including but not limited to:

- Bladder cancer (e.g., Decipher Bladder) (NCCN, 2019a)
- Cancers of unknown primary site (e.g., Response Dx, CancerTYPE ID, Rosetta Cancer Origin, ProOnc, SourceDX, Pathfinder TG)
- Colorectal cancer (e.g., Oncotype DX Colon Cancer Assay, Colorectal Cancer DSA, GeneFx Colon, OncoDefender-CRC)
- Gene panels of >50 genes
- Leukemia other than Chromosome Microarray (e.g., FoundationOne® Heme)
- Melanoma (e.g., Decision Dx – Melanoma, Decision Dx-UM, DermTech PLA)
- Multiple myeloma (e.g., MyPRS/MyPRS Plus)
- Prostate cancer (e.g., Oncotype DX Prostate Cancer Assay, TMPRSS2 fusion gene, Prolaris Prostate Cancer Test, Decipher Prostate Cancer Classifier)
- Uveal melanoma (e.g., Decision Dx-UM)
- Whole Exome Sequencing (WES) and Whole Genomic Sequencing (WGS) of tumors

Definitions

Chromosome Microarray: A laboratory analysis that identifies genome wide copy number variations at the chromosome level, such as aneuploidies, microdeletions and duplications, rearrangements, and amplification. CGH is one technology that can be used for a Chromosome Microarray test, and another example is a single nucleotide polymorphism (SNP) array (Peterson et al., 2018).

Comparative Genome Hybridization (CGH): CGH is a technology that can be used for the detection of genomic copy number variations (CNVs). Tests can use a variety of probes or single nucleotide polymorphisms (SNPs) to provide copy number and gene differentiating information. All platforms share in common that tumor (patient) and reference DNA are labelled with dyes or fluorescing probes and hybridized on the array, and a scanner measures differences in intensity between the probes, and the data is expressed as having greater or less intensity than the reference DNA (Cooley et al; 2013).

Gene Expression Testing: A laboratory test that analyzes mRNA patterns to determine gene activity (Kim et al. 2010).

Next Generation Sequencing (NGS): New sequencing techniques that can quickly analyze multiple sections of DNA at the same time. Older forms of sequencing could only analyze one section of DNA at once (Kamps, et al. 2017).

Whole Exome Sequencing (WES): About 1% of a person's DNA makes protein. These protein making sections are called exons. All the exons together are called the exome. WES is a DNA analysis technique that looks at all of the exons in a person, or a tissue type such as a tumor, at one time, rather than gene by gene (U.S. National Library of Medicine, 2017A).

Whole Genome Sequencing (WGS): WGS determines the sequence of the entire DNA in a person, or a tissue type, such as a tumor, which includes the protein making (coding) as well as non-coding DNA elements (U.S. National Library of Medicine, 2017B).

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
0005U	Oncology (prostate) gene expression profile by real-time RT-PCR of 3 genes (ERG, PCA3, and SPDEF), urine, algorithm reported as risk score
0011M	Oncology, prostate cancer, mRNA expression assay of 12 genes (10 content and 2 housekeeping), RT-PCR test utilizing blood plasma and urine, algorithms to predict high-grade prostate cancer risk
0012M	Oncology (urothelial), mRNA, gene expression profiling by real-time quantitative PCR of five genes (MDK, HOXA13, CDC2 [CDK1], IGFBP5, and CXCR2), utilizing urine, algorithm reported as a risk score for having urothelial carcinoma
0013M	Oncology (urothelial), mRNA, gene expression profiling by real-time quantitative PCR of five genes (MDK, HOXA13, CDC2 [CDK1], IGFBP5, and CXCR2), utilizing urine, algorithm reported as a risk score for having recurrent urothelial carcinoma
0013U	Oncology (solid organ neoplasia), gene rearrangement detection by whole genome next-generation sequencing, DNA, fresh or frozen tissue or cells, report of specific gene rearrangement(s)
0014U	Hematology (hematolymphoid neoplasia), gene rearrangement detection by whole genome next-generation sequencing, DNA, whole blood or bone marrow, report of specific gene rearrangement(s)
0016M	Oncology (bladder), mRNA, microarray gene expression profiling of 209 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as molecular subtype (luminal, luminal infiltrated, basal, basal claudin-low, neuroendocrine-like)

CPT Code	Description
0017M	Oncology (diffuse large B-cell lymphoma [DLBCL]), mRNA, gene expression profiling by fluorescent probe hybridization of 20 genes, formalin-fixed paraffin-embedded tissue, algorithm reported as cell of origin
0018U	Oncology (thyroid), microRNA profiling by RT-PCR of 10 microRNA sequences, utilizing fine needle aspirate, algorithm reported as a positive or negative result for moderate to high risk of malignancy
0019U	Oncology, RNA, gene expression by whole transcriptome sequencing, formalin-fixed paraffin embedded tissue or fresh frozen tissue, predictive algorithm reported as potential targets for therapeutic agents
0021U	Oncology (prostate), detection of 8 autoantibodies (ARF 6, NKX3-1, 5'-UTR-BMI1, CEP 164, 3'-UTR-Ropporin, Desmocollin, AURKAIP-1, CSNK2A2), multiplexed immunoassay and flow cytometry serum, algorithm reported as risk score
0022U	Targeted genomic sequence analysis panel, non-small cell lung neoplasia, DNA and RNA analysis, 23 genes, interrogation for sequence variants and rearrangements, reported as presence/absence of variants and associated therapy(ies) to consider
0026U	Oncology (thyroid), DNA and mRNA of 112 genes, next-generation sequencing, fine needle aspirate of thyroid nodule, algorithmic analysis reported as a categorical result ("Positive, high probability of malignancy" or "Negative, low probability of malignancy")
0036U	Exome (i.e., somatic mutations), paired formalin-fixed paraffin-embedded tumor tissue and normal specimen, sequence analyses
0037U	Targeted genomic sequence analysis, solid organ neoplasm, DNA analysis of 324 genes, interrogation for sequence variants, gene copy number amplifications, gene rearrangements, microsatellite instability and tumor mutational burden
0045U	Oncology (breast ductal carcinoma in situ), mRNA, gene expression profiling by real-time RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence score
0047U	Oncology (prostate), mRNA, gene expression profiling by real-time RT-PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a risk score
0048U	Oncology (solid organ neoplasia), DNA, targeted sequencing of protein-coding exons of 468 cancer-associated genes, including interrogation for somatic mutations and microsatellite instability, matched with normal specimens, utilizing formalin-fixed paraffin-embedded tumor tissue, report of clinically significant mutation(s)
0050U	Targeted genomic sequence analysis panel, acute myelogenous leukemia, DNA analysis, 194 genes, interrogation for sequence variants, copy number variants or rearrangements
0056U	Hematology (acute myelogenous leukemia), DNA, whole genome next-generation sequencing to detect gene rearrangement(s), blood or bone marrow, report of specific gene rearrangement(s)
0069U	Oncology (colorectal), microRNA, RT-PCR expression profiling of miR-31-3p, formalin-fixed paraffin-embedded tissue, algorithm reported as an expression score
0089U	Oncology (melanoma) gene expression profiling by RTqPCR PRAME and LINC00518 superficial collection using adhesive patch(es)
0090U	Oncology (cutaneous melanoma) mRNA gene expression profiling by RT-PCR of 23 genes (14 content and 9 housekeeping) utilizing formalin-fixed paraffin-embedded tissue algorithm reported as a categorical result (i.e., benign indeterminate malignant)
0091U	Oncology (colorectal) screening cell enumeration of circulating tumor cells utilizing whole blood algorithm for the presence of adenoma or cancer reported as a positive or negative result
0113U	Oncology (prostate), measurement of PCA3 and TMPRSS2-ERG in urine and PSA in serum following prostatic massage, by RNA amplification and fluorescence-based detection, algorithm reported as risk score
0118U	Transplantation medicine, quantification of donor-derived cell-free DNA using whole genome next-generation sequencing, plasma, reported as percentage of donor-derived cell-free DNA in the total cell-free DNA

CPT Code	Description
0153U	Oncology (breast), mRNA, gene expression profiling by next-generation sequencing of 101 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a triple negative breast cancer clinical subtype(s) with information on immune cell involvement
0171U	Targeted genomic sequence analysis panel, acute myeloid leukemia, myelodysplastic syndrome, and myeloproliferative neoplasms, DNA analysis, 23 genes, interrogation for sequence variants, rearrangements and minimal residual disease, reported as presence/absence
0179U	Oncology (non-small cell lung cancer), cell-free DNA, targeted sequence analysis of 23 genes (single nucleotide variations, insertions and deletions, fusions without prior knowledge of partner/breakpoint, copy number variations), with report of significant mutation(s)
0204U	Oncology (thyroid), mRNA, gene expression analysis of 593 genes (including BRAF, RAS, RET, PAX8, and NTRK) for sequence variants and rearrangements, utilizing fine needle aspirate, reported as detected or not detected
0208U	Oncology (medullary thyroid carcinoma), mRNA, gene expression analysis of 108 genes, utilizing fine needle aspirate, algorithm reported as positive or negative for medullary thyroid carcinoma
0211U	Oncology (pan-tumor), DNA and RNA by next-generation sequencing, utilizing formalin-fixed paraffin-embedded tissue, interpretative report for single nucleotide variants, copy number alterations, tumor mutational burden, and microsatellite instability, with therapy association
0239U	Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free DNA, analysis of 311 or more genes, interrogation for sequence variants, including substitutions, insertions, deletions, select rearrangements, and copy number variations
0242U	Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free circulating DNA analysis of 55-74 genes, interrogation for sequence variants, gene copy number amplifications, and gene rearrangements
0244U	Oncology (solid organ), DNA, comprehensive genomic profiling, 257 genes, interrogation for single-nucleotide variants, insertions/deletions, copy number alterations, gene rearrangements, tumor-mutational burden and microsatellite instability, utilizing formalin-fixed paraffin-embedded tumor tissue
0245U	Oncology (thyroid), mutation analysis of 10 genes and 37 RNA fusions and expression of 4 mRNA markers using next-generation sequencing, fine needle aspirate, report includes associated risk of malignancy expressed as a percentage
0250U	Oncology (solid organ neoplasm), targeted genomic sequence DNA analysis of 505 genes, interrogation for somatic alterations (SNVs [single nucleotide variant], small insertions and deletions, one amplification, and four translocations), microsatellite instability and tumor-mutation burden
0262U	Oncology (solid tumor), gene expression profiling by real-time RT-PCR of 7 gene pathways (ER, AR, PI3K, MAPK, HH, TGFB, Notch), formalin-fixed paraffin-embedded (FFPE), algorithm reported as gene pathway activity score
81228	Cytogenomic constitutional (genome-wide) microarray analysis; interrogation of genomic regions for copy number variants (e.g., bacterial artificial chromosome [BAC] or oligo-based comparative genomic hybridization [CGH] microarray analysis)
81229	Cytogenomic constitutional (genome-wide) microarray analysis; interrogation of genomic regions for copy number and single nucleotide polymorphism (SNP) variants for chromosomal abnormalities
81277	Cytogenomic neoplasia (genome-wide) microarray analysis, interrogation of genomic regions for copy number and loss-of-heterozygosity variants for chromosomal abnormalities
81425	Genome (e.g., unexplained constitutional or heritable disorder or syndrome); sequence analysis
81426	Genome (e.g., unexplained constitutional or heritable disorder or syndrome); sequence analysis, each comparator genome (e.g., parents, siblings) (List separately in addition to code for primary procedure)
81427	Genome (e.g., unexplained constitutional or heritable disorder or syndrome); re-evaluation of previously obtained genome sequence (e.g., updated knowledge or unrelated condition/syndrome)

CPT Code	Description
81445	Targeted genomic sequence analysis panel, solid organ neoplasm, DNA analysis, and RNA analysis when performed, 5-50 genes (e.g., ALK, BRAF, CDKN2A, EGFR, ERBB2, KIT, KRAS, NRAS, MET, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, if performed
81450	Targeted genomic sequence analysis panel, hematolymphoid neoplasm or disorder, DNA analysis, and RNA analysis when performed, 5-50 genes (e.g., BRAF, CEBPA, DNMT3A, EZH2, FLT3, IDH1, IDH2, JAK2, KRAS, KIT, MLL, NRAS, NPM1, NOTCH1), interrogation for sequence variants, and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed
81455	Targeted genomic sequence analysis panel, solid organ or hematolymphoid neoplasm, DNA analysis, and RNA analysis when performed, 51 or greater genes (e.g., ALK, BRAF, CDKN2A, CEBPA, DNMT3A, EGFR, ERBB2, EZH2, FLT3, IDH1, IDH2, JAK2, KIT, KRAS, MLL, NPM1, NRAS, MET, NOTCH1, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, if performed
81479	Unlisted molecular pathology procedure
81504	Oncology (tissue of origin), microarray gene expression profiling of > 2000 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as tissue similarity scores
81518	Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 11 genes (7 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithms reported as percentage risk for metastatic recurrence and likelihood of benefit from extended endocrine therapy
81519	Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 21 genes, utilizing formalin-fixed paraffin embedded tissue, algorithm reported as recurrence score
81520	Oncology (breast), mRNA gene expression profiling by hybrid capture of 58 genes (50 content and 8 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence risk score
81521	Oncology (breast), mRNA, microarray gene expression profiling of 70 content genes and 465 housekeeping genes, utilizing fresh frozen or formalin-fixed paraffin-embedded tissue, algorithm reported as index related to risk of distant metastasis
81522	Oncology (breast), mRNA, gene expression profiling by RT-PCR of 12 genes (8 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk score
81525	Oncology (colon), mRNA, gene expression profiling by real-time RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence score
81540	Oncology (tumor of unknown origin), mRNA, gene expression profiling by real-time RT-PCR of 92 genes (87 content and 5 housekeeping) to classify tumor into main cancer type and subtype, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a probability of a predicted main cancer type and subtype
81541	Oncology (prostate), mRNA gene expression profiling by real-time RT-PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a disease-specific mortality risk score
81542	Oncology (prostate), mRNA, microarray gene expression profiling of 22 content genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as metastasis risk score
81545	Oncology (thyroid), gene expression analysis of 142 genes, utilizing fine needle aspirate, algorithm reported as a categorical result (e.g., benign or suspicious)
81551	Oncology (prostate), promoter methylation profiling by real-time PCR of 3 genes (GSTP1, APC, RASSF1), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a likelihood of prostate cancer detection on repeat biopsy

CPT Code	Description
81552	Oncology (uveal melanoma), mRNA, gene expression profiling by real-time RT-PCR of 15 genes (12 content and 3 housekeeping), utilizing fine needle aspirate or formalin-fixed paraffin-embedded tissue, algorithm reported as risk of metastasis
81599	Unlisted multianalyte assay with algorithmic analysis
86152	Cell enumeration using immunologic selection and identification in fluid specimen (e.g., circulating tumor cells in blood)
86153	Cell enumeration using immunologic selection and identification in fluid specimen (e.g., circulating tumor cells in blood); physician interpretation and report, when required

CPT® is a registered trademark of the American Medical Association

HCPCS Code	Description
G0327	Colorectal cancer screening; blood-based biomarker

Diagnosis Code	Description
C90.10	Plasma cell leukemia not having achieved remission
C90.11	Plasma cell leukemia in remission
C90.12	Plasma cell leukemia in relapse
C91.00	Acute lymphoblastic leukemia not having achieved remission
C91.01	Acute lymphoblastic leukemia, in remission
C91.40	Hairy cell leukemia not having achieved remission
C91.41	Hairy cell leukemia, in remission
C91.42	Hairy cell leukemia, in relapse
C92.02	Acute myeloblastic leukemia, in relapse
C92.40	Acute promyelocytic leukemia, not having achieved remission
C92.41	Acute promyelocytic leukemia, in remission
C92.42	Acute promyelocytic leukemia, in relapse
C92.50	Acute myelomonocytic leukemia, not having achieved remission
C92.51	Acute myelomonocytic leukemia, in remission
C92.52	Acute myelomonocytic leukemia, in relapse
C92.60	Acute myeloid leukemia with 11q23-abnormality not having achieved remission
C92.61	Acute myeloid leukemia with 11q23-abnormality in remission
C92.62	Acute myeloid leukemia with 11q23-abnormality in relapse
C92.A0	Acute myeloid leukemia with multilineage dysplasia, not having achieved remission
C92.A1	Acute myeloid leukemia with multilineage dysplasia, in remission
C92.A2	Acute myeloid leukemia with multilineage dysplasia, in relapse
C95.90	Leukemia, unspecified not having achieved remission
C95.91	Leukemia, unspecified, in remission
C95.92	Leukemia, unspecified, in relapse

Description of Services

Technologies used for molecular profiling of cancers vary, and can include, but are not limited to, tests that evaluate variations in the genes, such as Chromosome Microarray and Next Generation Sequencing, as well as others that assess the gene products, such as gene expression arrays and microRNA analysis. The number of genes evaluated can range from a single gene to the whole exome or genome of a tumor. For the purposes of this policy, multi-gene analysis generally refers to a gene

panel containing five or more genes, though some exceptions may apply as noted specifically in the policy (e.g., epi-Colon, Clonoseq, DermTech PLA). In some tests, expression patterns of defined genes are combined in a defined manner to provide an expression signature, a score, or a classifier for potential diagnosis and or prognosis of disease or to predict impact of intervention. Results of molecular profiling may assist individuals and healthcare providers with determining prognosis and selection of more effective and targeted cancer therapies (Chantrill et al., 2015).

Clinical Evidence

Melanoma

Cutaneous Melanoma

Zager et al. (2018) conducted a multi-center trial of archived primary melanoma tumors from 523 patients, using a 31 gene expression classifier to classify patients as Class 1 (low risk) and Class 2 (high risk). The five-year recurrence free survival (RFS) rates for Class 1 and Class 2 were 88% and 52%, respectively. Distant metastasis-free survival rates (DMFS) were 93% for Class 1 versus 60% for Class 2. The gene expression classifier was a significant predictor of RFS and DMFS in univariate analysis in addition to with Breslow thickness, ulceration, mitotic rate, and sentinel lymph node (SLN) status. GEP, tumor thickness and SLN status were significant predictors of RFS and DMFS in a multivariate model that also included ulceration and mitotic rate. The authors concluded that the 31 gene expression classifier provided value to prognostication, and more prospective studies are needed.

Ardakani et al. (2017) assessed the ability of CGH to differentiate between melanocytic naevi and melanoma in cases where the two show overlapping histological features. Melanomas are characterized by CNVs, while naevi are normal. The team used 19 formalin fixed, paraffin embedded (FFPE) unambiguous naevi and 19 melanomas and tested them using a SurePrint G3 Human CGH 8x60K array. CGH was able to differentiate between the naevi and the melanoma in 95% of cases. One naevus showed two large CNV. The authors concluded that CGH may be a good adjunctive test to resolve histologically equivocal melanocytic samples.

The clinical utility of a two gene-gene expression assay by DermTech was studied by Ferris et al. (2017). A noninvasive adhesive skin patch test looking at the gene expression of *LINC/PRAME*, known as the pigmented lesion assay (PLA), was compared to the findings of 45 dermatologists who evaluated clinical and dermoscopic images of the same lesions and based on their observations, recommended biopsy or not. All samples were biopsied, and readers were blinded to the histopathology. Sixty samples were included that were obtain from March 2014 to November 2015 and represented which eight were melanomas and 52 were nonmelanomas. The biopsy concordance using only the dermatologist review was 95%. When the PLA results were included, the biopsy concordance improved to 98.6%. This clinical utility was further explored in a real-world analysis in an observational cohort of 381 patients (Ferris, et al., 2018). The PLA test was positive in 51 patients, and all had a biopsy that resulted in 37% diagnosed with melanoma. In the 330 negative PLA group, nearly all were managed by monitoring. Three had biopsies, and none were found to be melanoma. The authors calculate that 93% of samples diagnosed histologically as melanoma were positive for both *LINC/PRAME*. *PRAME* only and *LINC* only positives were histologically melanomas in 50% and 7% of cases respectively.

Berger et al. (2016) conducted a retrospective analysis to ascertain clinical management changes to 156 patients with cutaneous melanoma, based on the outcome of DecisionDx-Melanoma. Molecular risk classification by gene expression profiling has clinical impact and influences physicians to direct clinical management of CM patients. The vast majority of the changes implemented after the receipt of test results were reflective of the low or high recurrence risk associated with the patient's molecular classification. Because follow-up data was not collected for this patient cohort, the study is limited for the assessment of the impact of gene expression profile-based management changes on healthcare resource utilization and patient outcome.

Wiesner et al. (2016) provided a review on the diagnostic, prognostic, and therapeutic value of understanding genomic alterations in spitzoid tumors. Spitzoid tumors are composed of large spindle shaped or epithelioid melanocytes and are biologically distinct from melanocytic naevi and melanoma. Naevi and melanoma may have BRAF, NRAS mutations or inactivation of NF-1, Spitz tumors often have genomic rearrangements or HRAS mutation, or inactivation of BAP1. The number of genomic alterations correlates with the degree of abnormal histology and CGH analysis or FISH can accurately classify benign and malignant Spitz tumors. However, the vast majority of melanocytic tumors are histologically distinguishable as benign or malignant, so CGH provides no diagnostic value in these situations. Limited data exists on using CGH to differentiate

benign from malignant in ambiguous melanocytes, but the authors report that prior publications and their own experience shows that ambiguous tumors have more genetic aberrations than benign lesions, but fewer than malignant, so the value is limited to up grading or down grading the risk of malignancy, but doesn't necessarily give clear answers.

Kutzner et al. (2012) evaluated the use of CGH and FISH in evaluating 27 histologically ambiguous "distinct morphological variant of superficial spreading melanoma, termed 'melanomas composed exclusively or predominantly of large nests' (MLN)". Of the 27 original samples, the authors concluded that 11 met the definition of a MLN. The others were considered to be conventional spreading melanomas. They were found equally in men and women, and the average age was 61 years. The majority of MLN mirrored the typical features of melanoma, and some clinicians in the group noted that in patients with multiple melanocytic lesions, the MLNs were very different from other pigmented lesions and raised the clinical suspicion of melanoma. Eight of the 11 MLN were evaluated by CGH and 10 were also evaluated by FISH. All cases analyzed by CGH had multiple chromosome aberrations, and no one aberration was associated with the morphology of large nests and was similar to the group of conventional spreading melanomas. FISH was only positive in four cases, which were also abnormal on CGH. Cases that were abnormal by CGH, but normal on FISH, were abnormal in chromosomal regions not covered by FISH. The authors concluded that while the histological appearance created difficulties in a definitive diagnosis, "most of the MLN were correctly diagnosed as malignant melanomas by clinicians on the basis of clinical criteria." In cases that continue to confound after conventional histological examination, CGH can be useful to confirm a diagnosis.

Raskin et al. (2011) used CGH and FISH to evaluate atypical Spitz tumors in order to differentiate between melanoma and Spitz nevi. Sixteen patients with histologically ambivalent melanocytes were evaluated in the study, and of these, eight has positive sentinel lymph node biopsy, one of which also had distant metastasis. Also evaluated were eight patients with Spitz nevi, and three patients with melanoma (two spitzoid, one superficial spreading). Chromosomal aberrations were found in seven patients with ambivalent melanocytes, and there was no difference between the positive and negative lymph node biopsy groups. One had a fatal outcome. Chromosome abnormalities were also found in two spitzoid melanomas, and one conventional melanoma. The majority of aberrations found in the ambivalent group were not the ones commonly found in melanomas, suggesting that this may be a unique clinical entity. FISH failed to detect one spitzoid melanoma, one fatal metastatic case, and the other chromosomally aberrant ambivalent cases. It was positive in one spitzoid melanoma and one conventional. Overall the authors concluded that CGH may offer better diagnostic aid with better sensitivity and specificity than FISH in atypical Spitz tumors.

NCCN (2019d) clinical practice guidelines for melanoma note that molecular profiling using a variety of tests ranging from cytogenetics to chromosome microarray to gene expression is more available to help stage indeterminate melanocytic neoplasms, at this time they offer only complementary information and their clinical utility is still under investigation. Gene expression tests are available that are being marketed to help determine prognosis, but it has not yet been established that these tests provide clinically actionable information beyond known nomograms using patient demographics and histopathology. The authors do note, however, that the analysis of some somatic genetic alterations such as *BRAF* and *KIT* may be useful to guide treatment decisions, but using whole genome or exome sequencing, or next generation sequencing panels to determine mutation burden, remain investigational.

Uveal Melanoma

Klufas et al. (2017) retrospectively reviewed the role of gene expression profile analysis (GEP) vs. chromosome 3 specific analysis. Records of consecutive patients diagnosed with posterior uveal melanoma who underwent intraoperative fine needle aspiration biopsy prior to placement of an iodine-125 radioactive plaque between 2012 and 2014 were reviewed. Two cohorts of patients were identified. Cohort 1 had 44 patients, and tumors had both GEP and FISH analysis. Cohort 2 had 43 patients, and those tumors had GEP and multiplex ligation-dependent probe amplification (MLPA) results were obtained. Discordance between GEP and chromosome 3 status by FISH and MLPA occurred in the series at a rate of 15.9 and 16.3%, respectively. The authors concluded that caution must be advised when counseling a patient with a good-prognosis GEP "Class 1" result that the uveal tumor may actually harbor monosomy 3, which is associated with a poor prognosis for metastasis in nearly 20% of the patients.

Plasseraud et al. (2016) evaluated the clinical validity and utility of DecisionDx-UM in a prospective, multicenter, study (supported by Castle Biosciences, Inc.). 70 patients were enrolled to document patient management differences and clinical outcomes associated with low-risk Class 1 and high-risk Class 2 results indicated by DecisionDx-UM testing. Thirty-seven patients in the prospective study were Class 1 and 33 were Class 2. Class 1 patients had 100% 3-year metastasis-free survival compared to 63% for Class 2 (log rank test $p = 0.003$) with 27.3 median follow-up months in this interim analysis. Class 2 patients received significantly higher-intensity monitoring and more oncology/clinical trial referrals compared to Class 1

patients (Fisher's exact test $p = 2.1 \times 10^{-13}$ and $p = 0.04$, resp.). In the authors' opinion, the results of this study provide additional, prospective evidence in an independent cohort of patients for which Class 1 and Class 2 patients are managed according to the differential metastatic risk indicated by DecisionDx-UM. A study limitation is financial sponsorship/support by the manufacturer which increases the risk of bias.

Minca et al. (2014) noted that monosomy 3 and MYC amplification at 8q24 are strong prognosticators of outcomes in uveal melanoma (UVM) and is commonly detected by FISH. They hypothesized that CMA would be an alternative to FISH and have advantages in identifying loss of heterozygosity, partial chromosome loss and other aberrations that FISH can't detect. They analyzed CMA using SNP+CGH (Roche-NimbleGen OncoChip) on enucleations from formalin-fixed paraffin-embedded tissue (FFPET) for 34 patients and/or frozen tissue (FZT) for 41 patients. CMA was successful in 30 of 30 of 34 FFPET and all 41 FZT. In 27 paired FFPET/FZT samples 96% were concordant for at least four of six major chromosome abnormalities, and 93% were concordant for one (- chromosome 3). CMA was concordant with FISH in 90% of FFPET and 93% of FZT. Partial -3q was detected in two FISH negative cases and whole chromosome LOH for 3, 4 and 6 in one case. Results of UVM SNP+CGH genotyping were significantly correlated with clinical outcome and reliably predicted metastasis, time to progression, and survival. The authors concluded that SNP+CGH is a practical method for UVM prognostication, and provides additional data with relevance to biology, diagnosis and prognosis.

In a prospective multi-center validation study, Onken et al. (2012) evaluated the prognostic performance of a 15 gene expression profiling (GEP) assay that assigned primary posterior uveal melanomas to prognostic subgroups: class 1 (low metastatic risk) and class 2 (high metastatic risk). A total of 459 patients were enrolled. Analysis was performed to compare the prognostic accuracy of GEP with Tumor-Node-Metastasis (TNM) classification and chromosome 3 status. Patients were managed for their primary tumor and monitored for metastasis. The GEP assay successfully classified 446 of 459 cases (97.2%). The authors concluded that the GEP assay had a high technical success rate and was the most accurate prognostic marker among all of the factors analyzed. The GEP provided a highly significant improvement in prognostic accuracy over clinical TNM classification and chromosome 3 status. Further studies are needed to determine the clinical utility of these tests and the role they have in clinical decision-making.

NCCN introduced guidelines in 2018 for the staging and management of uveal melanoma. While they note that tumor specimens may be sent for chromosome analysis and/or gene expression profiling, they note that biopsy is usually not necessary for initial diagnosis or to make a treatment selection and does not impact patient outcomes. (NCCN, 2018b)

Clinical Practice Guidelines

American Academy of Dermatology (AAD)

The AAD does not address molecular testing in their guidelines at any point, and states in general that baseline laboratory tests are generally not recommended in asymptomatic patients with newly diagnosed primary melanoma of any thickness, and that such tests have low yield for detection of metastatic disease and are associated with relatively high false-positive rates (Bichakjian, 2011).

European Society for Medical Oncology (ESMO)

In their 2015 guidelines, ESMO states that genetic testing is generally not recommended for melanoma diagnoses, but notes that biomarkers such as mutations (NRAS, c-Kit, BRAF) are already indispensable today for a personalized medicine approach in advanced melanoma. Broader panels are not recommended, though it is noted that additional mutations and the overall mutation rate might provide additional molecular predictive markers in the near future (Drumer, 2015).

Cancers of Unknown Primary Site

Varadhachary and Raber (2014) reviewed the research, diagnosis and treatment of CUP, noting that the performance of tissue-of-origin molecular-profiling assays in known cancers has been validated with the use of independent, blinded evaluation of sets of tumor samples, with an accuracy of approximately 90%. Based on these findings, the authors comment that the feasibility of using formalin-fixed samples obtained from small, core-needle biopsy or using samples obtained by means of fine-needle aspiration makes this method practical for use in the clinic setting. However, without randomized, controlled trials it is difficult to gauge the therapeutic effect of tissue-of-origin molecular-profiling assays. Further, they suggest that creative trial designs are urgently needed in order to study subsets of unknown primary cancers and the effect of these assays on survival and quality of life of patients.

Meleth et al. (2013) conducted a technology assessment on genetic testing or molecular pathology testing for cancer of unknown primary cancers with CancerTypeID, miRview, or PathworkDx to determine analytical validity, clinical validity, and clinical utility. The results showed that the clinical accuracy of all the three tests is similar, ranging from 85 percent to 88 percent. The evidence that the tests contribute to identifying a TOO is moderate; however, the researchers noted that they did not have sufficient evidence to assess the effect of the tests on treatment decision and outcomes.

In a systematic review of loss-of-heterozygosity based topographic genotyping with PathfinderTG®, Trikalinos et al. (2010) found no studies that demonstrated longer survival, longer time to tumor recurrence, or fewer adverse outcomes as a result attributable to unnecessary harmful interventions, as a result of this testing. The authors reported several limitations with eligible studies including limited sample size and lack of patient selection criteria.

In a guideline on the diagnosis and management of metastatic malignant disease of unknown primary origin in adults, the National Institute of Health and Care Excellence (NICE) (2010) does not recommend the use of gene-expression-based profiling to identify primary tumors in patients with provisional CUP. They also do not recommend the use of gene expression-based profiling when deciding which treatment to offer patients with confirmed CUP.

National Comprehensive Cancer Network (NCCN) clinical practice guidelines for occult primary (cancer of unknown primary site) state that while there is diagnostic benefit of gene expression profiling (GEP) assays, it is not different than immunohistochemical staining and a clinical benefit for GEP or next generation sequencing has not been demonstrated. Consequently, the panel does not recommend tumor sequencing and gene signature profiling for the identification of tissue of origin as standard management in the diagnostic workup of patients with occult primary tumors. In addition, the NCCN suggests that pathologists and oncologists collaborate on the judicious use of these modalities on a case-by-case basis, with the best individualized patient outcome in mind (NCCN, 2019f).

Clinical Practice Guidelines

European Society for Medical Oncology (ESMO)

In a clinical practice guideline for the diagnosis, treatment and follow-up on cancers of unknown primary (CUP) site, ESMO (Fizazi et al., 2015) did not identify any significant differences in the tumor microRNA expression profile when CUP metastases biologically assigned to a primary tissue of origin were compared with metastases from typical solid tumors of known origin. Although they noted that these tests may aid in the diagnosis of the putative primary tumor site in some patients, their impact on patient outcome via administration of primary site-specific therapy remains questionable and unproven in randomized trials.

Colorectal Cancer (CRC)

Zhang et al. (2016) retrospectively reviewed the prognostic role of CDX2 expression in patients with stage I and stage III metastatic colorectal cancer (CRC) after complete surgical resection. The patient cohort (n=145) included 66 patients with CDX2-negative metastatic CRC and a comparison cohort of 79 patients with CDX2-positive metastatic CRC. The prevalence of absent CDX2 expression in this cohort was 5.6%. After adjusting for covariates in a multivariate model, the association of a lack of CDX2 expression and OS remained statistically significant (HR, 4.52; 95% CI, 2.50-8.17; P < .0001). In addition, the median PFS (3 vs. 10 months; HR, 2.23; 95% CI, 1.52-3.27; P < .0001) for first-line chemotherapy was significantly decreased in patients with CDX2-negative metastatic CRC. The authors concluded that the results showed that a lack of CDX2 expression in metastatic CRC is an adverse prognostic feature and a potential negative predictor of the response to chemotherapy. Further research with randomized controlled trials is needed to validate these findings.

To evaluate whether patients with CDX2-negative tumors might benefit from adjuvant chemotherapy, Dalerba et al. (2016) investigated the association between CDX2 status, and assessed at either the mRNA or protein level, the disease-free survival among patients who either did or did not receive adjuvant chemotherapy. Reviewing a database of 669 patients with stage II colon cancer and 1,228 patients with stage III colon cancer, the authors reported that their results confirmed that treatment with CDX2 as a biomarker in colon cancer adjuvant chemotherapy was associated with a higher rate of disease-free survival in both the stage II subgroup (91% with chemotherapy vs. 56% with no chemotherapy, P = 0.006) and the stage III subgroup (74% with chemotherapy vs. 37% with no chemotherapy, P < 0.001) of the CDX2-negative patient population (Fig. 5). A test for the interaction between the biomarker and the treatment revealed that the benefit observed in CDX2-negative cohorts was superior to that observed in CDX2-positive cohorts in both the stage II subgroup (P = 0.02 for the interaction) and the stage III subgroup (P = 0.005 for the interaction). In the authors' opinion, their results indicate that patients with stage II or stage III CDX2-negative colon cancer might benefit from adjuvant chemotherapy and that adjuvant chemotherapy might be a treatment option for

patients with stage II CDX2-negative disease, who are commonly treated with surgery alone. Given the exploratory and retrospective design of this study, these results will need to be further validated through randomized, clinical trials, in conjunction with genomic DNA sequencing studies.

Yamanaka et al. (2016) evaluated the 12-gene Recurrence Score assay for stage II and III colon cancer without chemotherapy to reveal the natural course of recurrence risk in stage III disease (the Sunrise Study). A cohort-sampling design was used. From 1,487 consecutive patients with stage II to III disease who had surgery alone, 630 patients were sampled for inclusion with a 1:2 ratio of recurrence and nonrecurrence. Sampling was stratified by stage (II v III). The assay was performed on formalin-fixed, paraffin-embedded primary cancer tissue. Association of the Recurrence Score result with recurrence-free interval (RFI) was assessed by using weighted Cox proportional hazards regression. With respect to prespecified subgroups, as defined by low (< 30), intermediate (30 to 40), and high (≥ 41) Recurrence Score risk groups, patients with stage II disease in the high-risk group had a 5-year risk of recurrence similar to patients with stage IIIA to IIIB disease in the low-risk group (19% v 20%), whereas patients with stage IIIA to IIIB disease in the high-risk group had a recurrence risk similar to that of patients with stage IIIC disease in the low-risk group (approximately 38%). The authors conclude that this validation study of the 12-gene Recurrence Score assay in stage III colon cancer without chemotherapy showed the heterogeneity of recurrence risks in stage III as well as in stage II colon cancer.

Venook et al. (2013) conducted a validation study of the 12-gene recurrence score in cancer and leukemia group B (CALGB) 9,581 of 1,713 randomly assigned patients with stage II colon cancer to treatment with edrecolomab or observation and found no survival difference. The analysis reported included all patients with available tissue and recurrence (n=162) and a random (approximately 1:3) selection of nonrecurring patients. RS was assessed in 690 formalin-fixed paraffin-embedded tumor samples with quantitative reverse transcriptase polymerase chain reaction by using prespecified genes and a previously validated algorithm. Association of RS and recurrence was analyzed by weighted Cox proportional hazards regression. The researchers concluded that 12-gene RS predicts recurrence in stage II colon cancer in CALGB 9581, which is consistent with the importance of stromal response and cell cycle gene expression in colon tumor recurrence. RS appears to be most discerning for patients with T3 MMR-I tumors, although markers such as grade and lymphovascular invasion did not add value in this subset of patients.

In a validation study of the 12-gene colon cancer recurrence score in NSABP C-07 as a predictor of recurrence in patients with stage II and III colon cancer treated with fluorouracil and leucovorin (FU/LV) and FU/LV plus oxaliplatin, Yothers et al. (2013). Recurrence Score was assessed in 892 fixed, paraffin-embedded tumor specimens (randomly selected 50% of patients with tissue). Data were analyzed by Cox regression adjusting for stage and treatment. Based on the results, the authors concluded that 12-gene Recurrence Score predicts recurrence risk in stage II and stage III colon cancer and provides additional information beyond conventional clinical and pathologic factors. Incorporating Recurrence Score into the clinical context may better inform adjuvant therapy decisions in stage III as well as stage II colon cancer.

NCCN clinical practice guidelines for colon cancer review several multigene panels for prognosis and recurrence, including Oncotype Dx Colon, ColoPrint, and ColDx. The panels state that there is insufficient data to recommend the use of multigene assay panels to determine adjuvant therapy in colon cancer patients (NCCN, 2019c, 2019h).

Molecular technologies are also under investigation to screen for colon cancer, such as the Epi proColon 2.0 assay that measures the methylated Septin9 (SEPT9), a circulating tumor cell marker. The premise of this test is that during colorectal cancer development, the tumor will release cell free DNA (cfDNA) into the bloodstream, and the ratio of SEPT9 DNA be detected through specialized techniques and can predict the presence of early colorectal cancer. A meta-analysis of one cohort study and thirteen case-controlled studies representing 9870 cases demonstrated a pooled sensitivity of 0.66 and specificity of 0.91. The authors compared this to data available for the gold standard test, fecal occult blood testing (FOBT) of a sensitivity of 0.60 and specificity of 0.91, equal to SEPT9. The authors combined the results of FOBT and SEPT9 and achieved a detection rate of colorectal cancer of 88.7% with a specificity of 78.8%. They concluded that FOBT and SEPT9 complement each other, but further studies are needed to determine the best screening tests and approaches (Yan et al., 2016).

He et al. (2018) examined the clinicopathological features that could impact the sensitivity and specificity of SEPT9 analysis. A total of 1,160 patients were included in the study from hospitals in China, which included 300 patients with colorectal cancer, 122 patients with adenoma, 103 patients with hyperplastic polyps, 568 normal participants (no evidence of disease), and 67 patients with other gastrointestinal diseases. Overall, the sensitivity and specificity of SEPT9 was impacted by cancer stage, size, invasion depth, classification, differentiation and metastasis. It was also noted that SEPT9 detected adenomas,

hyperplastic polyps and other gastrointestinal diseases such as inflammatory bowel disease. When screening an average risk population, these non-colorectal cancer disorders are much more common and could lead to false positives and unnecessary intervention.

ColonSentry is a blood-based gene expression test that assesses the expression of ANXA3, CLEC4D, LMNB1, PRRG4, TNFAIP6, VNN1, and IL2RB genes using real time PCR, and reports results as a cumulative relative risk score (CURR). In a 2014 evaluation of available data, Heichman (2014) reviewed the work of Han et al. (2008) and Marshall et al. (2010) that explored the clinical utility of the test and reported that in a case controlled study of 202 colorectal cancer patients and 208 matched healthy controls, a specificity of 70% for distinguishing cancer from healthy controls, and a sensitivity of 72% for identifying colorectal cancer. Larger, prospective studies are needed to further confirm the performance of this test.

Prostate Cancer

A study from McKiernan et al (2016) evaluated the performance of the ExoDx Prostate IntelliScore urine exosome assay. The study compared those patients who received standard of care (SOC) alone to those who received SOC plus this novel exosome assay. SOC was defined as PSA levels, age, race, and family history. ExoDx Prostate IntelliScore urine exosome assay is a noninvasive, urinary 3-gene expression assay that is designed to discriminate high-grade (> Gleason Score 7) from low-grade (Gleason Score 6) and benign disease. The researchers compared the urine exosome gene expression assay with biopsy outcomes in 499 patients with PSA levels of 2 to 20 ng/ml. After this first phase, the derived prognostic score was validated in 1064 patients that included PCA-free men, 50 years or older, scheduled for an initial or repeated prostate needle biopsy due to suspicious digital rectal examination (DRE) findings and/or PSA levels (limit range, 2.0-20.0 ng/mL). This study found that in 255 men in the training target population (median age 62 years and median PSA level 5.0 ng/mL, and initial biopsy), the urine exosome gene expression assay plus SOC was associated with enhanced discrimination between GS7 or greater and GS6 and benign disease (AUC 0.77 (95% CI, 0.71-0.83) vs SOC AUC 0.66 (95% CI, 0.58-0.72) (P < .001)). The validation study found that in 519 patients' urine exosome gene expression assay plus SOC AUC 0.73 (95% CI, 0.68-0.77) was superior to SOC AUC 0.63 (95% CI, 0.58-0.68) (P < .001). Using a predefined cut point, 138 of 519 (27%) biopsies would have been avoided, missing only 5% of patients with dominant pattern 4 high-risk GS7 disease. This study concluded that the urine exosome gene expression assay was associated with improved identification of patients with higher-grade prostate cancer among men with elevated PSA levels and could reduce the total number of unnecessary biopsies.

McKiernan et al (2018) assessed the performance and utility of ExoDx Prostate (IntelliScore) (EPI) urine exosome gene expression assay versus SOC parameters for discriminating grades of prostate cancer from benign disease was evaluated. This study compared EPI results with biopsy outcomes in men with age \geq 50 yr. and prostate-specific antigen (PSA) 2–10 ng/ml, scheduled for initial prostate biopsy. The results were that in a total of 503 patients, with median age of 64 yr., median PSA 5.4 ng/ml, 14% African American, 70% Caucasian, 53% positive biopsy rate (22% GG1, 17% GG2, and 15% \geq GG3), EPI was superior to SOC with an area under the curve (AUC) 0.70 versus 0.62, respectively, comparable with previously published results (n=519 patients, EPI AUC 0.71). Using a validated cut-point 15.6 would have avoided 26% of unnecessary prostate biopsies and 20% of total biopsies, with negative predictive value (NPV) 89% and missing 7% of \geq GG2 PCa. Setting a different cut-point 20 would avoid 40% of unnecessary biopsies and 31% of total biopsies, with NPV 89% and missing 11% of \geq GG2 PCa. This study concluded that EPI has been validated in over 1000 patients across two prospective validation trials for risk stratification of high-grade and low-grade from benign disease. The use of test may improve identification of patient with higher grade disease and could reduce unnecessary biopsies; although 10% of prostate cancer cases would be missed based on the test characteristics.

Klein et al. (2016) retrospectively analyzed prostatectomy tissue of 337 Gleason 3+3 patients. To compare clinico-pathologic variables across pathologic Gleason score categories, Fisher's exact test or analysis of variance F test were used. Distributions of Decipher scores among different clinico-pathologic groups were compared using Wilcoxon rank sum test. The association of Decipher score and adverse pathology was examined using logistic regression models. Among men who had Gleason 3+3=6 disease only, 269 (80%) had low Decipher scores with 43 (13%) and 25 (7%) harboring intermediate and high scores respectively. Thus, a small proportion of histologic Gleason 6 tumors harbor molecular characteristics of aggressive cancer. The authors note that molecular profiling of such tumors at diagnosis may better select patients for active surveillance at the time of diagnosis and trigger appropriate intervention during follow-up.

Glass et al. (2016) published long-term outcomes to a previously reported validation study on Decipher. Study subjects (n=224) had aggressive prostate cancer with at least one of several criteria such as preoperative prostate specific antigen 20 ng/ml or greater, pathological Gleason score 8 or greater, stage pT3 disease or positive surgical margins at prostatectomy. Of the 224

patients treated 12 experienced clinical recurrence, 68 had biochemical recurrence and 34 experienced salvage treatment failure. At 10 years after prostatectomy the recurrence rate was 2.6% among patients with low Decipher scores but 13.6% among those with high Decipher scores ($p=0.02$). When CAPRA-S and Decipher scores were considered together, the discrimination accuracy of the ROC curve was increased by 0.11 compared to the CAPRA-S score alone (combined c-index 0.84 at 10 years after radical prostatectomy) for clinical recurrence. The authors conclude that Decipher improves the ability to predict clinical recurrence in prostate cancer and adds precision to conventional pathological prognostic measures. Long-term studies are needed to validate these results.

Den et al. (2016) conducted a retrospective review of 2,341 consecutive radical prostatectomy patients to understand the relationship between the Decipher classifier test and patient tumor characteristics. Decipher score had a positive correlation with pathologic Gleason score (PGS; $r = 0.37$, 95% confidence interval (CI) 0.34 – 0.41), pathologic T-stage ($r = 0.31$, 95% CI 0.28 – 0.35), CAPRA-S ($r = 0.32$, 95% CI 0.28 – 0.37) and patient age ($r = 0.09$, 95% CI 0.05-0.13). Decipher reclassified 52%, 76% and 40% of patients in CAPRA-S low-, intermediate- and high-risk groups, respectively. The authors detected a 28% incidence of high-risk disease through the Decipher score in pT2 patients and 7% low risk in pT3b/pT4, PGS 8–10 patients. There was no significant difference in the Decipher score between patients from community centers and those from academic centers ($P = 0.82$). The authors concluded that although Decipher correlated with baseline tumor characteristics for over 2 000 patients, there was significant reclassification of tumor aggressiveness as compared to clinical parameters alone. In their opinion, utilization of the Decipher genomic classifier can have major implications in assessment of postoperative risk that may impact physician-patient decision making and ultimately patient management.

Marrone et al. (2015) did a literature review of the Decipher test, a 22 gene expression assay designed to predict the metastatic rate of prostate cancer within five years of a radical prostatectomy. They utilized PubMed to search for peer reviewed literature that discussed the analytic validity, clinical validity and clinical utility of Decipher. Eight studies were identified, but no guidelines. Analytical validity was identified by the authors in a single conference abstract, and the correlation between genomic classifier scores between matched biopsies was 74%. Clinical validity was described in all included studies, and the authors found that the data represented that the genomic classifier was able to adequately discriminate between those men that developed metastatic prostate cancer within five years and those that did not. Clinical utility was another matter, however. The authors found that additional evidence was needed to show that outcomes were improved in men whose post-surgical treatment was guided by Decipher results when compared to standard of care.

Oderda et al. (2016) assessed whether cell-cycle progression (CCP)-score (Polaris) can improve the current risk assessment in newly diagnosed prostate cancer (PCa) patients. The CCP-score at biopsy was evaluated in 52 patients newly diagnosed with PCa who underwent radical prostatectomy. CCP-score was calculated as average RNA expression of 31 CCP genes, normalized to 15 housekeeping genes. The predictive ability of CCP-score was assessed in univariate and multivariate analyses and compared to that of Ki-67 levels and traditional clinical variables including prostate-specific antigen, Gleason score, stage, and percentage of positive cores at biopsy. The authors reported that in spite of an overall good accuracy in attributing the correct risk class, 7 high-risk and 13 intermediate-risk patients were misclassified by the Polaris test, which is a limitation to this study. On analysis of variance, mean CCP-score significantly differed across different risk classes based on pathologic results (-1.2 in low risk, -0.444 in intermediate risk, 0.208 in high risk). CCP-score was a significant predictor of high-risk PCa both on univariate and multivariate analyses, after adjusting for clinical variables. Combining CCP-score and the European Association of Urology clinical risk assessment improved the accuracy of risk attribution by around 10%, up to 87.8%. CCP-score was a significant predictor of biochemical recurrence, but only on univariate analysis. The authors conclude that the CCP-score might provide important new information to risk assessment of newly diagnosed PCa in addition to traditional clinical variables. A correct risk attribution is essential to tailor the best treatment for each patient. Additional studies with larger patient sample sizes are needed to determine whether the use of this test in making treatment decisions improves patient outcomes.

Shore et al. (2014) evaluated the clinical utility of the CCP score in a U.S.-based clinical setting. Urologists who participated in a prospective clinical study were sent a retrospective questionnaire to assess the value of the CCP test results. Fifteen urologists participated in the study, representing 15 distinct urology group practices. Questionnaires were received for 294 evaluable patients. All patients had localized prostate cancer. Physicians found the CCP score valuable and indicated that 55% of tests generated a mortality risk that was either higher or lower than expected. Physicians also indicated that 32% of test results would lead to a definite or possible change in treatment. The data suggest that the test would have the net effect of shifting patients from more aggressive treatment to more conservative treatment. This was evidenced by the significant association between change in treatment and lower CCP scores. Results of this survey study provide only indirect evidence of clinical utility as the study measured the likelihood of change in treatment as estimated by the physician, not the actual change in treatment. The

authors concluded that real-world use of the test is likely to lead to a change in treatment in a significant portion of tested patients, particularly by shifting patients towards more conservative management.

Crawford et al. (2014) conducted a prospective survey study evaluating the impact of the CCP score on physician treatment recommendations for prostate cancer. Physicians ordering the test completed surveys regarding treatment recommendations before and after they received and discussed test results with patients. Clinicians also rated the influence of the test result on treatment decisions. For patients originally targeted for interventional therapy, results of the CCP test led to a 37.2% reduction of interventional therapy. For patients originally targeted for noninterventional therapy, 23.4% of patients had treatment changes to interventional therapy based on test results. Overall, surgical interventions were reduced by 49.5%, and radiation treatment was reduced by 29.6%. Author-reported limitations included physician selection of patients for testing, no evaluation of patient input in therapeutic choice and other potential treatment decision factors not queried by the survey. Results of this survey study provide only indirect evidence of clinical utility since it does not capture clinical outcomes.

Brand et al. (2016) performed a meta-analysis of two independent clinical validation studies of a 17-gene biopsy-based genomic assay (Oncotype Dx Prostate Cancer Assay) as a predictor of favorable pathology at radical prostatectomy. Patient-specific meta-analysis was performed on data from two studies (732 patients) using the Genomic Prostate Score (GPS; scale 0-100) together with Cancer of the Prostate Risk Assessment (CAPRA) score or National Comprehensive Cancer Network (NCCN) risk group as predictors of the likelihood of favorable pathology (LFP). Risk profile curves associating GPS with LFP by CAPRA score and NCCN risk group were generated. Patient-specific meta-analysis generated risk profiles ensure more precise estimates of LFP with narrower confidence intervals either study alone. GPS added significant predictive value to each clinical classifier. The authors concluded that a model utilizing GPS and CAPRA provided the most risk discrimination, and in a decision curve analysis, greater net benefit was shown when combining GPS with each clinical classifier compared with the classifier alone. Although the clinical characteristics of the two patient cohorts were similar, there were nonetheless some key differences in the representation of different racial groups and higher risk patients. The risk estimates were numerically different in the two studies, although the confidence levels overlapped.

In a review of tissue-based genomic biomarkers for prostate cancer, Moschini et al. (2016), report that available genomic assays have improved the prognostic ability over clinicopathologic parameters of localized prostate cancer (PCa). However, these assays should be prospectively applied, or even retrospectively applied to prospective studies, to validate their clinical utility in prognostication and even prediction in terms of what treatment should be applied either at a new diagnosis or post-RP.

Na et al. (2016) reviewed the literature on clinically available RNA profiling tests (Oncotype Dx, Prolaris, and Decipher) of prostate tumors. They concluded that these RNA profiling panels have shown promising results in regard to clinical utility, several limitations are worth noting: (1) the current studies are retrospective with relatively small sample sizes, so larger-scale prospective randomized trials are necessary for validation; (2) RNA quality varies among panels (e.g., microdissection is needed for Decipher [some medical center may not have the equipment], while for Prolaris, tissue extraction relies on the instruction from pathologist, which will lead to heterogeneity of the testing results); and (3) the relatively high prices limit potential use of the panels, will necessitate further evaluation of their cost-effective values.

NCCN clinical practice guidelines for prostate cancer state that molecular profiling of biopsies may be considered in men with low and favorable intermediate risk prostate cancer and a life expectancy greater than or equal to ten years to help guide decision-making on treatment. However, NCCN cautions that these tests (Decipher, Oncotype Dx Prostate, Prolaris or ProMark) have been developed with extensive industry support, guidance and involvement and have been marketed under the less rigorous FDA regulatory pathway for biomarkers. In addition, full assessment of their clinical utility requires prospective, randomized clinical trials. (NCCN, 2019g).

Clinical Practice Guidelines

American Urological Association (AUA) with the American Society for Radiation Oncology (ASTRO) and the Society for Urologic Oncology (SUO)

Sanda et al. (2018) published the joint AUA/ASTRO/SUO guidelines for clinical localized prostate cancer. The guidelines stated that tissue based genomic biomarkers have not shown a clear role in active surveillance for localized prostate cancer and are not necessary for follow up.

American Urological Association (AUA)

In a clinical practice guideline on early detection of prostate cancer (Carter et al., 2013; reviewed and confirmed 2018) based on a systematic review and meta-analysis, the AUA notes that an improved understanding of the interaction between inherited risk alleles and the environment (lifestyle choices) could provide a potential means of prevention. Future studies of the genetic and epigenetic basis of disease development and progression may provide biomarkers and/or panels of biomarkers with improved specificity when compared to PSA. When available, risk assessment tools combining multiple predictors will need to be evaluated in carefully designed trials to be generalizable to the population in which they would be used.

American Society of Clinical Oncology (ASCO)

In 2018, Bekelman et al. (2018) published the ASCO endorsement of their guidelines, developed in 2017, for managing clinically localized prostate cancer (Sanda et al., 2018). This guideline stated that tissue based genomic biomarkers have not shown a clear role in active surveillance and not necessary for follow up.

In an endorsement of Cancer Care Ontario's guideline on active surveillance of localized prostate cancer, ASCO comments that ancillary radiologic and genomic tests are investigational but may have a role in patients with discordant clinical and/or pathologic findings. Prospective validation of these tests is needed to assess their impact on patient outcomes such as survival (Chen et al., 2016).

Lung Cancer

NCCN guidelines for NSCLC Panel have added a section on Plasma Cell-Free/Circulating Tumor DNA Testing that states that cell-free/circulating tumor DNA testing should not be used in lieu of a tissue diagnosis as the analytical standards have not been established. However, NCCN also suggests that the use of cell-free/circulating tumor DNA testing can be considered in specific clinical circumstances, most notably: if a patient is medically unfit for invasive tissue sampling; or in the initial diagnostic setting, if following pathologic confirmation of a NSCLC diagnosis there is insufficient material for molecular analysis, cell-free/circulating tumor DNA should be used only if follow up tissue based analysis is planned for all patients in which an oncogenic driver is not identified (NCCN, 2019e).

Drilon et al. (2015) identified 31 patients with lung adenocarcinoma with a ≤ 15 pack-year smoking history whose tumors previously tested "negative" for alterations in 11 genes (mutations in EGFR, ERBB2, KRAS, NRAS, BRAF, MAP2K1, PIK3CA, and AKT1 and fusions involving ALK, ROS1, and RET) via multiple non-NGS methods. A broad, hybrid capture-based NGS assay (FoundationOne) was performed (4,557 exons of 287 cancer-related genes and 47 introns of 19 genes frequently rearranged in solid tumors). A genomic alteration with a corresponding targeted therapeutic based on the National Comprehensive Cancer Network (NCCN) guidelines for non-small cell lung cancer (NSCLC) was found in 26% (n=8 of 31) of patients. The drivers identified in tumors from these 8 patients were EGFR G719A, BRAF V600E, SOCS5-ALK, HIP1-ALK, CD74-ROS1, KIF5B-RET (n=2), and CCDC6-RET. Six of these patients went on to receive targeted therapy. The authors noted that the reasons for non-detection of these genomic alterations via non-NGS testing can be varied such as lower sensitivity, complex rearrangements undetectable by standard FISH, and, possibly, heterogeneity between different tumor biopsies or sites. They concluded that broad, hybrid capture-based NGS assays have the potential to uncover clinically actionable genomic alterations in never smokers or ≤ 15 pack-year smokers whose lung adenocarcinomas do not harbor a potential driver via non-NGS testing.

Clinical Practice Guidelines

American College of Chest Physicians (ACCP)

In an evidence-based clinical practice guideline for the diagnosis and management of lung cancer, the ACCP states that the epidemiology of lung cancer is an active field. According to the ACCP, researchers in the area of molecular epidemiology are making advances in the identification of biomarkers of risk and for early detection, although these are not yet mature enough for clinical application (Detterbeck et al., 2013).

American Society of Clinical Oncology (ASCO)

ASCO endorsed the College of American Pathologists/International Association for the Study of Lung Cancer/Association for Molecular Pathology Clinical Practice Guideline Update with minor modifications (Kalemkerian et al., 2018). The guidelines, supported by ASCO, include the following relevant points, considered to be 'expert consensus opinion'.

- Physicians may use molecular biomarker testing in tumors with:
 - An adenocarcinoma component;

- Nonsquamous, non–small-cell histology;
- Any non–small-cell histology when clinical features indicate a higher probability of an oncogenic driver (e.g., young age [< 50 years]; light or absent tobacco exposure).
- BRAF testing should be performed on all patients with advanced lung adenocarcinoma, irrespective of clinical characteristics. RET, or KRAS, or MET molecular testing are not recommended as single gene routine stand-alone assays outside the context of a clinical trial. It is appropriate to include these as part of larger testing panels performed either initially or when routine EGFR, ALK, BRAF, and ROS1 testing is negative.
- Multiplexed genetic sequencing panels are preferred where available over multiple single-gene tests to identify other treatment options beyond EGFR, ALK, BRAF, and ROS1.
- Circulating tumor cell free DNA testing, also called a liquid biopsy, should not be routinely considered due to lack of evidence of efficacy. However, the expert consensus opinion provided is that cfDNA may be used in some clinical settings in which tissue is limited and/or insufficient for molecular testing to identify EGFR mutations.

National Comprehensive Cancer Network (NCCN)

NCCN guidelines for NSCLC (NCCN, 2017) strongly endorse the use of broad molecular profiling (also known as precision medicine) to detect certain rare mutations using multiplex or NGS. Presence of EGFR-activating mutations represents a critical biological determinant for proper therapy selection in patients with lung cancer, stating “determination of the specific molecular abnormalities of the tumor is critical for predicting sensitivity or resistance to an increasing number of drugable targets, primarily tyrosine kinase inhibitors (TKIs)”. Data has shown that targeted therapy is potentially very effective in patients with specific gene mutations or rearrangements. The guidelines specifically report that “EGFR and ALK testing be conducted as part of broad molecular profiling.” The NCCN Panel states that such testing would ensure that patients receive the most effective available targeted treatment for NSCLC.

Thyroid Cancer

There have been multiple studies, prospective and retrospective, for the commercially available molecular classifiers for indeterminate and suspected malignant thyroid nodules, such as the Afirma Gene Expression Classifier, and next generation sequencing test panels, such as ThyGenX/ThyraMir and ThyroSeq v3.

The Afirma gene classifier, a gene expression analysis of 167 genes, has a sensitivity of 92% with a negative predictive value (NPV) of 93% in the largest prospective study of indeterminate nodules to date (Alexander et al., 2012). However, a study performed in a community hospital–based thyroid surgery practice (Harell and Bimston, 2014) showed a lower NPV (89.6%) than other studies in the literature, leading some to conclude (Zhang and Lin, 2016, Marti et al., 2015) that the Afirma test will only provide the most useful information in a practice setting with a prevalence of malignancy in indeterminate thyroid lesions of 15% to 21% where a NPV $>95\%$ and PPV $>25\%$ would be expected. Outside this range it is unlikely the test can provide information that would alter management. Marti et al. (2015) conducted a retrospective review of the Afirma gene classifier at two institutions from February 2013 to December 2014 and found that there were wide variations in the Afirma GEC-benign call rate, PPV, and NPV between the two institutions; one a comprehensive health system with a TMC prevalence of 30–38% and the second a tertiary referral cancer center with a prevalence 10–19%. Each had differing rates of malignancy in indeterminate thyroid nodules and Afirma did not routinely alter management in both institutions, and the NPV ranged from 86-98%. In addition, the Afirma 167 gene classifier appears to be less accurate in nodules with that contain benign Hurthle cells. In several studies that examined the cytology population percentage of Hurthle cells, the test was more likely to report a suspicious for malignancy result for which the patient was sent for surgery, and therefore limited the clinical utility of the test (Harrell and Bimston, 2014, Brauner et al., 2015, Lastra et al., 2014).

Deaver et al. (2018) conducted a retrospective analysis of 2019 thyroid FNA from 2011 to 2015. The samples were categorized using the Bethesda System for reporting thyroid cytology into B3 and B4 nodules. GEC results from Afirma were available for 54% of B3 cases, with about half having a benign classification. In the B4 group, 52% had GEC, with 28.6% classified as benign. The authors followed 73 benign GEC cases. Five underwent surgery and no malignancy was found. The remainder continued to have a stable size, and in those that had repeat FNA, about 72%, no malignancy was noted. The authors concluded that GEC results accurately predicted benign thyroid nodules.

In a meta-analysis of the gene expression classifier(GEC) for the diagnosis of indeterminate thyroid nodules, Santhanam et al. (2016) evaluated 7 out of 58 potential studies. The reference standard for determination of benign or malignant nodules was the histopathology of the thyroidectomy specimen. A QUADAS-2 report for all studies included in the final analysis was tabulated

for risk of bias and applicability. The pooled sensitivity of the GEC was 95.7% (95% CI 92.2-97.9, I (2) value 45.4%, $p = 0.09$), and the pooled specificity was 30.5% (95% CI 26.0-35.3, I (2) value 92.1%, $p < 0.01$). Overall, the diagnostic odds ratio was 7.9 (95% CI 4.1-15.1). Although the meta-analysis revealed a high pooled sensitivity and low specificity for the Afirma GEC, patients with a benign GEC were not followed long enough to ascertain the actual false-negative rates of the index test.

Partyka et al. (2018) recently conducted a small retrospective study on 10 archived FNA samples comparing two commercially available miRNA tests, ThyraMir and RosettaGxReveal. The samples represented follicular lesion of undetermined significance (FLUS, $n=5$), follicular neoplasm/suspicious for follicular neoplasm (FN/SFN, $n=4$), and suspicious for malignancy (SM, $n=1$). Of the seven cases with benign histology, six smears were classified as benign by the RosettaGX microRNA classifier, and one case was designated as suspicious. RosettaGX showed a 75% positive predictive value in comparison to 60% for ThyGenX/ThyraMIR, and both tests demonstrated a 100% NPV.

Next-generation-sequencing (NGS) tests that identify variants in genes associated with thyroid cancer have also been used to help resolve the clinical dilemma presented by indeterminate cytology on thyroid nodules. At least seven genes have been found to have a high degree of specificity for thyroid malignancy, including BRAF, RAS, HRAS, and NRAS mutations and the gene fusions RET/PTC1, RET/PTC3 and PAX8/PPAR γ (Zhang and Linm 2016, Beaudenon-Huibregtse et al., 2014). Rapid technological advances have allowed laboratories the opportunity to add many more genes to their sequencing platforms and may additionally analyze micro-RNA simultaneously. For example, the ThyroSeq v3 assay analyzes 112 genes, providing information on >12,000 mutation hotspots and >120 gene fusion types. In a publication describing the validation of the assay, Nikiforova et al. (2018) reported that in a training set of 238 tissue samples and 175 FNA samples with known surgical follow-up, the test was able to distinguish cancer from benign tissue nodules with 93.9% sensitivity, 89.4% specificity, and 92.1% accuracy. In FNA the authors report a sensitivity of 98.0%, a specificity of 81.8%, and accuracy of 90.9%. Additional studies are necessary to determine the real-world analytical validity and clinical utility of this test.

In a cross-sectional cohort study, Duick et al. (2012) demonstrated that obtaining a GEC test (Afirma) in patients with cytologically indeterminate nodules was associated with a reduction in the rate of diagnostic thyroidectomies. The authors reported that approximately one surgery was avoided for every two GEC tests run on thyroid fine-needle aspirations (FNA) with indeterminate cytology. Data was contributed retrospectively by 51 endocrinologists at 21 practice sites. Compared to a 74% previous historical rate of surgery for cytologically indeterminate nodules, the operative rate fell to 7.6% during the period that GEC tests were obtained. The rate of surgery on cytologically indeterminate nodules that were benign by the GEC reading did not differ from the historically reported rate of operation on cytologically benign nodules. The four primary reasons reported by the physicians for operating on nodules with a benign GEC reading were, in descending order, large nodule size (46.4%), symptomatic nodules (25.0%), rapidly growing nodules (10.7%) or a second suspicious or malignant nodule in the same patient (10.7%). According to the authors, these reasons are concordant with those typically given for operation on cytologically benign nodules.

In a retrospective analysis of 189 thyroid FNAs with indeterminate cytology, Yang et al. (2016) examined the refining role of the Afirma GEC test in a 20-month period after implementation. Correlation with surgical follow-up, when available, was performed. The excisional rate of atypia of undetermined significance-follicular lesion of undetermined significance in the pre-GEC category was 63%, which decreased to 35% in the post-GEC category, whereas the malignancy rate in the excised thyroids increased from 35% in the pre-GEC category to 47% in the post-GEC category. Similar findings also were obtained for suspicious for follicular neoplasm-follicular neoplasm lesions. The authors concluded that the strength of the GEC test appears to lie in its ability to reclassify 42% of indeterminate cytology cases as benign, thereby decreasing the number of unnecessary surgical procedures.

Pagan et al. (2016) investigated the prevalence of genetic alterations in diverse subtypes of thyroid nodules beyond papillary thyroid carcinomas (PTC) in 851 variants and 133 fusions in 524 genes. After adding a cohort of tissue samples, the authors found 38/76 (50%) of histopathology malignant samples and 15/75 (20%) of benign samples to harbor a genetic alteration. In a direct comparison of the same FNA also tested by an RNA-based gene expression classifier (GEC), the sensitivity of genetic alterations alone was 42%, compared to the 91% sensitivity achieved by the GEC. The specificity based only on genetic alterations was 84%, compared to 77% specificity with the GEC. Due to the finding that variants are also found in benign nodules, the authors conclude that testing only GEC suspicious nodules may be helpful in avoiding false positives and altering the extent of treatment when selected mutations are found.

Sipos et al. (2016) retrospectively evaluated the long-term follow-up of patients with a 'benign' Afirma GEC to determine impact on management compared to published data. During 36 months of follow-up, 17 of 98 patients (17.3%) had thyroid surgery; the majority (88%) being performed within two years. According to the authors, this represents a reduction in thyroid surgeries compared to patients that did not have a GEC performed on suspicious lesions. Limitations of this study are small patient population and non-randomization of patients.

MicroRNAs (miRNA) are small noncoding RNAs that regulate gene expression. Research has demonstrated that a number of miRNAs are differentially expressed between benign and malignant thyroid nodules which have led to the development of miRNA based diagnostic lab tests, and in some cases, labs may offer miRNA testing in conjunction with gene variant and expression analysis. Wylie et al. (2016) conducted a study examining genetic variant and miRNA analysis on archived pathology samples from the University of Michigan. The samples consisted of an initial set of 235 aspirates representing 118 nodules with benign cytology, including 13 with surgical outcome (12 benign, 1 malignant), 73 with malignant cytology, including 51 with surgical outcome (1 benign, 50 malignant), and 44 with indeterminate cytology, all with available surgical outcome. The second set of aspirates consisted of 42 distinct nodules with indeterminate cytology and surgical outcome. Thirty-one miRNAs were analyzed as well as 17 genetic alterations in the BRAF, RAS, RET and PAX8 genes, considered standard mutation testing. Furthermore, 54 samples that were negative by the 17-mutation panel were interrogated using a miRNA classification algorithm, commercially available as the ThyraMIR Thyroid miRNA Classifier, which analyzes in parallel 20 genes through next generation sequencing and 46 mRNA transcripts. The authors found that standard mutation testing alone had a sensitivity of 61%, consistent with the literature. Machine learning was utilized to group miRNA analysis into two groups of miRNAs, classifier A and classifier B. When miRNA classifier A was included in the analysis, the sensitivity rose to 78%, and 94% with classifier B. The authors calculated that this leads to a low residual risk of cancer (8%) among specimens negative by mutation and miRNA testing and corresponds to a calculated improvement from 78–90% NPV to 94–98% NPV at 20–40% cancer prevalence. These results contributed to the development of ThyraMIR. In the small cohort that underwent evaluation by ThyraMIR, the authors report a diagnostic sensitivity of 85% and specificity of 95%.

Labourier et al (2015) studied surgical specimens and preoperative FNAs (n=638) for 17 validated gene alterations in the BRAF, RAS, RET and PAX8 genes combined with a 10-miRNA gene expression classifier that provided positive (malignant) or negative (benign) results. Mutations were detected in 69% of nodules with malignant outcome. Among mutation-negative specimens, miRNA testing correctly identified 64% of malignant cases and 98% of benign cases. The authors reported the diagnostic sensitivity and specificity of the combined algorithm was 89% and 85%, respectively. They calculated that with a thyroid cancer prevalence of 32%, the NPV would be 94%, and could help reduce unnecessary surgeries by 69%.

The National Comprehensive Cancer Network (NCCN) guidelines for Thyroid Carcinoma (NCCN, 2019i) recommend molecular profiling for thyroid nodules with indeterminate or suspicious for follicular neoplasm cytology. They note to use molecular markers with caveat and caution. Molecular profiling is not recommended for Hurthle cell neoplasms. Molecular testing of single genes, especially BRAF, or a multigene panel that includes BRAF, NRAS, HRAS, KRAS, RET/PTC1, RET/PTC3, and PAX8/PPAR γ or a gene expression classifier test may be considered, and should be selected by the clinician based on the clinical question being asked.

Clinical Practice Guidelines

American Thyroid Association (ATA)

In this guideline on the clinical management of thyroid nodules, Haugen et al. (2016) provide the following recommendations regarding the use of molecular profiling:

- Nondiagnostic Cytology – Some studies suggest that use of a thyroid core needle biopsy with *BRAF* testing, a gene panel, or a gene expression analysis may provide clinical guidance in these cases, but the full clinical impact of these approaches for nodules with nondiagnostic cytology remains unknown. If molecular testing is being considered, patients should be counseled regarding the potential benefits and limitations of testing and about the possible uncertainties in the therapeutic and long-term clinical implications of results.
- Atypia of Undetermined Significance/Follicular Lesion of Undetermined Significance (AUS/FLUS) – Investigations such as repeat FNA or molecular testing may be used to supplement malignancy risk assessment in lieu of proceeding directly with a strategy of either surveillance or diagnostic surgery. Informed patient preference and feasibility should be considered in clinical decision-making. The authors reviewed available data for multi-gene panels of BRAF, NRAS, HRAS, and KRAS point mutations, as well as RET/PTC1 and RET/PTC3, with or without PAX8/PPAR γ rearrangements, and a mRNA expression profile of 167 genes, and concluded that more data was needed to fully understand how such tests can impact

clinical management. They conclude that there is currently no single optimal molecular test that can definitively rule in or rule out malignancy in all cases of indeterminate cytology.

- Follicular Neoplasm/Suspicious for Follicular Neoplasm Cytology – After consideration of clinical and sonographic features, molecular testing may be used to supplement malignancy risk assessment data in lieu of proceeding directly with surgery.
- Suspicious for Malignant Cytology – After consideration of clinical and sonographic features, mutational testing for *BRAF* or the seven-gene mutation marker panel (*BRAF*, *RAS*, *RET/PTC*, *PAX8/PPAR γ*) may be considered in nodules with SUSP cytology if such data would be expected to alter surgical decision-making. Molecular testing using the 167 GEC has a PPV that is similar to cytology alone (76%) and a NPV of 85% and it is therefore not indicated in patients with this cytological diagnosis.
- Malignant Cytology – While studies have been presented in the literature that suggest that *BRAF* and other multi-gene panels may be useful in prognosis and treatment decisions, more studies are needed to establish the impact of molecular profiling involving multiple mutations or other genetic alterations on clinical management of patients with primary thyroid medullary cancer.
- Post-Operative Radioiodine (RAI) Therapy – Molecular testing to guide postoperative RAI use is not recommended at this time.

American Association of Clinical Endocrinologists, American College of Endocrinology, and Associazione Medici Endocrinologi (AAACE/ACE/AME)

The AAACE/ACE/AME updated their guidelines on the management of thyroid nodules in 2016 (Gharib et al., 2016). They state that molecular profiling should be considered in nodules with indeterminate cytology, and not in those who are found to be clearly benign or malignant. They favor profiles that include *BRAF*, *RET/PTC*, *PAX8/PPARG* and *RAS* mutations. They find that there is insufficient evidence either for, or against, gene expression classifiers. There is insufficient evidence to use molecular profiling to determine the extent of surgical interventions, or for use with low risk indeterminate cytology cases.

Hematological Malignancies

Leukemias

Peterson et al (2015) conducted a study to determine the clinical utility and diagnostic yield, plus examine the rationale, of including microarray analysis in the diagnosis of hematological neoplasias. Twenty-seven patients with hematological malignancies were evaluated by chromosome analysis, FISH and CGH or CGH+SNP arrays. Nearly 90% of chromosome abnormalities found in the patients were also identified by microarray. Of 183 CNVs found, 52% were additional anomalies that were not found by routine cytogenetics or FISH. 65% were <10 Mb in size. Balanced rearrangements were not found by microarray, but of 19 rearrangements that appeared “balanced” by routine cytogenetics, seven had alterations found by microarray at the breakpoints. The authors concluded that CGH provided clinicians with advantages in identification of cryptic imbalances and clonal abnormalities in non-dividing cells with poor chromosome morphology and therefore had potential to be integrated as a patient management tool.

Laurie et al. (2015) compared the SNP array results of 278 symptomatic CLL patients with >50,000 subjects from the GENEVA consortium of genome wide association studies, which analyzed people with a range of medical conditions and healthy controls. The CLL patients were also analyzed by FISH to determine performance and concordance between the SNP array and FISH. When a parameter of 20% abnormal cells was used as a cutoff, the concordance rate between the SNP array and FISH was 98.9%. The array found 8.4% of cases with UPD which cannot be detected by FISH. In 214 CLL patients with SNP results, 1,112 genetic anomalies were found, of which 628 were considered acquired. This was a higher percentage and anomalies were unique in the CLL group when compared to the GENEVA cohort and suggests that late stage CLL has recurrent acquired anomalies that do not occur in precursor conditions or in the general population. The clinical significance of this finding is not clear, however, SNP based array was demonstrated to be a valid analysis tool.

Koh et al. (2014) utilized a CGH+SNP array platform to study the presence of CNVs and LOH in 15 children with acute myeloid leukemia (AML) and three with myelodysplastic syndrome (MDS). Cytogenetic analysis revealed CNV in 11 regions in eight patients. SNP+CGH found 14 CNV in nine patients, and cryptic LOHs in three of five patients with normal cytogenetics. Overall, nine patients were found to have abnormalities not detected by routine cytogenetics. three patients with AML and terminal LOH of >10Mb had significantly inferior relapse-free survival time, suggesting that SNP+CGH testing can provide additional prognostic information.

Puiggros et al. (2012) studied 70 patients with chronic lymphocytic leukemia (CLL) by routine cytogenetics, FISH, and genomic arrays to determine if genomic arrays could replace current testing standards. Routine cytogenetics found 31% genomic anomalies in patients, and FISH found 69%. Genomic arrays, CytoScan HD Array and CytoScan HD Array, found anomalies in 79% and 80%, respectively. Arrays missed small deletions at 11q and 17p due to their limited sensitivity in these regions. The authors concluded that arrays should remain a complementary tool to routine cytogenetics and FISH to prevent a negative impact on patients who harbor genetic anomalies that would be missed by this technology.

Hagenkord et al. (2010) examined the optimal SNP array probe density for clinical use in CLL to identify actionable genetic variation missed by FISH and conventional chromosome analysis. The validation cohort consisted of 18 archived sample and 11 clinical samples that were simultaneously tested with standard FISH for CLL. Where possible, cytogenetic and flow cytometry was also performed. Affymetrix SNP arrays of low (10K2.0), medium (250K Nsp) and high (SNP6.0) density were utilized. Ultimately the medium density array was validated for clinical use and was found in 98.5% concordance with standard FISH. In particular, a region of acquired uniparental disomy (UPD) with two mutation copies of TP53 was identified that was not found by FISH or routine cytogenetics. The authors concluded that SNP array karyotyping provides high resolution CNV analysis, identification of UPD and detects lesions missed by FISH.

Boulwood et al. (2010) used a SNP array to analyze 41 chronic myeloid leukemia (CML) patients using 53 bone marrow or blood sample. 32 were in chronic phase and 21 were in blast crises. The samples were analyzed for uniparental disomy (UPD) and copy number variants, with quality control comparisons with 100 healthy controls of different ethnicities for SNP array hybridization intensities, and 45 healthy controls as a reference set. Across the samples 44 regions of UPD were identified, with chromosome 8 having the highest frequency. 10 regions of copy number variation were identified in four of 21 patients with blast crises, and none were observed for those in chronic phase. The authors noted that 32 regions of UPD were noted in 23 of 45 healthy controls on chromosomes 15 and 22. Therefore only regions of UPD were reported for CML patients that weren't found in the controls, and this emphasized to the authors that SNP analysis, particularly for UPD, requires inclusion of constitutional controls. UPD is not identifiable by other testing methods but is important as the acquired homozygosity of disease genes may contribute to disease progression. In this cohort, UPD was found in one patient at 20q11 that includes the ASXL1 gene, a tumor suppressor gene associated with early events in CML. Sequencing exon 12 in all patients found that 6 of 41 had ASXL1 mutations, which is likely a newly identified molecular abnormality for CML.

Clinical Practice Guidelines

College of American Pathologists (CAP) and American Society of Hematology (ASH)

CAP and ASH convened a panel of experts to review the literature and establish a guideline for appropriate lab testing for the initial diagnosis of acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL) and ambiguous acute leukemias (ALs). The experts reviewed the literature and using an evidence-based methodology intended to meet recommendations from the Institute of Medicine, a set of guidelines was developed. The guidelines were reviewed by an independent panel and were made available for public comment. The outcome was 27 guidelines addressing clinical information required by the pathologist and recommended laboratory testing. Chromosome microarray is broadly addressed as one potential test in several statements that refer to "molecular genetic testing," which may also include FISH, RT-PCR, or DNA methylation studies. These include:

- "In addition to morphologic assessment (blood and BM), the pathologist or treating clinician should obtain sufficient samples and perform conventional cytogenetic analysis (i.e., karyotype), appropriate molecular-genetic and/or FISH testing, and FCI. The flow cytometry panel should be sufficient to distinguish acute myeloid leukemia (including acute promyelocytic leukemia), T-ALL (including early T-cell precursor leukemias), B-cell precursor ALL (B-ALL), and AL of ambiguous lineage for all patients diagnosed with AL. Molecular genetic and/or FISH testing does not, however, replace conventional cytogenetic analysis." [Statement 5. Strong Recommendation].
- "For patients who present with extramedullary disease without BM or blood involvement, the pathologist should evaluate a tissue biopsy and process it for morphologic, immunophenotypic, cytogenetic, and molecular genetic studies, as recommended for the BM." [Statement 11. Strong Recommendation].
- "For patients with suspected or confirmed AL, the pathologist or treating clinician should ensure that flow cytometry analysis or molecular characterization is comprehensive enough to allow subsequent detection of MRD". [Statement 12. Strong Recommendation] (Arber et al., 2017).

Myelodysplastic Syndrome

Song et al. (2017a) conducted a review of the literature comparing the clinical utility of a variety of genomic profiling techniques in the treatment of myelodysplasias (MDS). They noted that the common defects in MDS that should be identified are del5q,

trisomy 8, del20q, del7q, monosomy 7 and complex karyotypes. Each aberration has different prognostic and management challenges, so accurate identification of genomic abnormalities is important for a clear diagnosis and to optimize treatment strategies. The authors compared findings from the literature for routine cytogenetics, FISH, spectral karyotyping (SKY), SNP array, CGH, and SNP+CGH for the ability to detect the common defects in MDS. The authors concluded that no single technology provides all the information necessary for the clinician to create informed treatment plans, and that a combination of techniques is required. The authors favored routine cytogenetics, FISH and SNP+CGH, but noted that additional efforts are needed to standardize testing and bioinformatics, and further technological advances are needed to overcome the limitations of diverse techniques.

Evans et al. (2016) studied the diagnostic utility of SNP+CGH array to identify unexplained cytopenia in 83 MDS patients and compared results with 18 normal bone marrow controls. Array analysis was done in parallel with standard cytogenetics, FISH, flow cytometry, and morphology. Forty-five percent of patients were diagnosed with MDS, 33% were normal, and 8% had other pathological disorders. 57% of the MDS patients had normal cytogenetics, but the SNP+CGH array found significant cryptic chromosome aberrations. In MDS patients with abnormal cytogenetics, the array essentially matched the chromosome results and didn't add any new information. Overall, the SNP+CGH array analysis contributed significantly to the diagnostic yield in indeterminate morphology cytopenic patients.

Kolquist et al. (2011) examined the clinical utility of CGH in myelodysplasias. They noted that only half of myelodysplasias (MDS) patients show genomic abnormalities using routine cytogenetics, yet this group of patients is characterized by ineffective hematopoiesis, cytopenia, and a 30% risk of developing acute myeloid leukemia (AML). They hypothesized that using CGH to test patients who were cytogenetically normal would reveal cryptic genomic alternations that would improve prognosis, managing disease progression, and determining the suitability and efficacy of molecularly targeted therapy. They analyzed 35 samples by CGH derived from patients with a diagnosis and suspicion of MDS who also had known abnormal karyotypes. 80% of samples had new chromosomal aberrations that had not been revealed by cytogenetics or FISH. An additional 132 cryptic abnormalities were found including deletions of known oncogenes, such as NF1, RUNX1, RASSF1, CCND1, TET2, DNMT3A, HRAS, PDGFRA and FIP1L1. Overall, the authors concluded that CGH in combination with routine cytogenetics provided additional clinically relevant information that could better direct the care of the patients analyzed.

Thiel et al. (2011) notes that 40% of those with MDS have a normal karyotype and may have a different prognosis than those who have an abnormal karyotype. The availability of CGH now allows for the identification of cryptic genomic abnormalities and having this information may have a prognostic or treatment impact. They studied 107 MDS patients with a normal karyotype and found that 39% of patients had cryptic genomic imbalances, including regions that are known to be impacted in MDS such as del4q, del5q, and del7q. Most alterations were verified by other methods. Overall, these patients had inferior survival and outcomes similar to those with cytogenetically visible aberrations when compared to the rest of the patients in this cohort with no identifiable cytogenetic abnormalities.

Multiple Myeloma

Weinhold et al. (2016) reported clinical outcomes of GEP testing in relation to treatment type for subgroups of patients (n=1217) with multiple myeloma (MM) who participated in the University of Arkansas for Medical Sciences Total Therapy (TT) trials. Using log-rank tests for GEP data, the researchers identified 70 genes linked to early disease-related death. The UAMS GEP70 risk score is based on the ratio of the mean expression level of up-regulated to down-regulated genes among the 70 genes. Most up-regulated genes are located on chromosome 1q, and many down-regulated genes map to chromosome 1p. The predictor enabled the reliable identification of patients with shorter durations of complete remission, event-free survival, and overall survival that constitute 10 – 15% of newly diagnosed MM patients. The authors' reported that impact of treatment differs between molecular subtypes of MM and that GEP gives important information that can help in clinical decision-making and treatment selection. Future studies should address whether strategies maximizing exposure to proteasome-inhibitors can further improve outcome in the MM subgroup. The authors' note that comparison of GEP data of multiple paired samples showed differences in risk signatures, indicating the co-existence of HiR and LoR subclones (manuscript in preparation). Possibly, cells of a LoR subclone were collected at relapse in these patients. The addition of thalidomide significantly improved outcome of LoR cases from maintenance and that outcome of LoR was improved further by the addition of bortezomib. The authors comment that they could not detect a significant improvement for HiR cases, but this may be due to a lack of statistical power.

Tiu et al. (2011) examined the analytical validity and clinical utility of SNP arrays in individuals with myelodysplastic syndromes when performed in parallel with cytogenetics vs. cytogenetics alone. They analyzed 430 patients within the MDS spectrum

which included 250 with MDS, 95 with MDS/myeloproliferative overlap neoplasm, and 85 with acute subsequent AML. Overall, the combined SNP array+karyotype had a higher diagnostic yield of chromosomal defects at 74%, compared to karyotype alone at 44%. Novel lesions were identified by array in 54% with normal cytogenetics and 62% of those with abnormal cytogenetics. The presence and number of SNP identified lesions proved to be an independent predictor of outcome and tended to have worse survival outcomes. The authors concluded that concurrent use of routine cytogenetics with a SNP array improves diagnostic yield and prognostic information compared to cytogenetics alone.

NCCN clinical practice guidelines for multiple myeloma state that gene expression profiling (GEP) has the potential to provide additional prognostic value to further refine risk-stratification, help therapeutic decisions and inform novel drug design and development. The NCCN panel unanimously agreed that although GEP is not routinely used in clinical practice during diagnostic workup, it may be helpful in selected patients to estimate the aggressiveness of the disease and individualize treatment. No patient selection criteria were provided (NCCN, 2019d).

Detection of Minimal Residual Disease (MRD) in Hematologic Malignancies

Ladetto et al. (2014) compared real time quantitative polymerase chain reaction (RQ-PCR) to NGS for identifying clonotype identification, clonotype identity and comparability of MRD results. A total of 378 samples from 55 patients with acute lymphoblastic leukemia (ALL), mantle cell lymphoma (MCL) or multiple myeloma (MM) were analyzed. RQ-PCR identified 45 clonotypes, and NGS found 49, and were identical or >97% homologous in all cases. Both consistently had a sensitivity level of 1×10^{-5} and MRD results were concordant in 79.6% of cases. NGS showed at least the same level of sensitivity as RQ-PCR without the need for patient specific reagents, and may be a useful tool for monitoring in ALL, MCL and MM.

Avet-Loiseau et al. (2015) reported on the use of FC and NGS in the Intergroupe Francophone du Myélome/ Dana-Farber Cancer Institute (IFM/DFCI) 2009 trial to measure MRD in the IFM arm of the study. This trial enrolled 700 patients under 66 years of age and randomized them to either receive either 8 cycles of VRD (Velcade-Revlimid-Dexamethasone) (arm A), or 3 VRD cycles, high-dose melphalan, followed by two consolidation VRD cycles (arm B). All patients received a lenalidomide maintenance for 12 months. A total of 246 patients were evaluated by NGS using the LymphoSight platform, and before maintenance, 87 patients were negative, 80 were low-positive, and 79 were positive. After maintenance, 178 were tested, and 86 patients were negative, 52 were low-positive, and 40 were positive. Using a cutoff of 10^{-6} , patients below this threshold had a pre-maintenance progression free survival (PFS) of 86%, vs 53% for patient $>10^{-6}$. In the post-maintenance group, these numbers were 90% and 59% respectively. When compared with results from 7 color FC, of 72 patients who were positive with FC, 67 were also positive with NGS. In the FC negative group, of the 163 patients, 51 were positive by NGS. In this subgroup, the 3-year PFS was 86% for the NGS negative patients compared to 66% for the NGS negative patients in the pre-maintenance group. In the post-maintenance group, the numbers were 91% and 65% respectively. The authors concluded that NGS was able to predict PFS in this study.

The efficacy of targeted NGS to identify MRD in patients with acute myeloid leukemia (AML) was studied by Jongen-Lavrencic et al. (2018). Between 2001 and 2013, a total of 482 patients ranging in age from 18-65 with newly diagnosed AML were included. NGS of 54 genes that are often present in AML patients was performed at diagnosis and after induction therapy during complete remission. The end points analyzed were four-year relapse, relapse free survival and overall survival. Results were compared with flow cytometry (FC). The authors discovered an average of 2.9 mutations per patient, of which at least one single mutation could serve as an indicator of residual disease, in 430 patients. These patients then had NGS testing repeated on bone marrow after induction therapy and they were in complete remission. Persistent mutations were found in 51.4% and were highly variable across the genes analyzed. DTA mutations were most common, persisting at rates of 78.7%, whereas *RAS* pathway mutations cleared, persisting at an average rate of about 9%. The authors noted that DTA mutations are common gene mutations in individuals with age related clonal hematopoiesis, and likely represent non-leukemic clones rather than persistent malignant disease. After DTA mutations were excluded, the detection of MRD was associated with a significantly higher relapse rate than no detection (55% vs. 32%), lower relapse-free survival (37% vs. 58%) and overall survival (42% vs. 66%). The results of NGS were compared to FC in a subset of 340 patients. Concordant results for detection or non-detection of MRD were found in 69% of patients. The four year relapse rate was 73% among patients in whom both assays were positive, 52% among those who had residual disease on sequencing but not on flow cytometry, 49% among those who had residual disease on flow cytometry but not on sequencing, and 27% among those in whom both assays were negative. Multivariate analysis found that combining the two assays gave a high prognostic value to the rate of relapse ($p < .001$), relapse free survival ($p < .001$) and overall survival ($p = .003$). The authors concluded that persistent mutations associated with clonal hematopoiesis did not have prognostic value, whereas the detection of MRD during complete remission using NGS with FC had significant additive prognostic value.

The Food and Drug Administration (FDA) reviewed data submitted by Adaptive Technologies on their ClonoSeq assay, which included data from currently ongoing studies (FDA, 2018). They noted that clinical validity was demonstrated in a retrospective analysis of 273 patients with ALL, on ongoing study of 323 patients with multiple myeloma, and separate study of 706 patients with multiple myeloma. Patients who had a negative MRD results had a longer event free survival.

An important prognostic factor in B-lymphoblastic leukemia (B-ALL) is early response to combination induction chemotherapy. End of induction response is typically measured by multiparametric flow cytometry (FC) or allele-specific oligonucleotide polymerase chain reaction (ASO-PCR). The analytical sensitivity for FC is 0.01%, and ASO-PCR is .001%, but requires the development of patient specific probes. Wood et al. (2018) reviewed the clinical validity of a new technical approach of using high throughput sequencing (HTS) of IGH and TRG genes to FC for determining minimal residual disease (MRD). The study used 619 paired pretreatment and end-of-induction bone marrow samples from Children's Oncology Group studies AALL0331 and AALL0232 (clinicaltrials.gov). The samples were evaluated by HTS and FC for event free survival and overall survival. Using an MRD threshold of 0.01%, HTS and FC show similar 5-year event free survival and overall survival rates. There was high discordance between HTS and FC in number of patients identified; HTS identified 55 more patients (38.7%), and these patients had worse outcomes than FC MRD negative patients. HTS also identified 19% of standard risk patients without MRD at any detectable level, which was correlated with excellent outcomes. Overall HTS had a high sensitivity and lower false-negative rate than FC in this analysis.

Determining the response to treatment is an important aspect of managing multiple myeloma, and NCCN guidelines recently adding assessing MRD to the management algorithm. NCCN notes that a validated next generation sequencing assay or next generation flow could be used for determining MRD. The ideal time is after each treatment stage for individuals that have undergone autologous or allogenic bone marrow transplant. Two consecutive assessments are not necessary, one test is sufficient after each treatment stage. Only individuals who appear to have complete response and have no evidence of progression or new bone lesions should have MRD assessment (NCCN, 2018a). NCCN does not have the same stance on using NGS for MRD detection in AML (NCCN AML 2019). They note that the use of MRD as a management or prognostic tool for AML is still emerging, and that common assessments are RT-PCR for NPM1, CBFβ-MYH11, and RUNX1-RUNX1t1 and flow cytometry. The sensitivity of these two approaches is considered superior at this time to NGS using targeted panels of 20-50 genes

Other Cancers and Clinical Indications

Molecular profiling has many theoretical clinical applications in the field of oncology. Published clinical studies have addressed the use of molecular profiling for the following:

- Acute myeloid leukemia (Port et al., 2014; Link et al., 2012)
- Adrenocortical cancer (Zheng et al., 2016; Ross et al., 2014a)
- Breast cancer (Ganesan et al., 2014; Wheler et al., 2014)
- Circulating tumor cells (Yang et al., 2018; Merker et al., 2018)
- Gastric and gastrointestinal cancer (West et al., 2017; Ali et al., 2015, Vignot et al., 2015; Miura et al., 2014)
- Head and neck cancer (Wang et al., 2017; Chung et al., 2015)
- Gynecological cancer (Rodriguez-Rodriguez et al., 2016; Ross et al., 2013)
- Non-melanoma skin cancers
- Pancreatic cancer (Zhou et al., 2017; Chmielecki et al., 2014; Chantrill et al., 2015)
- Urothelial carcinoma/urinary bladder adenocarcinoma (Roy et al., 2017; Ross et al., 2014b; Millis et al., 2015)

There is insufficient published evidence to support the use of molecular profiling for these cancers, technologies or sample types. The main evidence deficiencies are insufficient data on analytical validity, clinical validity, and clinical utility.

Hirshfield et al. (2016) conducted a prospective clinical study on 100 patients with diverse-histology, rare, or poor-prognosis cancers to evaluate the clinical implications of a comprehensive genomic profiling assay (FoundationOne), using formalin-fixed, paraffin-embedded tumors. The primary objectives were to assess utility, feasibility, and limitations of genomic sequencing for genomically guided therapy or other clinical purpose in the setting of a multidisciplinary molecular tumor board. Of the tumors from the 92 patients with sufficient tissue, 88 (96%) had at least one genomic alteration (average 3.6, range 0–10). Use of comprehensive profiling led to implementable clinical action in 35% of tumors with genomic alterations, including genomically guided therapy, diagnostic modification, and trigger for germline genetic testing. Although use of targeted next-generation sequencing in the setting of an institutional molecular tumor board led to implementable clinical action in more than one third of patients with rare and poor-prognosis cancers, major barriers to implementation of genomically guided therapy were clinical

status of the patient and drug access. Early and serial sequencing in the clinical course and expanded access to genomically guided early-phase clinical trials and targeted agents may increase clinical application.

Kato et al. (2015) investigated the clinical correlates of CDK4/6 and CDKN2A/B abnormalities in diverse malignancies. Patients with various cancers who underwent molecular profiling by targeted next generation sequencing (Foundation Medicine; 182 or 236 cancer-related genes) were reviewed. Of 347 patients analyzed, 79 (22.8%) had aberrant CDK 4/6 or CDKN2A/B. Only TP53 mutations occurred more frequently than those in CDK elements. Aberrations were most frequent in glioblastomas (21/26 patients; 81%) and least frequent in colorectal cancers (0/26 patients). Aberrant CDK elements were independently associated with EGFR and ARID1A gene abnormalities. CDK aberrations were associated with poor overall survival. In multivariate analysis, PTEN and TP53 aberrations were independently associated with poorer survival; CDK aberrations showed a trend toward worse survival. There was also a trend toward worse progression-free survival (PFS) with platinum-containing regimens in patients with abnormal CDK elements (3.5 versus 5.0 months). In conclusion, aberrations in the CDK pathway were some of the most common in cancer and independently associated with EGFR and ARID1A alterations. Patients with abnormal CDK pathway genes showed a trend toward poorer survival, as well as worse PFS on platinum-containing regimens. According to the authors, further investigation of the prognostic and predictive impact of CDK alterations across cancers is warranted. This study was limited due to it being performed retrospectively in a single institution with a relatively limited number of patients.

Johnson et al. (2014) retrospectively assessed demographics, next-generation sequencing (NGS) results, and therapies received for patients undergoing targeted NGS using the FoundationOne test. Co-primary endpoints were the percentage of patients with targeted therapy options uncovered by mutational profiling and the percentage who received genotype-directed therapy. Samples from 103 patients were tested; most frequently breast carcinoma (26%), head and neck cancers (23%), and melanoma (10%). Most patients (83%) were found to harbor potentially actionable genetic alterations, involving cell-cycle regulation (44%), phosphatidylinositol 3-kinase-AKT (31%), and mitogen-activated protein kinase (19%) pathways. With median follow-up of 4.1 months, 21% received genotype-directed treatments, most in clinical trials (61%), leading to significant benefit in several cases. The most common reasons for not receiving genotype-directed therapy were selection of standard therapy (35%) and clinical deterioration (13%). The authors concluded that mutational profiling using a targeted NGS panel identified potentially actionable alterations in a majority of advanced cancer patients. The assay identified additional therapeutic options and facilitated clinical trial enrollment. According to the authors, there are many unanswered questions regarding implementation of this technology. First, based on this study, some patients with potentially actionable alterations did not respond to genotype-directed therapy, highlighting the still underdeveloped understanding of the pathophysiologic implications of many genetic alterations. Second, the most appropriate indications for obtaining targeted NGS are not yet clear. Third, randomized studies in the future will need to assess whether targeted NGS improves overall outcomes.

Frampton and colleagues (2013) conducted an analytical and clinical validation study to evaluate massively parallel DNA sequencing using the FoundationOne assay to characterize base substitutions, indels, copy number alterations, and selected fusions across 287 cancer-related genes from routine formalin-fixed and paraffin-embedded (FFPE) clinical specimens. The authors implemented a validation strategy with reference samples of pooled cell lines that modeled key drivers of test accuracy, including mutant allele frequency, indel length and amplitude of copy change. Test sensitivity achieved was 95% to 99% across alteration types, with high specificity (positive predictive value [PPV] >99%). The authors confirmed accuracy using 249 FFPE cancer specimens characterized by established assays. Application of the test to 2,221 clinical cases revealed clinically actionable alterations in 76% of tumors, three times the number of actionable alterations detected by current diagnostic tests. This study did not evaluate the clinical utility of such findings in improving care and outcome of patients by tailoring treatments or predicting response to treatment. Hence, it is important to note that the clinical utility of genomic profiling using massively parallel DNA sequencing remains unknown. In addition, study authors colleagues did not categorize the data regarding sensitivity, specificity, and positive predictive value (PPV) by cancer type.

O’Kane et al. (2019) reported on the COMPASS trial for pancreatic ductal adenocarcinoma (PDAC). Patients were recruited before chemotherapy for whole genome sequencing (WGS) and RNA sequencing (RNASeq). The tumor tissue was analyzed, and tumor responses and clinical outcomes were correlated. Of the 157 patients that had a tumor biopsy, 141 genomes were reported. Twenty-five (21%) had a Moffitt basal-like RNA signature which is usually associated with chemotherapy resistance. GATA6 expression was able to separate the Moffitt subgroup from those with classical tumors. Also, 30% of patients had potentially actionable genetic alterations including BRAF variants (n=4) and a NTRK3-EML4 fusion in *KRAS*WT tumors (8%). The researchers concluded that there are subsets of patients with advanced PDAC that have actionable variants.

Another COMPASS trial publication described the use of real-time WGS and RNASeq of advanced PDAC to identify predictive mutational and transcriptional features for better treatment selection (Aung et al., 2018). Sixty-three patients underwent a tumor biopsy and WGS and RNASeq were successful in 62 (98%) and 60 (95%), respectively. PDAC RNA subtypes were compared to basal-like subtypes for chemotherapy response. GATA6 expression in tumor measured by RNA in situ hybridization was found to be a robust surrogate biomarker for differentiating classical and basal-like PDAC subtypes. These potentially actionable genetic alterations were found in 30% of patients.

Singhi et al. (2018) studied the clinical validity of using pre-operative pancreatic cyst fluid (PCF) for next generation sequencing (NGS) of KRAS, GNAS, TP53, PIK3CA and PTEN genes in order to predict benign vs. malignant lesions. PCF samples from 595 patients (626 samples) were obtained through fine needle aspiration and subjected to NGS for the five genes. A different cohort of 159 PCF specimens was also evaluated for KRAS/GNAS mutations by Sanger sequencing. Of the 595 patients, 308 (49%) had KRAS or GNAS mutations and 35 had a mutation in TP53, PIK3CA, or PTEN. Follow up diagnostic pathology was available in 102 patients. For these 102 patients, NGS testing of PCF for KRAS/GNAS had a 100% sensitivity (n=56) and 96% specificity for an intraductal papillary mucinous neoplasm. In the separate cohort of Sanger sequencing patients, KRAS/GNAS mutations detection had a 65% sensitivity and 100% specificity. By NGS, the combination of KRAS/GNAS mutations and alterations in TP53/PIK3CA/PTEN had an 89% sensitivity and 100% specificity for advanced cancer. The study concluded that in comparison to Sanger sequencing, preoperative NGS of PCF for KRAS/GNAS mutations is highly sensitive for IPMNs and specific for mucinous PCs. In addition, the combination of TP53/PIK3CA/PTEN alterations is a useful preoperative marker for advanced cancer.

In a guideline from ASCO in 2016, clinical decision support was outlined for metastatic pancreatic cancer. Sohal et al. (2018) published an update to this guideline that incorporated new evidence. The researchers conducted a literature review and found two new studies to include. The recommendations included that select patients should be tested for mismatch repair deficiency or microsatellite instability, and pembrolizumab is recommended for patients with mismatch repair deficiency or high microsatellite instability tumors.

Lowery et al. (2018) performed comprehensive germline testing (GT) in a cohort of patients with exocrine pancreatic neoplasms. The genotype and phenotype associations were used to identify biomarkers for therapy response. Six hundred fifteen patients were prospectively tested for somatic tumor and matched sample profiling for 410-468 genes. PGAs were present in 122 (19.8%) of 615 patients involving 24 different genes, including BRCA1/2, ATM, PALB2, and multiple additional genes associated with the DNA damage response pathway. Of these patients with germline alterations, 41.8% did not meet current guidelines for GT. The study concluded that the data supported routinely offering GT in all pancreatic ductal adenocarcinoma patients with a broad panel of known hereditary cancer predisposition genes.

Wong et al. (2019) reported on ampullary cancer (AC) and germline alterations in BRCA2, ERBB2, and ELF3. Forty-five patients with pathologically confirmed AC were tested with the Memorial Sloan Kettering Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT) test (410-468 genes). Twenty-three patients were also tested with GT with MSK-IMPACT (76-88 genes). Eight of 44 patients (18%) were identified as harboring pathogenic mutations in BRCA2, ATM, RAD50, and MUTYH. Additionally, they found a wide spectrum of SAs in genes such as KRAS, MDM2, ERBB2, ELF3, and PIK3CA. Two patients in the cohort underwent SA-targeted therapy, and 1 had a partial radiographic response.

Liquid Biopsy

Liquid biopsy is a non-invasive technique of obtaining bodily fluids, such as blood, urine, cerebrospinal fluid, saliva, and other aspirates, to analyze different types of biomolecules including circulating tumor DNA (ctDNA), circulating tumor cells (CTCs), and exosomes. Liquid biopsies have been investigated for a number of cancer types; however, this testing has not been widely accepted yet. Research continues to study this technique for non-invasive methods that may assist in therapeutic decisions without traditional biopsy.

Cohen et al. (2017) and Su et al. (2018) researched different methods for detection of T790M in plasma cell-free DNA for lung cancer. The researchers used a combination of peptide nucleic acid (PNA) and Matrix-Assisted Laser Desorption/Ionization Time-of-Flight Mass Spectrometry (MALDI-TOF MS) to monitor cell-free DNA T790M in EGFR-mutant patients. The cohort included 103 tumor and cell free DNA T790M samples. Detection sensitivity of cfDNA T790M was 67.4% and overall concordance was 78.6%. Among 65 T790M-positive tumors, 15 were negative in cfDNA (23.1%). Seven of 38 T790M-positive cfDNA samples were negative in the tumors (18.4%).

Oxnard et al (2016) studied whether noninvasive genotyping of cell-free plasma DNA (cfDNA) is a useful biomarker for prediction of outcome from a third-generation EGFR-TKI, osimertinib. All patients had plasma collected and genotyping was performed by using BEAMing. The use of plasma genotyping for detection of T790M had a sensitivity of 70%. Of 58 patients with T790M-negative tumors, T790M was detected in plasma of 18 (31%). This study suggested that the use of plasma T790M assays could help certain patients avoid a tumor biopsy for T790M genotyping. However due to the 30% false-negative rate of plasma genotyping, patients with T790M-negative plasma results still need a tumor biopsy to determine presence or absence of T790M.

Another study also evaluated rapid plasma genotyping for the detection of EGFR and KRAS in advanced lung cancer (Sacher et al. 2016). Blood samples were taken from patients with advanced nonsquamous non-small-cell lung cancer (NSCLC). The patients either (1) had a new diagnosis and were planned for initial therapy or (2) had developed acquired resistance to an EGFR kinase inhibitor and were planned for rebiopsy. Test was performed for EGFR exon 19 del, L858R, T790M, and/or KRAS G12X. All patients underwent biopsy for tissue genotyping, which was used as the reference standard for comparison. Of 180 patients with advanced NSCLC, 120 cases were newly diagnosed; 60 had acquired resistance. Tumor genotype included 80 EGFR exon 19/L858R mutants, 35 EGFR T790M, and 25 KRAS G12X mutants. The plasma test had a positive predictive value of 100% (95% CI, 91%-100%) for EGFR 19 del, 100% (95% CI, 85%-100%) for L858R, and 100% (95% CI, 79%-100%) for KRAS, but lower for T790M at 79% (95% CI, 62%-91%). The sensitivity was 82% (95% CI, 69%-91%) for EGFR 19 del, 74% (95% CI, 55%-88%) for L858R, and 77% (95% CI, 60%-90%) for T790M, but lower for KRAS at 64% (95% CI, 43%-82%). Sensitivity for EGFR or KRAS was higher in patients with multiple metastatic sites and those with hepatic or bone metastases, specifically. The researchers concluded that this rapid plasma testing detected EGFR and KRAS mutations rapidly with high specificity needed to select therapy and avoid repeat biopsies. In addition, this testing may also detect EGFR T790M missed by tissue genotyping due to tumor heterogeneity in resistant disease.

Riediger et al (2016) studied tumors over time through the use of plasma DNA. The researchers aimed to identify early indications for therapy response or tumor progression. Lung adenocarcinoma patients who were treated with TKIs had serial plasma samples taken. Through digital PCR, EGFR and KRAS mutations were quantified in the circulating DNA. The DNA levels were compared to the treatment courses and variations were found in 15 patients. The study concluded that serial assessment of EGFR mutations in the plasma of these lung cancer patients was able to determine treatment response and tumor progression earlier than other methods.

Guardant 360

One liquid biopsy test, Guardant360, evaluates cell-free tumor DNA for 73 different genes. The majority of studies with Guardant360 have focused on NSCLC; however, more research is being performed with other tumor types. A study by Yang, et al (2017) evaluated lung cancer and other solid tumors. Plasma from patients with lung cancer (n=103) and other solid tumors (n=74) was analyzed for ct (DNAs) using the Guardant360 test. In this cohort, mutations in TP53, EGFR, and KRAS genes were most often determined. Mutations in BRCA1, BRCA2, and ATM were found in 18.1% (32/177) of cases. Also, the researchers compared the ctDNA and tumor tissue of 37 lung cancer cases. This analysis found that key mutations could be found in plasma even if they were minor in the tumor tissue.

Dagogo-Jack et al (2019) performed a study on ROS1 fusions in NSCLC with the Guardant360 NGS assay and the Guardant Health plasma dataset. The assay part of the study aimed to detect potential genetic mediators of resistance in the plasma of patients with ROS-1 positive NSCLC who were relapsing on crizotinib. The researchers found that the sensitivity for detection of ROS1 fusions in plasma at relapse on crizotinib therapy was 50%. Of 18 post-crizotinib plasma specimens, 6 (33%) had ROS1 kinase domain mutations (5 were ROS1 G2032R). Two (11%) post-crizotinib plasma specimens had genetic alterations (n=1 each BRAF V600E and PIK3CA E545K). Additionally, the plasma dataset provided by Guardant Health was compared to institutional tissue data. There was 100% concordance between the specific tissue- and plasma-detected ROS1 fusion for seven patients genotyped with both methods.

McCoach et al (2018) evaluated patients with advanced NSCLC and with tumors that carried ALK gene fusions. The researchers sought to analyze the cfDNA to find a non-invasive way to identify these gene fusions. The study used the Guardant360 database of NSCLC cases to identify patients. Eighty-eight patients with 96 plasma detected ALK fusions were determined. The fusion partners identified included *EML4* (85.4%), *STRN* (6%), and *KCNQ, KLC1, KIF5B, PPM1B, and TGF* (totaling 8.3%). The study concluded that in this cohort, cfDNA was acceptable at detecting targetable alterations.

Another study by Lam et al (2019) studied lung squamous-cell carcinoma (LUSC) and cfDNA. The researchers retrospectively evaluated 492 LUSC patients with 410 patients (stage 3B or 4 LUSC) who were tested with a targeted cell-free circulating DNA NGS assay, and 82 patients (any stage) who were tested with a tissue NGS cancer panel. Overall, 467 patients (94.9%) had a diagnosis of LUSC, and 25 patients (5.1%) had mixed histology. Of the LUSC subgroup, a total of 10.5% had somatic alterations with therapeutic relevance, including in EGFR (2.8%), ALK/ROS1 (1.3%), BRAF (1.5%), and MET amplification or exon 14 skipping (5.1%). Three of these patients were treated with targeted therapy and all experienced a partial response. Of the group with mixed histology, 16% had an actionable alteration. The researchers found actionable alterations in genes that were clinically significant through this testing; however, they state that further evaluation is needed.

Kim, et al. (2017) performed a prospective study on solid tumor cancers and ctDNA guided matched therapy. The testing identified point mutations in 70 genes and indels, fusions, and copy number amplifications in selected genes. Alterations in somatic genes was detected in 59 patients with gastric cancer (78%), and 25 patients (33%) had targetable alterations (*ERBB2*, n=11; *MET*, n=5; *FGFR2*, n=3; *PIK3CA*, n=6). In NSCLC, 62 patients (85%) had somatic alterations, and 34 (47%) had targetable alterations (*EGFR*, n=29; *ALK*, n=2; *RET*, n=1; *ERBB2*, n=2). In a small subgroup of patients that had tissue available for confirmation (10 with gastric cancer and 17 with NSCLC), molecularly matched therapy was initiated. The response rate and disease control rate in this group was 67% and 100%, respectively, in gastric cancer and 87% and 100%, respectively, in NSCLC. Response was independent of targeted alteration variant allele fraction in NSCLC ($P = .63$). The researchers concluded that response rates in this analysis were similar to tissue-based targeted therapy studies.

Villafior, et al. (2016) reported on patients with NSCLC undergoing analysis of ctDNA using Guardant360. As part of clinical care, 90 patients submitted for ctDNA testing, but only 68 provided consent. These patients had lung adenocarcinoma (n=55, 81%), lung squamous cell carcinoma (n=12, 17.7%) and other lung cancers (n=1, 1.3%). Of these 68, 38 were tested using the 54-gene ctDNA panel and 31 were analyzed on the 68-gene ctDNA panel. Tissue-based testing was performed on 44 subjects using 9 different testing platforms. The researchers found that 83% of subjects had at least one genomic alteration and the most commonly mutated genes were TP53, KRAS and EGFR. Only 31 patients had matched tissue and blood samples, and, in those patients, an EGFR activating was found in both tissue and blood in five paired samples, and in tissue only in two samples (71% concordance). In 9 subjects with paired tissue and blood samples, an EGFR driver mutation was identified in plasma and tissue (n=5), plasma only (n=1) or tissue only (n=3). Overall, the investigators concluded that in this limited cohort, ctDNA is an option when tissue is unavailable.

Another study of ctDNA testing for 70 genes and NSCLC was performed by Thompson, et al. (2016). A total of 112 plasma samples were obtained from 102 prospectively enrolled patients with advanced NSCLC. Matched tissues samples, when available, were also evaluated. The investigators found 275 alterations in 45 genes, and at least one alteration in the ctDNA for 86 of 102 patients (84%), with EGFR variants being most common. This testing detected 50 driver and 12 resistance mutations, and mutations in 22 additional genes for which experimental therapies, including clinical trials, are available. Tissue sequencing was only successful for 50 patients (49%). Overall concordance for all variants covered and detected by both platforms was 60%. Actionable EGFR mutations were detected in 24 tissue and 19 ctDNA samples, yielding concordance of 79%.

InVisionFirst Lung

InVisionFirst is a liquid biopsy test that analyzes the presence of relevant genetic variants in the *ALK*, *BRAF*, *EGFR*, *ERBB2*, *KRAS*, *MET*, *ROS1* and *STK11* and 26 other genes in patients with non-small cell lung cancer. Plagnol et al. (2018) reported on the analytical validation of the TAM-Seq technology utilized in InVisionFirst Lung. At least two 10ml tubes of blood were collected from each donor into Streck Cell Free DNA Blood Collection tubes (BCT) and EDTA tubes. Ninety-five samples from healthy donors were analyzed for gene fusions, and no genetic variants were found. One hundred and nine samples from healthy donors were analyzed for SNVs, indels and amplications, and no copy number variants were found. Three splice site variants were found. Digital PCR (dPCR) was performed on these three and a *TP53* mutation was confirmed, but not the other two. A further 92 samples from healthy donors and 242 samples from untreated NSCLC patients were tested, and these three variants were not seen. In the affected group, twenty NSCLC patients were tested by both InVisionFirst and dPCR at two separate labs, who were blinded to each other's results. In this cohort, 40% of patients had a genetic variant. dPCR detected 19 of 20 expected changes. InVisionFirst identified a mutation in one sample not seen with dPCR, and the sample had a very low cfDNA fraction. It can't be determined if this was a true positive undetectable by dPCR or a false positive. In addition, contrived samples using various seeded cell lines and reference material were used to simulate a wide array of copy number and other genetic variations were tested in the same way. Overall, in the donor samples and contrived materials, the concordance rate between InVisionFirst and dPCR was high. InVisionFirst demonstrated a >99% sensitivity for SNVs and >92% for indels.

The NCCN NSCLC Guidelines Panel for NSCLC have added a section on Plasma Cell-Free/Circulating Tumor DNA Testing that states that cell-free/circulating tumor DNA testing should not be used in lieu of a tissue diagnosis as the analytical standards have not been established. However, NCCN also suggests that the use of cell-free/circulating tumor DNA testing can be considered in specific clinical circumstances, most notably: if a patient is medically unfit for invasive tissue sampling; or in the initial diagnostic setting, if following pathologic confirmation of a NSCLC diagnosis there is insufficient material for molecular analysis, cell-free/circulating tumor DNA should be used only if follow up tissue based analysis is planned for all patients in which an oncogenic driver is not identified.

In a 2017 study, Cohen et al. developed a noninvasive test for detection of pancreatic ductal adenocarcinoma. They combined blood tests for KRAS gene mutations with protein biomarkers as a testing method. They tested this assay on a cohort of 221 patients with resectable pancreatic ductal adenocarcinomas and 182 control patients without known cancer. In the plasma samples of 66 patients (30%), KRAS mutations were detected and every mutation found in the plasma was also detected in the primary tumor (100% concordance). This combination of tests increased the sensitivity to 64%. Only one of the control samples was positive for any of the DNA or protein biomarkers (99.5% specificity). The researchers concluded that this approach may prove useful for early cancer detection.

Sun et al. (2018) published a study examining liquid biopsies in colorectal cancer (CRC). The researchers analyzed blood from 140 CRC patients with matched tumor samples. Both the circulating tumor cells (CTC) and tumor DNA (ctDNA) were extracted before surgery and treatment. The samples were quantified and tested for mutations in KRAS, NRAS and BRAF. Within this sample cohort, there was good agreement between the CTC and the ctDNA (97% concordance). The researchers also determined that patients who were refractory to specific medications showed molecular profile changes and were positive for KRAS, NRAS or BRAF. This was noteworthy as the changes were detected in the circulating tumor cells first. The study concluded that using CTC and ctDNA for monitoring CRC patients molecular profile changes to treatment may be useful.

A study from Dieffenbacher et al. (2018) evaluated tumor tissue and liquid biopsies in metastatic clear cell renal cell cancer patients in the MORE-TRIAL. Samples were performed at baseline and first and second progression under treatment. The study stated that this relatively new technique may help to avoid the necessity for invasive biopsies in the future and a further aim of MORE is to study the reliability and relevance of ct-DNA in RCC patients.

Another renal cell carcinoma study by Yamamoto et al. (2019) evaluated circulating tumor DNA for clinical utility. Fifty-three patients histologically diagnosed with clear cell RCC were enrolled and sequencing was performed on plasma cell-free DNA (cfDNA) and tumor DNA. A total of 38 mutations across 16 (30%) patients were identified from cfDNA, including mutations in TP53 (n=6) and VHL (n=5), and median mutant allele frequency of ctDNA was 10%. The researchers concluded that this study shows the clinical utility of ctDNA for prognosis and disease monitoring in RCC.

Clinical Practice Guidelines

American Society of Clinical Oncology (ASCO)

Merker et al. (2018) published a joint review from the American Society of Clinical Oncology (ASCO) and the College of American Pathologists (CAP) to assess the clinical use of circulating tumor DNA (ctDNA). The researchers performed a literature review and identified 1,339 references. Of these references, 390, plus an additional 31 supplied by the researchers, were reviewed. The literature review included 77 references. The literature review stated that while some ctDNA tests have demonstrated clinical validity and utility with specific advanced stage cancer, overall, there is insufficient evidence of clinical validity and utility for the majority of these assays in this stage of cancer. The researchers also noted that there is no evidence of clinical utility and little evidence of clinical validity of ctDNA tests in early stage cancer, treatment monitoring, or residual disease detection. Likewise, no evidence of clinical validity and utility was demonstrated in the literature review for the use of ctDNA in cancer screening.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Laboratories that perform genetic tests are regulated under the Clinical Laboratory Improvement Amendments (CLIA) Act of 1988. More information is available at:

<https://www.fda.gov/medicaldevices/deviceregulationandguidance/ivdregulatoryassistance/ucm124105.htm>.

(Accessed April 5, 2019)

References

- Adalsteinsson VA, Ha G, Freeman SS, et al. Scalable whole-exome sequencing of cell-free DNA reveals high concordance with metastatic tumors. *Nature Communications*. 2017; 8:1324.
- Alexander EK, Kennedy GC, Baloch ZW, et al. Preoperative diagnosis of benign thyroid nodules with indeterminate cytology. *N Engl J Med*. 2012;367(8):705–715.
- Alexander EK, Schorr M, Klopper J, et al. Multicenter clinical experience with the Afirma gene expression classifier. *J Clin Endocrinol Metab*. 2014;99(1):119–125.
- Ali SM, Sanford EM, Klemptner SJ, et al. Prospective comprehensive genomic profiling of advanced gastric carcinoma cases reveals frequent clinically relevant genomic alterations and new routes for targeted therapies. *Oncologist*. 2015;20(5):499-507.
- Arber DA, Borowitz MJ, Cessna M, et al. Initial diagnostic workup of acute leukemia: Guideline from the College of American Pathologists and the American Society of Hematology. *Archives of Pathology & Laboratory Medicine*: October 2017, Vol. 141, No. 10, pp. 1342-1393.
- Ardakani MN, Thomas C, Robinson C, et al. Detection of copy number variations in melanocytic lesions utilising array based comparative genomic hybridisation. *Pathology*. 2017 Apr;49(3):285-291.
- Avet-Loiseau H, Corre L, Lauwers-Cances V, et al. Evaluation of minimal residual disease (MRD) by next generation sequencing (NGS) is highly predictive of PFS in the IFM/DFCI 2009 trial. *Blood*. 2015;126:191.
- Beaudenon-Huibregtse S, Alexander EK, Guttler RB, et al. Centralized molecular testing for oncogenic gene mutations complements the local cytopathologic diagnosis of thyroid nodules. *Thyroid*. 2014;24(10):1479–1487.
- Bekelman JE, Rumble RB, Freedland SJ. Clinically localized prostate cancer: ASCO Clinical Practice Guideline endorsement of an AUA/ASTRO/SUO Guideline Summary. *J Oncol Pract*. 2018 Oct;14(10):618-624.
- Beltran H, Yelensky R, Frampton GM, et al. Targeted next-generation sequencing of advanced prostate cancer identifies potential therapeutic targets and disease heterogeneity. *Eur Urol*. 2013 May;63(5):920-6.
- Berger AC, Davidson RS, Poitras JK, et al. Clinical impact of a 31-gene expression profile test for cutaneous melanoma in 156 prospectively and consecutively tested patients. *Curr Med Res Opin*. 2016 Sep;32(9):1599-604.
- Bichakjian CK, Halpern AC, Johnson TM, et al. Guidelines of care for the management of primary cutaneous melanoma. *American Academy of Dermatology*. *J Am Acad Dermatol*. 2011 Nov;65(5):1032-47.
- Boulwood J, Perry J, Zaman R., et al. High-density single nucleotide polymorphism array analysis and ASXL1 gene mutation screening in chronic myeloid leukemia during disease progression. *Leukemia*. 2010;24(6):1139–1145.
- Brand TC, Zhang N, Crager MR, et al. Patient-specific meta-analysis of 2 clinical validation studies to predict pathologic outcomes in prostate cancer using the 17-gene genomic prostate score. *Urology*. 2016 Mar;89:69-75.
- Brauner E, Holmes BJ, Krane JF, et al. Performance of the Afirma gene expression classifier in Hurthle cell thyroid nodules differs from other indeterminate thyroid nodules. *Thyroid*. 2015;25(7):789–796.
- Carter HB, Albertsen PC, Barry MJ, et al. American Urological Association (AUA). Early detection of prostate cancer. Published 2013; reviewed and validity confirmed 2018.
- Chantrill LA, Nagrial AM, Watson C, et al.; Australian Pancreatic Cancer Genome Initiative (APGI) and the Individualized Molecular Pancreatic Cancer Therapy (IMPACT) Trial Management Committee of the Australasian Gastrointestinal Trials Group (AGITG). Precision medicine for advanced pancreas cancer: The Individualized Molecular Pancreatic Cancer Therapy (IMPACT) Trial. *Clin Cancer Res*. 2015 May 1;21(9):2029-37.
- Chen RC, Rumble RB, Loblaw DA, et al. Active surveillance for the management of localized prostate cancer (Cancer Care Ontario guideline): American Society of Clinical Oncology Clinical practice guideline endorsement. *J Clin Oncol* 2016 34:2182-2190.
- Chmielecki J, Hutchinson KE, Frampton GM, et al. Comprehensive genomic profiling of pancreatic acinar cell carcinomas identifies recurrent RAF fusions and frequent inactivation of DNA repair genes. *Cancer Discov*. 2014 Dec;4(12):1398-405.
- Chua MLK, Lo W, Pintilie M, et al. Prostate cancer “nimbus”: genomic instability and schlaf1 dysregulation underpin aggression of intraductal and cribriform subpathologies. *Eur Urol*. 2017 May 13. [Epub ahead of print].

Chung CH, Guthrie VB, Masica DL, et al. Genomic alterations in head and neck squamous cell carcinoma determined by cancer gene-targeted sequencing. *Ann Oncol*. 2015 Feb 23. pii: mdv109.

Cohen JD, Javed AA, Thoburn C, et al. Combined circulating tumor DNA and protein biomarker-based liquid biopsy for the earlier detection of pancreatic cancers. *Proc Natl Acad Sci U S A*. 2017 Sep 19;114(38):10202-10207.

Cooley LD, Lebo M, Li MM, et al. Working Group of the American College of Medical Genetics and Genomics (ACMG) Laboratory Quality Assurance Committee. American College of Medical Genetics and Genomics technical standards and guidelines: microarray analysis for chromosome abnormalities in neoplastic disorders. (2013) *Genet Med* ;15:484–494.

Crawford ED, Scholz MC, Kar AJ, et al. Cell cycle progression score and treatment decisions in prostate cancer: results from an ongoing registry. *Curr Med Res Opin*. 2014 Jun;30(6):1025-31.

Dagogo-Jack I, Rooney M, Nagy RJ, et al. Molecular analysis of plasma from patients with ROS1-Positive NSCLC. *J Thorac Oncol*. 2019 Jan 18. pii: S1556-0864(19)30027-9.

Dalerba P, Sahoo D, Paik S, et al. CDX2 as a prognostic biomarker in stage II and stage III colon cancer. *N Engl J Med* 2016;374:211-22.

Deaver KE, Haugen BR, Pozdeyev N, Marshall CB. Outcomes of Bethesda categories III and IV thyroid nodules over 5 years and performance of the Afirma gene expression classifier: a single-institution study. *Clin Endocrinol (Oxf)*. 2018 May 23.

Detterbeck FC, Lewis SZ, Diekemper R, et al. Executive Summary: Diagnosis and management of lung cancer, 3rd ED: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2013 May;143(5 Suppl):7S-37S.

Dieffenbacher S, Zschäbitz S, Hofer L, et al. Prospective single center trial of next-generation sequencing analysis in metastatic renal cell cancer: the MORE-TRIAL. *Future Sci OA*. 2018;4(5):FSO299. Published 2018 Mar 14.

Drilon A, Wang L, Arcila ME, et al. Hybrid capture–based next-generation sequencing identifies actionable genomic alterations in lung adenocarcinomas otherwise negative for such alterations by other genomic testing approaches. *Clin Cancer Res*. 2015 Aug 15;21(16):3631-9.

Duick DS, Klopper J, Diggins, J et al. The impact of benign gene expression classifier test results on the endocrinologist/patient decision to operate on patients with thyroid nodules with indeterminate fine-needle aspiration cytopathology. *Thyroid*. 2012 Oct; 22(10): 996–1001.

Dummer R, Hauschild A, Lindenblatt N, et al. ESMO Guidelines Committee. Cutaneous melanoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2015 Sep;26 Suppl 5:v126-32.

Evans AG, Ahmad A, Burack WR, Iqbal MA. “Combined comparative genomic hybridization and single-nucleotide polymorphism array detects cryptic chromosomal lesions in both myelodysplastic syndromes and cytopenias of undetermined significance.” *Modern pathology: an official journal of the United States and Canadian Academy of Pathology, Inc*. 2016 Oct 0; 29(10):1183-99.

Ferris LK, Gerami P, Skelsey MK, et al. Real-world performance and utility of a noninvasive gene expression assay to evaluate melanoma risk in pigmented lesions. *Melanoma Res*. 2018 Oct;28(5):478-482.

Ferris LK, Jansen B, Ho J, et al. Utility of a noninvasive 2-gene molecular assay for cutaneous melanoma and effect on the decision to biopsy. *JAMA Dermatol*. 2017;153(7):675–680.

Fizazi K, Greco FA, Pavlidis N, et al. Cancers of unknown primary site: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* (2015) 26 (suppl 5): v133-v138.

Food and Drug Administration (FDA) FDA authorizes first next generation sequencing-based test to detect very low levels of remaining cancer cells in patients with acute lymphoblastic leukemia or multiple myeloma. <https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm622004.htm>. Accessed June 20, 2019.

Frampton GM, Fichtenholtz A, Otto GA, et al. Development and validation of a clinical cancer genomic profiling test based on massively parallel DNA sequencing. *Nat Biotechnol*. 2013 Nov;31(11):1023-31.

Gatalica Z, Millis SZ, Vranic S, et al. Comprehensive tumor profiling identifies numerous biomarkers of drug response in cancers of unknown primary site: analysis of 1806 cases. *Oncotarget*. 2014 Dec 15;5(23):12440-7.

Gharib H, Papini E, Garber JR, et al. American Association of Clinical Endocrinologists, American College Of Endocrinology, And Associazione Medici Endocrinologi Medical Guidelines For Clinical Practice For The Diagnosis And Management Of Thyroid Nodules – 2016 Update. *Endocrine Practice* May 2016 Vol 22 (Suppl 1).

Glass AG, Leo MC, Haddad Z., et al. Validation of a genomic classifier for predicting post-prostatectomy recurrence in a community-based health care setting. *J Urol*. 2016 Jun;195(6):1748-53.

Hagenkord JM, Monzon FA, Kash SF, et al. Array-based karyotyping for prognostic assessment in chronic lymphocytic leukemia: Performance comparison of Affymetrix 10K2.0, 250K Nsp, and SNP6.0 Arrays. *The Journal of Molecular Diagnostics : JMD*. 2010;12(2):184-196.

Han M, Liew CT, Zhang HW, et al. Novel, blood-based five-gene panel biomarker set for the detection of colorectal cancer. *Clin Cancer Res*. 2008;14:455–60.

Harrell RM, Bimston DN. Surgical utility of Afirma: effects of high cancer prevalence and oncocyctic cell types in patients with indeterminate thyroid cytology. *Endocr Pract*. 2014;20(4):364–369

Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid*. 2016;26(1):1-133.

He N, Song L, Kang Q, et al. The pathological features of colorectal cancer determine the detection performance on blood ctDNA. *Technol Cancer Res Treat*. 2018;17:1533033818791794.

Heichman K. Blood-based testing for colorectal cancer screening. *Molecular Diagnosis & Therapy*; Volume 18, Issue 2; p:127-135.

Hirshfield KM, Tolkunov D, Zhong H et al. Clinical actionability of comprehensive genomic profiling for management of rare or refractory cancers. *Oncologist*. 2016 Aug 26. pii: theoncologist. 2016-0049. [Epub ahead of print]

Hutchinson KE, Lipson D, Stephens PJ, et al. BRAF fusions define a distinct molecular subset of melanomas with potential sensitivity to MEK inhibition. *Clin Cancer Res*. 2013 Dec 15;19(24):6696-702.

Johnson DB, Dahlman KH, Knol J, et al. Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel. *Oncologist*. 2014 Jun;19(6):616-22.

Jongen-Lavrencic M, Grob T, Hanekamp D, et al. Molecular minimal residual disease in acute myeloid leukemia. *N Engl J Med* 378:1189-1199, 2018.

Kamps R, Brandão RD, van den Bosch BJ, et al. Next-generation sequencing in oncology: Genetic diagnosis, risk prediction and cancer classification. Cho WC, ed. *International Journal of Molecular Sciences*. 2017;18(2):308.

Kato S, Schwaederle M, Daniels GA, et al. Cyclin-dependent kinase pathway aberrations in diverse malignancies: clinical and molecular characteristics. *Cell Cycle*. 2015 Apr 18;14(8):1252-9.

Kalemkerian GP, Narula N, Kennedy EB et al. Molecular testing guideline for the selection of lung cancer patients for treatment with targeted tyrosine kinase inhibitors: American Society of Clinical Oncology Endorsement Summary of the College of American Pathologists/International Association for the Study of Lung Cancer/Association for Molecular Pathology Clinical Practice Guideline Update. *J Oncol Pract*. 2018 Mar 28;JOP1800035

Kim K, Zakharkin SO, Allison DB. Expectations, validity, and reality in gene expression profiling. *Journal of clinical epidemiology*. 2010;63(9):950-959.

Kim ST, Banks KC, Lee SH, et al. [Prospective feasibility study for using cell-free circulating tumor DNA-guided therapy in refractory metastatic solid cancers: an interim analysis](#). *JCO Precision Oncology* 2017;1, 1-15.

Klein EA, Santiago-Jiménez M, Yousefi K, et al. Molecular analysis of low-grade prostate cancer using a genomic classifier of metastatic potential. *J Urol*. 2016 Jan;197(1):122-128.

Klufas MA, Richter E, Itty S, et al. Comparison of gene expression profiling and chromosome 3 analysis by fluorescent in situ hybridization and multiplex ligation probe amplification in fine-needle aspiration biopsy specimens of uveal melanoma. *Ocul Oncol Pathol*. 2017 Dec;4(1):16-20.

Koh KN, Lee JO, Seo EJ, et al. Clinical significance of previously cryptic copy number alterations and loss of heterozygosity in pediatric acute myeloid leukemia and Myelodysplastic Syndrome determined using combined array comparative genomic hybridization plus single-nucleotide polymorphism microarray analyses. *Journal of Korean Medical Science*. 2014;29(7):926-933.

Kolquist KA, Schultz RA, Furrow A, et al. Microarray-based comparative genomic hybridization of cancer targets reveals novel, recurrent genetic aberrations in the myelodysplastic syndromes. *Cancer Genet* 2011;204(11):603-628.

Kutzner H, Metzler G, Argenyi Z, et al. Histological and genetic evidence for a variant of superficial spreading melanoma composed predominantly of large nests. *Mod Pathol*. 2012;25:838–45.

Labourier E, Shifrin A, Busseniers AE, et al. Molecular testing for miRNA, mRNA, and DNA on fine-needle aspiration improves the preoperative diagnosis of thyroid nodules with indeterminate Cytology. *J Clin Endocrinol Metab* 2015; 100: 2743–2750.

Ladetto M, Bruggemann M, Monitillo L, et al. Next-generation sequencing and real-time quantitative PCR for minimal residual disease detection in B-cell disorders. *Leukemia*. 2013;28:1299-1307.

Lam VK, Tran HT, Banks KC, et al. Targeted tissue and cell-free tumor DNA sequencing of advanced lung squamous-cell carcinoma reveals clinically significant prevalence of actionable alterations. *Clin Lung Cancer*. 2019 Jan;20(1):30-36.e3.

Lastra RR, Pramick MR, Crammer CJ, et al. Implications of a suspicious Afirma test result in thyroid fine-needle aspiration cytology: an institutional experience. *Cancer Cytopathol*. 2014;122(10):737–744.

Laurie CC, Laurie CA, Smoley SA, et al. Acquired chromosomal anomalies in chronic lymphocytic leukemia (CLL) patients compared to >50,000 quasi-normal Subjects. *Cancer Genetics*. 2014;207(0):19-30.

Lowery MA, Wong W, Jordan EJ, et al. Prospective evaluation of germline alterations in patients with exocrine pancreatic neoplasms. *J Natl Cancer Inst*. 2018 Oct 1;110(10):1067-1074.

Marrone M, Potosky AL, Penson D, Freedman AN. A 22 gene-expression assay, Decipher® (GenomeDx Biosciences) to predict five-year risk of metastatic prostate cancer in men treated with radical prostatectomy. *PLoS Currents*. 2015;7.

Marshall KW, Mohr S, El Khettabi F, et al. A blood-based biomarker panel for stratifying current risk for colorectal cancer. *Int J Cancer*. 2010;126:1177–86.

Marti JL, Avadhani V, Donatelli LA, et al. Wide inter-institutional variation in performance of a molecular classifier for indeterminate thyroid nodules. *Ann Surg Oncol*. 2015;22(12):3996–400.

McCoach CE, Blakely CM, Banks KC, et al. Clinical utility of cell-free DNA for the detection of *ALK* fusions and genomic mechanisms of ALK inhibitor resistance in non-small cell lung cancer. *Clin Cancer Res*. 2018 Jun 15;24(12):2758-2770.

McKiernan J, Donovan MJ, O’Neill V, et al. A novel urine exosome gene expression assay to predict high-grade prostate cancer at initial biopsy. *JAMA Oncol*. 2016 Jul 1;2(7):882-9.

McKiernan J, Donovan M, Margolis E, et al. A prospective adaptive utility trial to validate performance of a novel urine exosome gene expression assay to predict high-grade prostate cancer in patients with prostate-specific Antigen 2–10 ng/ml at initial biopsy. *European Urology* 2018; 74(6):731-738.

Meleth S, Whitehead N, Swinson T, et al. Technology assessment on genetic testing or molecular pathology testing of cancers with unknown primary site to determine origin. *Technology Assessment Report*. Rockville, MA: Agency for Healthcare Research and Quality. February 2013.

Merker JD, Oxnard GR, Compton C, et al. Circulating tumor DNA analysis in patients with cancer: American Society of Clinical Oncology and College of American Pathologists Joint Review. *J Clin Oncol*. 2018 Jun 1;36(16):1631-1641.

Millis SZ, Bryant D, Basu G, et al. Molecular profiling of infiltrating urothelial carcinoma of bladder and nonbladder origin. *Clin Genitourin Cancer*. 2015 Feb;13(1):e37-49.

Minca EC, Tubbs RR, Portier BP, et al. Genomic microarray analysis on formalin-fixed paraffin-embedded material for uveal melanoma prognostication. *Cancer Genet*. 2014;207:306–315.

Miura JT, Johnston FM, Thomas J, et al. Molecular profiling in gastric cancer: examining potential targets for chemotherapy. *J Surg Oncol*. 2014 Sep;110(3):302-6.

Moschini M, Spahn M, Mattei A, et al. Incorporation of tissue-based genomic biomarkers into localized prostate cancer clinics. *BMC Med*. 2016; 14: 67.

Na R, Wu Y, Ding Q, et al. Clinically available RNA profiling tests of prostate tumors: utility and comparison. *Asian J Androl*. 2016 Jul-Aug; 18(4): 575–579.

National Comprehensive Cancer Network (NCCN). *Clinical Practice Guidelines in Oncology. Bladder cancer. Version 2.2019a.*

National Comprehensive Cancer Network (NCCN). *Clinical Practice Guidelines in Oncology. Colon cancer. Version 1.2019c.*

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Melanoma. Version 2.2019d.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Multiple myeloma. Version 2.2019.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Non-small cell lung cancer. Version 3.2019e.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Occult primary (cancer of unknown primary [CUP]). V2.2019f.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Prostate cancer. Version 1.2019g.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Rectal cancer. Version 1.2019h.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Thyroid Carcinoma Version 1.2019i.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Uveal melanoma. Version 1.2018b.

National Institute for Health and Clinical Excellence (NICE). Metastatic malignant disease of unknown primary origin in adults: diagnosis and management. Clinical Guideline CG04. July 2010.

Nikiforova MN, Mercurio S, Wald AI, et al. Analytical performance of the ThyroSeq v3 genomic classifier for cancer diagnosis in thyroid nodules. *Cancer*. 2018 Apr 15;124(8):1682-1690.

Nishino M and Nikiforova M. Update on molecular testing for cytologically indeterminate thyroid nodules. *Archives of Pathology & Laboratory Medicine*: April 2018, Vol. 142, No. 4, pp. 446-457.

Oderda M, Cozzi G, Daniele L, et al. Cell-cycle progression-score might improve the current risk assessment in newly diagnosed prostate cancer patients. *Urology*. 2016 Nov 25.

Onken MD, Worley LA, Char DH, et al. Collaborative Ocular Oncology Group report number 1: prospective validation of a multi-gene prognostic assay in uveal melanoma. *Ophthalmology*. 2012 Aug;119(8):1596-603.

Oxnard GR, Thress KS, Alden RS, et al. Association between plasma genotyping and outcomes of treatment with osimertinib (AZD9291) in advanced non-small-cell lung cancer. *J Clin Oncol*. 2016 Oct 1;34(28):3375-82.

Pagan M, Kloos RT, Lin C-F, et al. The diagnostic application of RNA sequencing in patients with thyroid cancer: an analysis of 851 variants and 133 fusions in 524 genes. *BMC Bioinformatics*. 2016;17(Suppl 1):6.

Palmetto GBA. Local Coverage Article for MolDx: AFIRMA™ Assay by Veracyte Update (Article ID Number A54356). January 5, 2018.

Partyka KL, Randolph ML, Lawrence KA, et al. Utilization of direct smears of thyroid fine-needle aspirates for ancillary molecular testing: A comparison of two proprietary testing platforms. *Diagn Cytopathol*. 2018 Apr;46(4):320-325.

PDQ Adult Treatment Editorial Board. Intraocular (Uveal) Melanoma Treatment (PDQ®): Health Professional Version. 2018 Feb 8.

Peterson JF, Aggarwal N, Smith CA, et al. Integration of microarray analysis into the clinical diagnosis of hematological malignancies: How much can we improve cytogenetic testing? *Oncotarget*. 2015;6(22):18845-18862.

Peterson JF, Van Dyke DL, Hoppman NL, et al. The utilization of chromosomal microarray technologies chromosomal microarray technologies for hematologic neoplasms: Hematologic Neoplasms: An ACLPS critical review. *Am J Clin Pathol*. 2018 Oct 1;150(5):375-384.

Plagnol V, Woodhouse S, Howarth K, et al. Analytical validation of a next generation sequencing liquid biopsy assay for high sensitivity broad molecular profiling. *PLoS One*. 2018;13(3):e0193802. Published 2018 Mar 15.

Plasseraud KM, Cook RW, Tsai T, et al. Clinical performance and management outcomes with the DecisionDx-UM gene expression profile test in a prospective multicenter study. *J Oncol*. 2016;2016:5325762.

Port M, Böttcher M, Thol F, et al. Prognostic significance of FLT3 internal tandem duplication, nucleophosmin 1, and CEBPA gene mutations for acute myeloid leukemia patients with normal karyotype and younger than 60 years: a systematic review and meta-analysis. *Ann Hematol*. 2014 Aug;93(8):1279-86.

Puiggros A, Puigdecenet E, Salido M, et al. Genomic arrays in chronic lymphocytic leukemia routine clinical practice: are we ready to substitute conventional cytogenetics and fluorescence in situ hybridization techniques? *Leuk Lymphoma*. 2013;54:986-995.

Raskin L, Ludgate M, Iyer RK et al. Copy number variations and clinical outcome in Atypical Spitz Tumors. *The American Journal of Surgical Pathology*: 2011 35(2) : p 243–252.

Riediger AL, Dietz S, Schirmer U, et al. Mutation analysis of circulating plasma DNA to determine response to EGFR tyrosine kinase inhibitor therapy of lung adenocarcinoma patients. *Sci Rep*. 2016 Sep 19;6:33505.

Rodriguez-Rodriguez L, Hirshfield KM, Rojas V, et al. Use of comprehensive genomic profiling to direct point-of-care management of patients with gynecologic cancers. *Gynecol Oncol*. 2016;141(1):2-9.

Ross JS, Ali SM, Wang K, et al. Comprehensive genomic profiling of epithelial ovarian cancer by next generation sequencing-based diagnostic assay reveals new routes to targeted therapies. *Gynecol Oncol*. 2013 Sep;130(3):554-9.

Ross JS, Wang K, Al-Rohil RN, et al. Advanced urothelial carcinoma: next-generation sequencing reveals diverse genomic alterations and targets of therapy. *Mod Pathol*. 2014b Feb;27(2):271-80.

Ross JS, Wang K, Gay L et al. Comprehensive genomic profiling of carcinoma of unknown primary site: new routes to targeted therapies. *JAMA Oncol*. 2015;1(1):40-49.

Ross JS, Wang K, Rand JV, et al. Next-generation sequencing of adrenocortical carcinoma reveals new routes to targeted therapies. *J Clin Pathol*. 2014a Nov;67(11):968-73.

Roy S, Pradhan D, Ernst WL, et al. Next-generation sequencing-based molecular characterization of primary urinary bladder adenocarcinoma. *Mod Pathol*. 2017 May 26. [Epub ahead of print].

Sacher AG, Paweletz C, Dahlberg SE, et al. Prospective validation of rapid plasma genotyping for the detection of EGFR and KRAS mutations in advanced lung cancer. *JAMA Oncol*. 2016 Aug 1;2(8):1014-22.

Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically localized prostate cancer: AUA/ASTRO/SUO Guideline.Part I: risk stratification, shared decision making, and care options.Care Options. *J Urol*. 2018 Mar;199(3):683-690.

Santhanam P, Khthir R, Gress T, et al. Gene expression classifier for the diagnosis of indeterminate thyroid nodules: a meta-analysis. *Med Oncol*. 2016 Feb;33(2):14.

Shore N, Concepcion R, Saltzstein D, et al. Clinical utility of a biopsy-based cell cycle gene expression assay in localized prostate cancer. *Curr Med Res Opin*. 2014 Apr;30(4):547-53.

Sipos JA, Blevins TC, Shea HC, et al. Long-term nonoperative rate of thyroid nodules with benign results on the afirma gene expression classifier. *Endocr Pract*. 2016 Jun;22(6):666-72.

Sohal DPS, Kennedy EB, Khorana A, et al. Metastatic pancreatic cancer: ASCO Clinical Practice Guideline Update. *Journal of Clinical Oncology*, 2018 36:24,2545-2556.

Song Q, Chu Y, Yao Y, et al. Identify latent chromosomal aberrations relevant to myelodysplastic syndromes. *Scientific Reports*. 2017b;7:10354.

Song Q, Peng M, Chu Y, Huang S. Techniques for detecting chromosomal aberrations in myelodysplastic syndromes. *Oncotarget*. 2017a;8(37):62716-62729.

Su KY, Tseng JS, Liao KM, et al. Mutational monitoring of EGFR T790M in cfDNA for clinical outcome prediction in EGFR-mutant lung adenocarcinoma. *PLoS One*. 2018 Nov 16;13(11):e0207001.

Sun Q, Liu Y, Liu B, Liu Y. Use of liquid biopsy in monitoring colorectal cancer progression shows strong clinical correlation. *Am J Med Sci*. 2018 Mar;355(3):220-227.

Thiel A, Beier M, Ingenhag D, et al. Comprehensive array CGH of normal karyotype myelodysplastic syndromes reveals hidden recurrent and individual genomic copy number alterations with prognostic relevance. *Leukemia* 2011; 25: 387–399.

Thompson JC, Yee SS, Troxel AB, et al. Detection of therapeutically targetable driver and resistance mutations in lung cancer patients by next-generation sequencing of cell-free circulating tumor DNA. *Clin Cancer Res*. 2016;22(23):5772-5782.

Tiu RV, Gondek LP, O’Keefe CL, et al. Prognostic impact of SNP array karyotyping in myelodysplastic syndromes and related myeloid malignancies. *Blood*. 2011;117(17):4552-4560.

Trikalinos TA, Terasawa T, Raman G, et al. A systematic review of loss-of-heterozygosity based topographic genotyping with PathfinderTG. *Technology Assessment Report GEND0308*. Prepared by the Tufts Evidence-based Practice Center for the Agency for Healthcare Research and Quality AHRQ under Contract No HHSA 290 10055 I AHRQ March. 2010.

U.S. National Library of Medicine. Do all gene mutations affect health and development? Genetics Home Reference. October 2017B.

U.S. National Library of Medicine. What are whole exome sequencing and whole genome sequencing? Genetics Home Reference. October 2017A.

Varadhachary GR, Raber MN. Cancer of unknown primary site. *N Engl J Med* 2014;371:757-65.

Venook AP, Niedzwiecki D, Lopatin M, et al. Biologic determinants of tumor recurrence in stage II colon cancer: validation study of the 12-gene recurrence score in cancer and leukemia group B (CALGB) 9581. *J Clin Oncol*. 2013;31(14):1775-1781.

Vignot S, Lefebvre C, Frampton GM, et al. Comparative analysis of primary tumour and matched metastases in colorectal cancer patients: Evaluation of concordance between genomic and transcriptional profiles. *Eur J Cancer*. 2015 May;51(7):791-9.

Villaflor V, Won B, Nagy R, et al. Biopsy-free circulating tumor DNA assay identifies actionable mutations in lung cancer. *Oncotarget*. 2016;7(41):66880-66891.

Wang K, McDermott JD, Schrock AB, et al. Comprehensive genomic profiling of salivary mucoepidermoid carcinomas reveals frequent BAP1, PIK3CA, and other actionable genomic alterations. *Ann Oncol*. 2017 Apr 1;28(4):748-753.

Weinhold N, Heuck C, Rosenthal A, et al. The clinical value of molecular subtyping multiple myeloma using gene expression profiling. *Leukemia*. 2016 February; 30(2): 423–430.

West AC, Tang K, Tye H, et al. Identification of a TLR2-regulated gene signature associated with tumor cell growth in gastric cancer. *Oncogene*. 2017 May 8. [Epub ahead of print].

Wheler J, Yelensky R, Falchook G, et al. Next generation sequencing of exceptional responders with BRAF-mutant melanoma: implications for sensitivity and resistance. *BMC Cancer*. 2015 Feb 18;15:61.

Wiesner T, Kutzner H, Cerroni L, et al. Genomic aberrations in spitzoid tumours and their implications for diagnosis, prognosis and therapy. *Pathology*. 2016;48(2):113-131.

Wolmark N, Mamounas EP, Baehner FL et al. Prognostic impact of the combination of recurrence score and quantitative estrogen receptor expression (ESR1) on predicting late distant recurrence risk in estrogen receptor–positive breast cancer after 5 years of tamoxifen: Results from NRG Oncology/National Surgical Adjuvant Breast and Bowel Project B-28 and B-14 *Journal of Clinical Oncology* 34, no. 20 (July 2016) 2350-2358.

Wong W, Lowery MA, Berger MF, et al. Ampullary cancer: evaluation of somatic and germline genetic alterations and association with clinical outcomes. *Cancer*. 2019 Jan 8.

Wood B, Wu D, Crossley B, et al. Measurable residual disease detection by high-throughput sequencing improves risk stratification for pediatric B-ALL. *Blood*. 2018 Mar 22;131(12):1350-1359.

Wylie D, Beaudenon-Huibregtse S, Haynes B, et al. Molecular classification of thyroid lesions by combined testing for miRNA gene expression and somatic gene alterations. *The Journal of Pathology: Clinical Research*. 2016;2(2):93-103.

Yamanaka T, Oki E, Yamazaki K, et al. 12-gene recurrence score assay stratifies the recurrence risk in stage II/III colon cancer with surgery alone: the SUNRISE study. *J Clin Oncol*. 2016 Aug 20;34(24):2906-13.

Yamamoto Y, Uemura M, Fujita M, et al. Clinical significance of the mutational landscape and fragmentation of circulating tumor DNA in renal cell carcinoma. *Cancer Sci*. 2019;110(2):617-628.

Yan S, Liu Z, Yu S, Bao Y. Diagnostic value of methylated septin9 for colorectal cancer screening: a meta-analysis. *Med Sci Monit*. 2016;22:3409–3418.

Yang C, Zou K, Yuan Z, et al. Prognostic value of circulating tumor cells detected with the CellSearch System in patients with gastric cancer: evidence from a meta-analysis. *OncoTargets and therapy*. 2018;11:1013-1023.

Yang M, Topaloglu U, Petty WJ, et al. Circulating mutational portrait of cancer: manifestation of aggressive clonal events in both early and late stages. *J Hematol Oncol*. 2017 May 4;10(1):100.

Yang SE, Sullivan PS, Zhang J et al. Has Afirma gene expression classifier testing refined the indeterminate thyroid category in cytology? *Cancer Cytopathol*. 2016 Feb;124(2):100-9.

Yothers G, O’Connell MJ, Lee M, et al. Validation of the 12-gene colon cancer recurrence score in NSABP C-07 as a predictor of recurrence in patients with stage II and III colon cancer treated with fluorouracil and leucovorin (FU/LV) and FU/LV plus oxaliplatin. *J Clin Oncol*. 2013 Dec 20;31(36):4512-9.

Zager JS, Gastman BR, Leachman S, et al. Performance of a prognostic 31-gene expression profile in an independent cohort of 523 cutaneous melanoma patients. BMC Cancer. 2018;18:130.

Zhang BY, Jones JC, Briggler, et al. Lack of caudal-type homeobox transcription factor 2 expression as a prognostic biomarker in metastatic colorectal cancer. Clin Colorectal Cancer. 2016 Sep 17. [Epub ahead of print].

Zhang M and Lin O Molecular testing of thyroid nodules: A review of current available tests for fine-needle aspiration specimens. Archives of Pathology & Laboratory Medicine: December 2016, Vol. 140, No. 12, pp. 1338-1344.

Zheng S, Cherniack AD, Dewal N, et al. Comprehensive pan-genomic characterization of adrenocortical carcinoma. Cancer Cell. 2016 May 9; 29(5): 723–736.

Zhou H, Chen Q, Tan W, et al. Integrated clinicopathological features and gene microarray analysis of pancreatic neuroendocrine tumors. Gene. 2017 May 4. pii: S0378-1119(17)30332-3. [Epub ahead of print].

Policy History/Revision Information

Date	Summary of Changes
10/01/2021	Applicable Codes <ul style="list-style-type: none">Updated list of applicable CPT codes to reflect quarterly edits; added 0262U Supporting Information <ul style="list-style-type: none">Archived previous policy version CS152IN.02

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.