

Proton Beam Radiation Therapy (for Indiana Only)

Policy Number: CS105IN.02
 Effective Date: July 1, 2021

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Applicable Codes	2
Description of Services	3
Clinical Evidence	3
U.S. Food and Drug Administration	13
References	13
Policy History/Revision Information	17
Instructions for Use	17

Related Policy
<ul style="list-style-type: none"> Intensity Modulated Radiation Therapy

Application

This Medical Policy only applies to the state of Indiana.

Coverage Rationale

Proton Beam Radiation Therapy is proven and medically necessary under certain circumstances. For medical necessity clinical coverage criteria, refer to the InterQual® 2021, Apr. 2021 Release, CP: Procedures, Proton Beam Radiotherapy (PBRT).

Click [here](#) to view the InterQual® criteria.

PBT is unproven and not medically necessary due to insufficient evidence of efficacy for treating all other indications not listed above as proven, including but not limited to:

- Age related macular degeneration (AMD)
- Bladder cancer
- Brain and spinal cord tumors
- Breast cancer
- Choroidal hemangioma
- Esophageal cancer
- Gynecologic cancers
- Head and neck tumors not noted above as proven
- Lung cancer
- Lymphomas
- Pancreatic cancer
- Vestibular tumors (e.g., acoustic neuroma or vestibular schwannoma)
- PBT used in conjunction with IMRT

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
77301	Intensity modulated radiotherapy plan, including dose-volume histograms for target and critical structure partial tolerance specifications
77338	Multi-leaf collimator (MLC) device(s) for intensity modulated radiation therapy (IMRT), design and construction per IMRT plan
77385	Intensity modulated radiation treatment delivery (IMRT), includes guidance and tracking, when performed; simple
77386	Intensity modulated radiation treatment delivery (IMRT), includes guidance and tracking, when performed; complex
77387	Guidance for localization of target volume for delivery of radiation treatment, includes intrafraction tracking, when performed
77520	Proton treatment delivery; simple, without compensation
77522	Proton treatment delivery; simple, with compensation
77523	Proton treatment delivery; intermediate
77525	Proton treatment delivery; complex

CPT® is a registered trademark of the American Medical Association

HCPCS Code	Description
G6015	Intensity modulated treatment delivery, single or multiple fields/arcs, via narrow spatially and temporally modulated beams, binary, dynamic MLC, per treatment session
G6016	Compensator-based beam modulation treatment delivery of inverse planned treatment using 3 or more high resolution (milled or cast) compensator, convergent beam modulated fields, per treatment session
G6017	Intra-fraction localization and tracking of target or patient motion during delivery of radiation therapy (e.g., 3D positional tracking, gating, 3D surface tracking), each fraction of treatment

Diagnosis Code	Description
C11.0	Malignant neoplasm of superior wall of nasopharynx
C11.1	Malignant neoplasm of posterior wall of nasopharynx
C11.2	Malignant neoplasm of lateral wall of nasopharynx
C11.3	Malignant neoplasm of anterior wall of nasopharynx
C11.8	Malignant neoplasm of overlapping sites of nasopharynx
C11.9	Malignant neoplasm of nasopharynx, unspecified
C22.0	Liver cell carcinoma
C30.0	Malignant neoplasm of nasal cavity
C31.0	Malignant neoplasm of maxillary sinus
C31.1	Malignant neoplasm of ethmoidal sinus
C31.2	Malignant neoplasm of frontal sinus
C31.3	Malignant neoplasm of sphenoid sinus
C31.8	Malignant neoplasm of overlapping sites of accessory sinuses

Diagnosis Code	Description
C31.9	Malignant neoplasm of accessory sinus, unspecified
C41.0	Malignant neoplasm of bones of skull and face
C61.0	Malignant neoplasm of prostate
C69.30	Malignant neoplasm of unspecified choroid
C69.31	Malignant neoplasm of right choroid
C69.32	Malignant neoplasm of left choroid
C69.40	Malignant neoplasm of unspecified ciliary body
C69.41	Malignant neoplasm of right ciliary body
C69.42	Malignant neoplasm of left ciliary body
D09.20	Carcinoma in situ of unspecified eye
D09.21	Carcinoma in situ of right eye
D09.22	Carcinoma in situ of left eye
D14.0	Benign neoplasm of middle ear, nasal cavity and accessory sinuses
D16.4	Benign neoplasm of bones of skull and face
D31.30	Benign neoplasm of unspecified choroid
D31.31	Benign neoplasm of right choroid
D31.32	Benign neoplasm of left choroid
D31.40	Benign neoplasm of unspecified ciliary body
D31.41	Benign neoplasm of right ciliary body
D31.42	Benign neoplasm of left ciliary body
Q28.2	Arteriovenous malformation of cerebral vessels
Q28.3	Other malformations of cerebral vessels

Description of Services

Unlike other types of radiation therapy (RT) that use x-rays or photons to destroy cancer cells, proton beam therapy (PBT) uses a beam of special particles (protons) that carry a positive charge. There is no significant difference in the biological effects of protons versus photons; however, protons can deliver a dose of radiation in a more confined way to the tumor tissue than photons. After they enter the body, protons release most of their energy within the tumor region and, unlike photons, deliver only a minimal dose beyond the tumor boundaries (American College of Radiology (ACR) website, updated 2019).

Proton beam radiation therapy (PBRT) is intended to deliver higher, more targeted radiation with less damage to collateral healthy tissue than external beam radiation therapy (EBRT) using photons (x-rays) when used to treat solid tumors. While PBRT has been used for several solid cancer tumor types (e.g., breast, lung, prostate, head and neck, central nervous system (CNS)) in adults and in certain pediatric cancers, evidence is lacking regarding clear benefits over EBRT (ECRI, 2017).

Clinical Evidence

Unproven Indications

Age-Related Macular Degeneration (AMD)

Evans et al. (2020) updated a previously conducted systematic review (Evans, 2010) that examined the effects of radiotherapy on neovascular AMD. A search was conducted using CENTRAL, MEDLINE, Embase, LILACS and three trials registers for randomized controlled trials in which radiotherapy was compared to another treatment, sham treatment, low dosage irradiation or no treatment in people with choroidal neovascularisation (CNV) secondary to AMD. Outcomes included best-corrected visual acuity (BCVA) (loss of 3 or more lines, change in visual acuity), contrast sensitivity, new vessel growth, quality of life and adverse effects at any time point. A total of 18 studies (n=2,430 people, 2,432 eyes) were included, and the radiation therapy

with dosages ranging from 7.5 to 24 Gy. Three of these studies investigated brachytherapy (plaque and epimacular), the rest were studies of external beam radiotherapy (EBM) including one trial of stereotactic radiotherapy. The authors concluded that the evidence is uncertain regarding the use of radiotherapy for neovascular AMD. They stated that: 1) most studies took place before the routine use of anti-VEGF, and before the development of modern radiotherapy techniques such as stereotactic radiotherapy; 2) visual outcomes with epimacular brachytherapy are likely to be worse, with an increased risk of adverse events, probably related to vitrectomy; 3) the role of stereotactic radiotherapy combined with anti-VEGF is currently uncertain; and 4) further research on radiotherapy for neovascular AMD may not be justified until current ongoing studies have reported their results.

In a systematic review, Bekkering et al. (2009) evaluated the effects and side effects of PBT for indications of the eye. All studies that included at least 10 patients and that assessed the efficacy or safety of PBT for any indication of the eye were included. Five controlled trials, 2 comparative studies and 30 case series were found, most often reporting on uveal melanoma, choroidal melanoma and AMD. Methodological quality of these studies was poor. Studies were characterized by large differences in radiation techniques applied within the studies, and by variation in patient characteristics within and between studies. Results for uveal melanoma and choroidal melanoma suggest favorable survival, although side effects are significant. Results for choroidal hemangioma and AMD did not reveal beneficial effects from proton radiation. There is limited evidence on the effectiveness and safety of due to the lack of well-designed and well-reported studies.

A RCT by Zambarakji et al. (2006) studied 166 patients with angiographic evidence of classic choroidal neovascularization resulting from AMD and best-corrected visual acuity of 20/320 or better. Patients were assigned randomly (1:1) to receive 16-cobalt gray equivalent (CGE) or 24-CGE PBT in 2 equal fractions. Complete ophthalmological examinations, color fundus photography, and fluorescein angiography were performed before and 3, 6, 12, 18, and 24 months after treatment. At 12 months after treatment, 36 eyes (42%) and 27 eyes (35%) lost three or more lines of vision in the 16-CGE and 24-CGE groups, respectively. Rates increased to 62% in the 16-CGE group and 53% in the 24-CGE group by 24 months after treatment. Radiation complications developed in 15.7% of patients receiving 16-CGE and 14.8% of patients receiving 24-CGE. The authors concluded that no significant differences in rates of visual loss were found between the 2 dose groups.

Clinical Practice Guidelines

American Academy of Ophthalmology (AAO)

AAO preferred practice patterns state that RT is not recommended in the treatment of AMD (2019).

Bladder Cancer

Takaoka and colleagues (2017) conducted a retrospective review to assess outcomes, prognostic factors and toxicities of PBT as a component of trimodal bladder-preserving therapy for muscle-invasive bladder cancer. Trimodal bladder-preserving therapy consisted of maximal transurethral resection of the bladder tumor, small pelvis (conventional) photon radiation, intra-arterial chemotherapy and PBT. Seventy patients with cT2-3N0M0 muscle-invasive bladder cancer were included who received treatment from 1990 to 2015 at a single institution. The OS and PFS rate, time to progression, predictive factors for progression and toxicities were analyzed. Progression was defined as when muscle-invasive recurrence, distant metastasis or upper urinary tract recurrence was observed. The patients' median age was 65 (range 36-85) years. The median follow-up period was 3.4 years (range 0.6-19.5 years). The 5-year cumulative OS rate, PFS rate and time to progression rate were 82%, 77%, and 82%, respectively. In univariate and multivariate analyses, tumor multiplicity and tumor size (≥ 5 cm) were significant and independent factors associated with progression (hazard ratio 3.5, 95% confidence interval 1.1-12; hazard ratio 5.0, 95% confidence interval 1.3-17; $P < 0.05$ for all). As for toxicity, 26 (18%) patients had grade 3-4 acute hematologic toxicities and 2 (3%) patients had grade 3 late GU toxicity. No patient had to discontinue the treatment due to acute toxicity. The authors concluded that trimodal therapy including both conventional and proton radiation was well tolerated and may be an effective treatment option for selected muscle-invasive bladder cancer patients. Further studies are needed to determine whether PBT is integral to this multimodality therapy.

Miyanaga et al. (2000) conducted a small prospective uncontrolled clinical study to assess the efficacy and safety of PBT and/or conventional photon therapy for bladder cancer. The study involved 42 patients who received PBT to the small pelvic space following intra-arterial chemotherapy. At 5-year follow-up, the bladder was preserved in 76% of patients and 65% were free of disease. The disease-specific survival rate was 91%. Patients with large and multiple tumors were more at risk of cancer

recurrence than patients with single, small tumors. Nausea and vomiting, irritable bladder and ischialgia were the main side effects.

NCCN guidelines do not address the use of PBT for treating bladder cancer (2020).

Brain and Spinal Cord Tumors

Petr et al. (2018) assessed structural and hemodynamic changes of healthy brain tissue in the cerebral hemisphere contralateral to the tumor following conventional (photon) and proton radiation with concurrent chemotherapy. Sixty seven adult patients diagnosed with glioblastoma undergoing adjuvant conventional (n=47) or proton (n=19) radiotherapy with temozolomide after tumor resection underwent T1-weighted and arterial spin labeling magnetic resonance imaging. Changes in volume and perfusion before and 3-6 months after were compared between therapies. A decrease in gray matter (GM) and white matter (WM) volume was observed in patients receiving conventional radiation compared to the pre-RT baseline. In contrast, for the proton therapy group, no significant differences in GM or WM volume were observed. GM volume decreased with 0.9% per 10 Gy dose increase and differed between the radiation modalities. Perfusion decreased in conventional radiation therapy patients, whereas the decrease in proton therapy patients was not statistically significant. There was no correlation between perfusion decrease and either dose or radiation modality. The authors concluded that proton therapy may reduce brain volume loss compared to photon therapy, with decrease in perfusion being comparable for both modalities. As this was an uncontrolled retrospective study with a surrogate end-point (brain volume loss on imaging), prospective randomized trials are needed to compare the effect of proton and conventional radiotherapy on imaging and clinical outcomes.

Kabolizadeh et al. (2017) conducted a single-center, retrospective, case series to evaluate local control (LC), overall survival (OS), disease-specific survival, and distant failure in 40 patients with unresected chordoma and treated with photon/proton radiation therapy. Tumor response was assessed using the modified Response Evaluation Criteria in Solid Tumors (mRECIST). To characterize tumor response the soft tissue and bone compartments of the tumor were defined separately as the soft tissue target volume, bone target volume and combined total target volume. Twenty-seven patients had sacrococcygeal chordoma, and the remaining patients had mobile spine tumors, which included 9 cervical, 1 thoracic, and 3 lumbar. Thirty-nine patients underwent proton therapy only or predominantly proton therapy mixed with photons to limit the radiation dose to adjacent critical normal structures. Only 4 patients received either concurrent or neoadjuvant systemic treatments. The median age was 67 years (range, 36-94 years) and median follow-up, after completion of radiation therapy, was 50.3 months (range, 2–216.4 months). At 5-years, LC, OS, disease-specific survival, and distant failure were 85.4%, 81.9%, 89.4%, and 20.2%, respectively. Nineteen patients had complete sets of regular imaging scans (a total of 84 CT and MRI scans were reviewed) and of those, only 4 local failures had occurred at 34, 46, 78 and 82 months after treatment. The authors concluded that their results support the use of high-dose definitive radiation therapy in select patients with unresected spine and sacral chordomas, and that soft tissue target volume is the best indicator of tumor response. Limitations of this study include its design, the small number of patients with local failure and limited follow-up periods.

Indelicato et al. (2016) conducted descriptive analysis using data from a single-institution. In this prospective case series study, researchers sought to evaluate the effectiveness of definitive or adjuvant external beam proton therapy in patients with chordomas and chondrosarcomas of the spine. Outcomes of interest included distant metastases (DM), overall survival (OS), cause-specific survival, local control (LC) and disease-free survival (DFS). A total of 51 patients participated with a median age of 58 years (range, 22-83 years) and median follow-up of 3.7 years (range, 0.3-7.7 years). There were 34 patients with chordomas, and 17 patients with chondrosarcomas, which were all grade 2 or higher. The anatomic distribution was as follows: sacrum (n=21), cervical spine (n=20), and thoracolumbar spine (n=10). The median dose of radiation therapy was 70.2 Gy (range, 64.2-75.6 Gy). The 4-year LC, freedom from distant metastases, DFS, cause-specific survival, and OS rates were 58%, 86%, 57%, 72%, and 72%, respectively. A total of 25 patients experienced disease recurrence: 18 local recurrences, 6 local and distant recurrences, and 1 DM. In patients with a local relapse, the median time to progression was 1.7 years (range, 0.2-6 years). The median survival after local progression was 1.7 years (range, 0.1-4.9+ years). Regression analysis results showed that younger patients had a significantly higher risk for local reoccurrence and that patients whose initial management was only surgery also had a higher rate of reoccurrence however, these patients may represent a high-risk subset. The authors concluded that high-dose proton therapy controls more than half of spinal chordomas and chondrosarcomas and compares favorably with historic photon data. Local progression is the dominant mode of treatment failure and it may be reduced by treating patients at the time of initial diagnosis. Limitations of this study include its design, small sample size and small number of select events, which may have impacted the statistical validity of the regression analysis results.

Noel et al. (2002) conducted a retrospective review of 17 patients with meningioma to evaluate the efficacy and the tolerance of an escalated dose of external conformal fractionated RT combining photons and protons. Five patients presented a histologically atypical or malignant meningioma, 12 patients had a benign tumor that was recurrent or rapidly progressive. In 2 cases, RT was administered in the initial course of the disease and in 15 cases at the time of relapse. A highly conformal approach was used combining high-energy photons and protons for approximately 2/3 and 1/3 of the total dose. The median total dose delivered within gross tumor volume was 61 CGE (25-69). Median follow-up was 37 months (17-60). The 4-year LC and OS rates were 87.5 +/- 12% and 88.9 +/- 11%, respectively. Radiologically, there were 11 stable diseases and 5 partial responses. The authors concluded that in both benign and more aggressive meningiomas, the combination of conformal photons and protons with a dose escalated by 10-15% offers clinical improvements in most patients as well as radiological long-term stabilization. Limitations of this study include small sample size and study design.

NCCN guidelines for CNS cancers states that when toxicity is a concern during management of spinal ependymoma or medulloblastoma in adults, PBRT should be considered if available (2020).

Several clinical trials studying PBT in patients with various types of brain tumors are active or recruiting. For more information, go to www.clinicaltrials.gov. (Accessed February 16, 2021)

Breast Cancer

DeCesaris (2019) conducted single-institution, retrospective cohort analysis to evaluate acute skin toxicity, i.e., radiation dermatitis (RD) or skin hyperpigmentation (SH) in patients with primary invasive breast cancer who underwent radiation therapy with either photon or proton radiation therapy. Skin toxicity was recorded using Common Terminology Criteria for Adverse Events version 4.0 criteria and scored by treating physicians on a weekly basis. For each patient, the highest recorded grades of RD and SH were analyzed. A total of 86 patients received treatment with a median age of 53 years (range, 24 – 78 years) and median RT dose of 60 Gy (range, 45 – 70 Gy). Of those, 47 (55%) received photon beam therapy and 39 (45%) received proton beam therapy. Patients treated with proton beam radiation therapy had a statistically significant higher rate of grade \geq 2 RD compared with patients who were treated with photon radiation therapy (69.2% vs. 29.8%, $p < 0.001$). There was no difference in the rates of grade 3 RD or SH between the modalities. The authors concluded that women who will be undergoing proton beam radiation therapy should receive counseling regarding its potential for grade \geq 2 skin toxicities. Limitations of this study include its design, use of subjective assessments, and that during treatment optically stimulated luminescent dosimeters were not used to measure patients' radiation exposure.

Verma et al. (2017) conducted a single-institution retrospective cohort study to evaluate acute toxicity in patients with locally advanced breast cancer and receiving comprehensive regional nodal irradiation (CRNI) with adjuvant PBT from 2011–2016. PBT targeting the intact breast/chest wall and CRNI including the axilla, supraclavicular fossa, and internal mammary lymph nodes consisted of a 3-dimensional uniform scanning technique. In 2016, the institution transitioned to a pencil beam scanning (PBS) technique. The change in technique was driven by anticipated dosimetric advantages including decreased dose to the skin surface and to cardiopulmonary organs, and shorter planning and treatment delivery time. Toxicities were assessed weekly during treatment, one month following treatment completion, and then, every 6 months. A total of 91 patients were treated with a median follow-up period of 15.5 months. The most common toxicities were dermatitis and/or skin infections, but esophagitis and fatigue were also observed. Acute dermatitis of grades 1, 2, and 3 occurred in 23%, 72%, and 5%, respectively. Eight percent ($n=7$) required treatment breaks due to dermatitis and the median time to resolution of acute skin toxicity was 32 days. Grades 1, 2, and 3 esophagitis developed in 31%, 33%, and 0%, respectively. The authors concluded that PBT for breast cancer as part of CRNI appears to have toxicity rates comparable to prior published studies e.g., Cuaron et al. (2015) reported 71.4% of those who received PBT developed grade 2 dermatitis however, Bradley et al. (2016) reported 100% developed grade 2 dermatitis. While the use of PBT with CRNI may have dosimetric advantages, particularly to the heart and other organs at risk, toxicities observed with its use demonstrates the need for randomized controlled trials comparing PBT to other radiation modalities.

Verma et al. (2016a) performed a systematic review of clinical outcomes and toxicity of PBT for treating breast cancer. Nine original studies were analyzed, however the types of studies and the volume of patients in those studies were not specifically cited by the authors. Conventionally fractionated breast/chest wall PBT produced grade 1 dermatitis rates of approximately 25% and grade 2 dermatitis in 71%-75%. This is comparable or improved over the published rates for photons. The incidence of esophagitis was decreased if the target coverage was compromised in the medial supraclavicular volume, a finding that echoes previous results with photon RT. From the limited available data, the rate of grade 2 esophagitis ranged from 12% to 29%. Using

PBT-based accelerated partial breast irradiation (PBI), the rates of seroma/hematoma and fat necrosis were comparable to those reported in the existing data. Radiation pneumonitis (RP) and rib fractures remain rare. PBT offers the potential to minimize the risk of cardiac events, keeping the mean heart dose at ≤ 1 Gy. However, definitive clinical experiences remain sparse. Results from clinical trials in progress, comparing protons to photons, will further aid in providing conclusions. Limitations to this review included a general lack of data and low number of participants in the available studies.

Bradley et al. (2016) conducted a prospective case series study to evaluate the clinical feasibility and potential benefits of PBT in breast cancer patients who were at risk for regional nodal disease. In this pilot study, the primary endpoint was cardiac V5, testing the hypothesis that PBT could reduce the volume of the heart receiving 5 Gy by $\geq 50\%$ when compared to CRT. The secondary endpoints included acute toxicity and other dosimetric parameters of target coverage and exposure to at-risk organs. PBT and CRT plans, targeting the regional nodes, were created for each patient. Patients were evaluated weekly while on RT, 4 weeks after RT was completed and at 6-month intervals thereafter. Toxicity was recorded using the Common Terminology Criteria for Adverse Events (CTCAE, v4.0). A total of 18 women enrolled with a median age of 51.8 years (range, 42–73 years) and a median follow-up period of 20 months (range, 2–31 months). Ten of the women received only PBT and 8 received combination therapy of PBT and photon beam RT. All patients had improved heart and lung dose with PBT. The primary endpoint, which was to determine if PBT could reduce cardiac V5 by $\geq 50\%$, was achieved. Of the nine patients with left-sided breast cancer, the median cardiac dose decreased from 5.9 Gy with CRT to 0.6 Gy with PBT ($p=0.004$). In patients with right-sided breast cancer, the median cardiac dose decreased from 2.9 Gy with CRT to 0.5 Gy with PBT ($p=0.004$). No patients developed grade 4+ toxicities. Four (22%) patients developed grade 3 dermatitis and of these, 3 were treated with PBT and 1 was treated with combination PBT and CRT. All of the patients developed grade 2 dermatitis, which resolved within 1 month of the completion of therapy. However, 1 patient developed cellulitis and required a course of antibiotics. Additional acute grade 2 toxicities included: fatigue ($n=6$), esophagitis ($n=5$), nausea ($n=1$) and dyspnea ($n=1$). The authors acknowledged that their rate of patients with grade 3 acute skin toxicity was not unexpected given the higher skin dose with PBT and concluded that PBT for regional node irradiation after mastectomy or breast conserving surgery offers a lower cardiac dose particularly for patients with left-sided breast cancer and without grade 4+ toxicities. Limitations of this study include its design, small sample size and higher toxicity rates compared with other forms of RT, e.g., intensity modulated RT (DeCesaris 2019).

Cuaron et al. (2015) conducted a single-institution case series study to report dosimetry and early toxicity data in patients with breast cancer. Retrospectively collected data from consecutive patients diagnosed with non-metastatic breast cancer, no prior history of chest wall radiation and treated with PBT postoperatively were studied. Patients with unfavorable cardiopulmonary anatomy were usually referred to this institution. Post-lumpectomy patients with large breast size were not offered treatment due to a higher propensity for day-to-day measurement differences in the target position. Patients were evaluated weekly while on RT, 4 weeks after RT was completed, and at 12–24 week intervals thereafter. Toxicity was recorded using the Common Terminology Criteria for Adverse Events (CTCAE, v4.0). A total of 30 women were included in the study with a median age of 49 years (range, 29–86 years), cancer staging was as follows: 8 had stage II, 20 had stage III and 2 had chest wall recurrence. The median follow-up was 9.3 months (range, 2.3–18.6 months). With PBT, full coverage of the planned target value was achieved, and it significantly spared the heart, lungs and contralateral breast. Of those with greater than 3 months of follow-up ($n=28$), 71.4% developed grade 2 dermatitis and of those, 28.6% experienced moist desquamation. Eight (28.6%) developed grade 2 esophagitis and 1 developed grade 3 reconstructive complications. The authors concluded that in this series of 30 patients, PBT achieved excellent coverage of the target volume while sparing the heart, lungs, and contralateral breast, that the treatment was well tolerated, and that additional studies assessing long-term outcomes and toxicity are needed. Limitations of this study include its design, exclusion of women with large breast size, and higher toxicity rates compared with other forms of RT, e.g., intensity modulated RT (DeCesaris 2019).

Bush et al. (2014) performed a single center study of 100 subjects who received postoperative PBI using PBT after undergoing partial mastectomy with negative margins and axillary lymph nodes. After following these individuals for an average of 5 years, the researchers concluded that ipsilateral recurrence-free survival with minimal toxicity was excellent. While the authors acknowledged that cosmetic results may be improved with PBT over those reported with photon-based techniques, there was nothing in the study demonstrating that PBT outcomes were superior to the current standard of care.

NCCN guidelines do not address the use of PBT for treating breast cancer (2020).

To further elucidate the clinical advantages and disadvantages between PBT and other types of radiation therapy used in breast cancer, additional clinical trials are underway, NCT02603341, NCT01245712, and NCT03391388, go to www.clinicaltrials.gov. (Accessed February 16, 2021).

Choroidal Hemangiomas

Hocht et al. (2006) conducted a single-center, retrospective study of 44 consecutive patients with choroid hemangiomas treated with photon therapy (n=19) or proton therapy (n=25). Outcomes were measured by visual acuity, tumor thickness, resolution of retinal detachment, and post-treatment complications. Mean follow-up was 38.9 months and 26.3 months, and median follow-up was 29 months and 23.7 months for photon and proton patients, respectively. Tumor thickness was greater in the photon group than in the proton group. In the collective groups, 91% were treated successfully, and there was no significant difference in the outcomes between the 2 groups. The authors concluded that RT is effective in treating choroidal hemangiomas with respect to visual acuity and tumor thickness, but a benefit of proton versus photon therapy could not be detected.

Three additional studies showed some improvement in tumor regression and visual acuity following PBT; however, these studies were small and retrospective in nature (Chan et al., 2010; Levy-Gabriel et al., 2009; Frau et al., 2004).

Gastrointestinal (GI) Cancers

Verma et al. (2016b) conducted a systematic review to identify studies on PBT and gastrointestinal malignancies. The search included PubMed, EMBASE, and abstracts from meetings of the American Society for Radiation Oncology, Particle Therapy Co-Operative Group, and American Society of Clinical Oncology. A total of 39 original investigations were analyzed. For esophageal cancer, 12 studies were analyzed and several of those reported that PBT resulted in a significant dose reduction to intrathoracic OARs and is associated with reduced toxicity, postoperative complications while achieving comparable local control and overall survival. However, for some of the studies, contemporaneous comparison groups were lacking or comparisons were made between PBT and x-ray radiotherapy (XRT), which consisted of either 3D-CRT or IMRT rather than IMRT only. For pancreatic cancer, 5 studies were analyzed. Survival for resected/unresected cases was similar to existing data where IMRT was used and nausea/emetesis were numerically lower than what had been reported among patients who received IMRT however, direct head-to-head comparisons were not made. For hepatocellular carcinoma, 10 studies were analyzed and these had the strongest evidence to support use of PBT. Those studies reported very low toxicities, and a phase III trial comparing PBT to transarterial chemoembolization (TACE) showed a trend toward better LC and PFS with PBT. For cholangiocarcinoma, liver metastases, and retroperitoneal sarcoma, survival and toxicity data is comparable to historical photon controls, and stomach and biliary system/gallbladder cancer studies consisted of case reports and small cohort experiences. The authors concluded that PBT offers the potential of lower toxicities without compromising survival or local control. However, there was limited high quality evidence for select gastrointestinal malignancies and that multi-institution, randomized controlled trials are needed.

NCCN guidelines do not address PBT in the treatment of gastric cancers (2020).

Esophageal Cancer

Xi et al. (2017) conducted a single-center retrospective cohort study to evaluate outcomes of patients diagnosed with esophageal cancer (EC) and treated with PBT or IMRT. Outcomes included treatment-related toxicity, OS, PFS, locoregional failure-free survival (LRFSS) and distant metastasis-free survival (DMFS). Patients were followed every three months for the first year after radiation therapy, every 6 months for the following 2 years and then yearly until 5 years. A total of 343 patients were included and of those ,211 received IMRT and 132 received PBT. The median follow-up period for the IMRT group was 65.1 months (range, 19.4-115.3) and for the PBT group was 44.8 months (range, 11.9 – 110.3 months). The median radiation dose was 50.4 Gy in both the IMRT and PBT groups (ranges, 41.4-66.0 Gy and 45.0-63.0 Gy, respectively). There was no difference in treatment-related toxicities between the groups. The PBT group had better OS (p=.011), PFS (p=.001), and DMFS (p=.031) compared with the IMRT group. In subset analyses, patients with stage I/II disease had no differences in survival. In patients with stage III disease, those who received PBT had higher rates of OS (34.6% vs. 25.0%, p=.038) and PFS (33.5% vs. 13.2%, p=.005). The authors concluded that PBT was associated with improved OS, PFS and LRFSS, particularly in EC patients with advanced disease and that their results may suggest a benefit of PBT over IMRT. Limitations of this study include its design, that the type of radiation therapy each patient received was based on the multidisciplinary team and the patients' intent rather than randomization, there were differences in patient demographics and baseline characteristics between the groups, and that

for some patients, accurate long term documentation was lacking. Prospective, randomized controlled studies are needed to clarify the role of PBT in EC.

Lin et al. (2017) conducted a multi-center retrospective cohort study of patients diagnosed with EC and treated with neoadjuvant chemoradiation. The purpose of this study was to assess the association between RT modality and postoperative outcomes. The outcomes included pulmonary, cardiac and wound complications, and length of stay (LOS), readmission and mortality. A total of 580 EC patients were included and of these, 214 (37%) received 3D-CRT, 255 (44%) received IMRT and 111 (19%) receive PBT. IMRT and PBT were associated with a reduced risk of pulmonary complications compared with 3D-CRT ($p=.001$), and PBT was trending toward being better than IMRT (OR 0.584, $p=.077$). Both IMRT and PBT were associated with a reduced risk of cardiac complications as were older age and history of coronary artery bypass grafting or atrial fibrillation. PBT was associated with a reduced risk of wound complications (OR 0.255, $p=0.006$, PBT vs. 3D-CRT; OR 0.276, $p=0.009$, PBT vs. IMRT) yet there was no difference between IMRT and 3D-CRT. Mean LOS was significantly associated with RT modality (13.2 days for 3D-CRT, 11.6 days for IMRT and 9.3 days for PBT ($p<0.0001$). There was no difference in 60-day readmission rates or deaths during the same hospitalization, or 30, 60 or 90-day postoperative mortality. The authors concluded that IMRT and PBT were associated with significantly reduced rates of postoperative complications compared to 3D-CRT, that these results may show an advantage of PBT over IMRT however, prospective randomized clinical trials will better establish the role of PBT in EC.

In a retrospective analysis, Wang et al. (2013a) reported that advanced radiation technologies such as IMRT or PBT significantly reduced postoperative pulmonary and GI complication rates compared to 3D-CRT in esophageal cancer patients. These results need to be confirmed in prospective studies.

Mizumoto et al. (2011) evaluated the efficacy and safety of hyperfractionated concomitant boost PBT in 19 patients with esophageal cancer. The overall 1- and 5-year actuarial survival rates for all 19 patients were 79% and 42.8%, respectively. The median survival time was 31.5 months. Of the 19 patients, 17 (89%) showed a complete response within 4 months after completing treatment and 2 (11%) showed a partial response, giving a response rate of 100% (19/19). The 1- and 5-year LC rates for all 19 patients were 93.8% and 84.4%, respectively. The results suggest that hyperfractionated PBT is safe and effective for patients with esophageal cancer. Further studies are needed to establish the appropriate role and treatment schedule for use of PBT for esophageal cancer.

Mizumoto et al. (2010) evaluated the efficacy and safety of PBT for locoregionally advanced esophageal cancer. Fifty-one patients were treated using PBT with or without X-rays. All but one had squamous cell carcinoma. Of the 51 patients, 33 received combinations of X-rays and protons as a boost. The other 18 patients received PBT alone. The overall 5-year actuarial survival rate for the 51 patients was 21.1% and the median survival time was 20.5 months. Of the 51 patients, 40 (78%) showed a complete response within 4 months after completing treatment and seven (14%) showed a partial response, giving a response rate of 92% (47/51). The 5-year LC rate for all 51 patients was 38% and the median LC time was 25.5 months. The authors concluded that these results suggest that PBT is an effective treatment for patients with locally advanced esophageal cancer. Further studies are required to determine the optimal total dose, fractionation schedules and best combination of proton therapy with chemotherapy.

NCCN guidelines state that PBT is appropriate when treating esophageal and esophagogastric junction cancers in settings where dose reduction to organs at risk is necessary and cannot be achieved by 3D-CRT. Because data is early and evolving, patients should receive PBT within a clinical trial (2019).

Gynecologic Cancers

The efficacy of PBT combined with photon radiation for the treatment of cervical cancer was investigated in a prospective uncontrolled study involving 25 patients (Kagei et al., 2003). In this study, 5-year and 10-year survival rates were similar to conventional therapies as reported in the literature. The 10-year survival rate was higher for patients with low stage (89%) compared with advanced stages (40%) of cervical cancer. The treatment caused severe late complications in 4% of patients.

NCCN guidelines do not address the use of PBT when treating any type of gynecologic cancer (i.e., Cervical Cancer (2021), Ovarian Cancer (2020), Uterine Neoplasms (2021) or Vulvar Cancer (2021).

Several clinical trials are recruiting or in progress studying the use of PBT in multiple types of gynecologic cancer (e.g., cervical, ovarian, and uterine). For more information, go to www.clinicaltrials.gov. (Accessed February 16, 2021).

Head and Neck Cancers (HNC) not listed in the coverage rationale as proven

A 2019 Hayes report, Proton Beam Therapy for Treatment of Head and Neck Cancer, assessed multiple clinical studies evaluating the efficacy and safety of PBT in patients with HNC. The majority of the evidence included retrospective studies, data analyses, and systematic reviews. They noted there was some overlap of investigators and, possibly, overlap of patient groups as well. The report concludes that the study abstracts present conflicting findings regarding the use of PBT for treatment of HNC.

Seeking to improve LC rate and reduce late AEs, Takayama et al. (2016) evaluated therapeutic results and toxicities of PBT combined with selective intra-arterial infusion chemotherapy (PBT-IACT) in patients with stage III-IVB squamous cell carcinoma of the tongue. Between February 2009 and September 2012, 33 patients were enrolled. After 2 systemic chemotherapy courses and whole-neck irradiation (36 Gy in 20 fractions), participants were administered concurrent chemoradiotherapy comprising PBT for the primary tumor and the metastatic neck lymph node with weekly retrograde IACT of cisplatin with sodium thiosulfate by continuous infusion. The median follow-up duration was 43 months. The 3-year OS, PFS, LC rate, and regional control rate for the neck were 87%, 74.1%, 86.6%, and 83.9%, respectively. Major acute toxicities > grade 3 included mucositis in 26 cases (79%), neutropenia in 17 cases (51%), and dermatitis in 11 cases (33%). Late grade 2 osteoradionecrosis was observed in 1 case (3%). The authors concluded that PBT-IACT for stage III-IVB tongue cancer has an acceptable toxicity profile and showed good treatment results, and that this protocol should be considered as a treatment option for locally advanced tongue cancer. This study is limited by the lack of data comparing toxicity to conventional radiation therapy.

NCCN's HNC guideline makes no mention of proton beam radiation therapy for cancer of the lip (mucosa), oral cavity, hypopharynx or glottic larynx. The guideline states that use of proton therapy is an active area of investigation, and that proton therapy may be considered when normal tissue constraints cannot be met by photon-based therapy in cancers of the oropharynx, nasopharynx, supraglottic larynx, paranasal/ethmoid sinus, maxillary sinus and salivary glands, as well as mucosal melanoma and other primary tumors of the head and neck (2020).

Clinical Practice Guidelines

American College of Radiology (ACR) / American Society for Radiation Oncology (ASTRO)

Regarding head and neck tumors, the ACR/ASTRO practice parameter states that PBRT reduces the dose delivered to critical normal structures in the head and neck region that may impact QOL, including optic nerves, optic chiasm, pituitary gland, brain, brainstem, spinal cord, salivary glands, pharyngeal constrictor muscles, oral cavity, and the emetogenic sites in the posterior fossa (2018).

American Society for Radiation Oncology (ASTRO)

ASTRO's model policy supports the use of PBT in the treatment of malignant and benign primary CNS tumors, advanced (e.g., T4) and/or unresectable head and neck cancers, and cancers of the paranasal sinuses and other accessory sinuses (2017). (Accessed February 17, 2021).

Lung Cancer

A Hayes report (2020) concluded that the best available studies of PBT for NSCLC do not provide sufficient evidence that PBT is safer or consistently more effective than CRT and IMRT in the treatment of NSCLC.

Liao et al. (2018) conducted a single-center randomized trial that compared outcomes of passive scattering proton therapy (PSPT) versus IMRT, both with concurrent chemotherapy, for inoperable non-small-cell lung cancer (NSCLC). The primary end point was the first occurrence of severe (grade ≥ 3) radiation pneumonitis (RP) or local failure (LF). Eligible patients had stage IIB to IIIB NSCLC (or stage IV NSCLC with a single brain metastasis or recurrent lung or mediastinal disease after surgery) and were candidates for concurrent chemoradiation therapy. Pairs of treatment plans for IMRT and PSPT were created for each patient. Patients were eligible for random assignment only if both plans satisfied the same prespecified dose-volume constraints for at-risk organs at the same tumor dose. Compared with IMRT (n=92), PSPT (n=57) exposed less lung tissue to doses of 5 to 10 Gy(RBE), which is the absorbed Gy dose multiplied by the relative biologic effectiveness (RBE) factor for protons; exposed more lung tissue to ≥ 20 Gy(RBE), but exposed less heart tissue at all dose levels between 5 and 80 Gy (RBE). The grade ≥ 3 RP was greater for PSPT than IMRT (6.5% for IMRT and 10.5% for PSPT) though the difference did not reach statistical

significance; there was no difference observed in LF (10.9% and 10.5% for IMRT and PSPT, respectively). Exploratory analysis showed that the RP and LF rates at 12 months for patients enrolled before versus after the trial midpoint were 21.1% (before) versus 18.2% (after) for the IMRT group and 31.0% (before) versus 13.1% (after) for the PSPT group suggesting that that outcomes for proton therapy improved over the course of the trial as the investigators gained experience. The authors stated that findings from 2 ongoing trials (NCT01993810 and NCT01629498) will provide additional evidence of the efficacy of proton and photon therapies.

Chang et al. (2017) reported 5-year results of a prospective phase II single-institution study evaluating chemotherapy with concurrent high dose PBT in 64 patients with unresectable phase III NSCLC. 5-year OS, PFS, actuarial distant metastases and locoregional recurrence were 29%, 22%, 54%, and 28%, respectively. Acute and late toxic effects with PBT (compared to historical studies with 3D-CRT and/or IMRT) with chemotherapy were very promising. The authors concluded that the study demonstrated that concurrent PBT and chemotherapy was safe and effective in the long term, and that further prospective studies are warranted.

Chi et al. (2017) conducted a systematic review and meta-analysis to assess hypo-fractionated PBT's efficacy relative to that of photon SBRT for early stage NSCLC. Seventy two SBRT studies and 9 hypo-fractionated PBT studies (mostly single-arm) were included. PBT was associated with improved OS and PFS in the univariate meta-analysis. The OS benefit did not reach its statistical significance after inclusion of operability into the final multivariate meta-analysis, while the 3-year LC still favored PBT. Researchers concluded that although hypo-fractionated PBT may lead to additional clinical benefit when compared with photon SBRT, no statistically significant survival benefit from PBT over photon SBRT was observed in the treatment of early stage NSCLC.

Harada et al. (2016) conducted a single-institutional, open label, dose escalation phase I trial to determine the recommended dose of PBT for inoperable stage III NSCLC. Two prescribed doses of PBT were tested: 66 Gy RBE in 33 fractions and 74 Gy RBE in 37 fractions in arms 1 and 2, respectively. The planning target volume included the primary tumor and metastatic lymph nodes with adequate margins. Concurrent chemotherapy included intravenous cisplatin (60 mg/m², day 1) and oral S-1 (80, 100 or 120 mg based on body surface area, days 1-14), repeated as 4 cycles every 4 weeks. Dose-limiting toxicity (DLT) was defined as grade 3 (severe) toxicities related to PBT during days 1-90. Each dose level was performed in 3 patients, and then escalated to the next level if no DLT occurred. When 1 patient developed a DLT, 3 additional patients were enrolled. Overall, 9 patients were enrolled, including 6 in Arm 1 and 3 in Arm 2. The median follow-up time was 43 months, and the median PFS was 15 months. In Arm 1, grade 3 infection occurred in 1 of 6 patients, but no other DLT was reported. Similarly, no DLT occurred in Arm 2. However, one patient in Arm 2 developed grade 3 esophageal fistula at 9 months after the initiation of PBT. From a clinical perspective, the authors concluded that 66 Gy RBE is the recommended dose.

Oshiro et al. (2014) initiated a phase II study to evaluate the safety and efficacy of high-dose PBT with concurrent chemotherapy for unresectable or medically inoperable advanced NSCLC. Patients (n=15) were treated with PBT and chemotherapy with monthly cisplatin (on Day 1) and vinorelbine (on Days 1 and 8). The treatment doses were 74 Gy RBE for the primary site and 66 Gy RBE for the lymph nodes without elective lymph nodes. The median follow-up period was 21.7 months. None of the patients experienced Grade 4 or 5 non-hematologic toxicities. Acute pneumonitis was observed in 3 patients (Grade 1 in one, and Grade 3 in two), but Grade 3 pneumonitis was considered to be non-proton-related. Grade 3 acute esophagitis and dermatitis were observed in 1 and 2 patients, respectively. Severe (\geq Grade 3) leukocytopenia, neutropenia and thrombocytopenia were observed in 10, 7, and 1 patients, respectively. Late RP (grades 2 and 3) was observed in one patient each. Six patients (40%) experienced local recurrence at the primary site and were treated with 74 Gy RBE. Disease progression was observed in 11 patients, with the mean survival time being 26.7 months. The authors cited short follow up period as a limitation to this study. They concluded that high-dose PBT with concurrent chemotherapy is safe and useful in the multimodality therapy for unresectable NSCLC.

Sejpal et al. (2011) conducted a single-center, retrospective case series study to evaluate the use of PBT plus concurrent chemotherapy in patients with SNCLC. Outcomes included acute and subacute toxicity and were evaluated using Common Terminology Criteria (version 3.0) at least weekly during treatment, at 4-6 weeks after treatment, every 3 months for 2 years and then, every 6 months. Survival, time to progression and failure patterns were also collected. Comparisons between other radiation treatment modalities (IMRT and 3D-CRT, each with concurrent chemotherapy) were made using historical controls from the same center. A total of 202 patients were included in the analysis: 74 received 3D-CRT, 66 IMRT and 62 PBT. Median follow-up periods were 17.9 months (3D-CRT), 17.4 months (IMRT) and 15.2 months (proton). Median total radiation dose was

higher in the PBT group at 74 Gy versus 63 Gy for the other groups. Despite the higher radiation dose in the PBT group, rates of severe (grade ≥ 3) pneumonitis and esophagitis were lower (2% and 5%, respectively) compared with the other groups (3D-CRT, 30% and 18%; IMRT, 9% and 44%, respectively). Due to the short follow-up periods, tumor control and survival were not reported. The authors concluded that in this early and promising study, higher doses of PBT could be delivered to lung tumors with a lower risk of esophagitis and pneumonitis, and that additional clinical trials may further clarify the benefits and risks of PBT in patients diagnosed with NSCLC.

Pijls-Johannesma et al. (2010) conducted a systematic review to test the theory that RT with beams of protons and heavier charged particles (e.g., carbon ions) leads to superior results, compared with photon beams. The authors searched for clinical evidence to justify implementation of particle therapy as standard treatment in lung cancer. Eleven studies, all dealing with NSCLC, mainly stage I, were identified. No phase III trials were found. For PBT, 2- to 5-year LC rates varied in the range of 57%-87%. The 2- and 5-year OS and 2- and 5-year cause-specific survival rates were 31%-74% and 23% and 58%-86% and 46%, respectively. RP was observed in about 10% of patients. For CIT, the overall LC rate was 77%, but it was 95% when using a hypofractionated radiation schedule. The 5-year OS and cause-specific survival rates were 42% and 60%, respectively. Slightly better results (at 50% and 76%, respectively) were reported when using hypofractionation. The results with protons and heavier charged particles are promising. However, the current lack of evidence on the clinical effectiveness of particle therapy emphasizes the need to further investigate the efficiency of particle therapy. The authors concluded that until these results are available for lung cancer, CPT should be considered experimental.

NCCN guidelines state that advanced technologies such as PBT are appropriate when needed to deliver curative RT safely when treating NSCLC. NCCN is silent on the use of PBT in the treatment of small cell lung cancer (2021).

A phase III RCT comparing photon to proton chemoradiotherapy for patients with inoperable NSCLC (NCT01993810) is in progress. For more information, go to www.clinicaltrials.gov. (Accessed February 17, 2021)

Clinical Practice Guidelines

American College of Radiology (ACR)

ACR appropriateness criteria addressing nonsurgical treatment for locally advanced NSCLC states that while PBT may have the potential to spare critical normal tissues, more prospective studies are needed (Chang, et al., 2014).

Lymphomas

Multiple small, lower quality studies have been published on the management of lymphomas with PBT, particularly focused on long term radiation toxicity (König et al., 2019; Horn et al., 2016; Sachsman et al., 2015; Hoppe et al., 2012). Early outcomes are encouraging, but larger prospective studies are needed to confirm long term efficacy.

NCCN guidelines for Hodgkin, B-cell, and T-cell lymphomas state that PBT may be appropriate, depending on clinical circumstances. It also states that advanced RT technologies, such as PBT, may offer significant and clinically relevant advantages in specific instances to spare important organs at risk and decrease the risk for late, normal tissue damage while still achieving the primary goal of LC. NCCN is silent on the use of PBT in the treatment of primary cutaneous lymphoma (2020-2021).

Pancreatic Cancer

There is a lack of robust clinical evidence evaluating PBT for treating pancreatic cancer although research continues (Kim et al., 2018, Hong et al., 2014; Terashima et al., 2012; Hong et al., 2011). Further larger scaled prospective studies are warranted to determine the long-term safety and efficacy of this treatment modality.

NCCN guidelines do not address PBT in the treatment of pancreatic adenocarcinoma (2021).

Numerous clinical trials are currently in progress studying the use of PBT in multiple types of GI cancer (e.g., esophageal, pancreatic, and retroperitoneal sarcoma). For more information, go to www.clinicaltrials.gov. (Accessed February 16, 2021)

Vestibular Tumors

In a critical review, Murphy and Suh (2011) summarized the radiotherapeutic options for treating vestibular schwannomas, including single-session stereotactic radiosurgery, fractionated conventional RT, fractionated stereotactic RT and PBT. The comparisons of the various modalities have been based on single-institution experiences, which have shown excellent tumor control rates of 91-100%. Early experience using PBT for treating vestibular schwannomas demonstrated LC rates of 84-100% but disappointing hearing preservation rates of 33-42%. The authors report that mixed data regarding the ideal hearing preservation therapy, inherent biases in patient selection and differences in outcome analysis have made comparison across radiotherapeutic modalities difficult.

The efficacy of PBT for the treatment of tumors of the vestibular system was assessed in 2 prospective uncontrolled studies involving 30 patients with acoustic neuromas (Bush et al., 2002) and 68 patients with vestibular schwannomas (Harsh et al., 2002). Fractionated PBT effectively controlled tumor growth in all patients with acoustic neuroma, and 37.5% of patients experienced tumor regression. Hearing was preserved in 31% of patients. The actuarial 5-year tumor control rate for patients with vestibular schwannomas was 84%; 54.7% of tumors regressed, 39.1% remained unchanged, and 3 tumors enlarged. The procedure caused some serious side effects in patients with vestibular schwannoma (severe facial weakness), but most side effects were either transient or could be successfully treated.

Combined Therapies

No evidence was identified in the clinical literature supporting the combined use of PBT and IMRT in a single treatment plan.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Radiation therapy is a procedure and, therefore, is not subject to FDA regulation. However, the accelerators and other equipment used to generate and deliver PBRT are regulated by the FDA. See the following website for more information (use product code LHN): <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed February 17, 2021)

References

- American Academy of Ophthalmology. Preferred Practice Pattern® Guidelines. Age-related macular degeneration. October 2019. (Accessed February 16, 2021)
- Barney CL, Brown AP, Grosshans DR, et al. Technique, outcomes, and acute toxicities in adults treated with proton beam craniospinal irradiation. *Neuro Oncol*. 2014 Jan;16(2):303-9.
- Bekkering GE, Rutjes AW, Vlassov VV, et al. The effectiveness and safety of proton radiation therapy for indications of the eye: a systematic review. *Strahlenther Onkol*. 2009 Apr;185(4):211-21.
- Bradley JA, Dagan R, Ho MW, et al. Initial report of a prospective dosimetric and clinical feasibility trial demonstrates the potential of protons to increase the therapeutic ratio in breast cancer compared with photons. *Int J Radiat Oncol Biol Phys*. 2016 May 1;95(1):411-21.
- Bryant C, Smith TL, Henderson RH, et al. Five-year biochemical results, toxicity, and patient-reported quality of life after delivery of dose-escalated image guided proton therapy for prostate cancer. *Int J Radiat Oncol Biol Phys*. 2016 May 1;95(1):422-34.
- Bush DA, Do S, Lum S, et al. Partial breast radiation therapy with proton beam: 5-year results with cosmetic outcomes. *Int J Radiat Oncol Biol Phys*. 2014 Nov 1;90(3):501-5.
- Bush DA, McAllister CJ, Loredon LN, et al. Fractionated proton beam radiotherapy for acoustic neuroma. *Neurosurgery*. 2002;50(2):270-275.
- Bush DA, Smith JC, Slater JD, et al. Randomized clinical trial comparing proton beam radiation therapy with transarterial chemoembolization for hepatocellular carcinoma: results of an interim analysis. *Int J Radiat Oncol Biol Phys*. 2016 May 1;95(1):477-82.
- Chan RV, Yonekawa Y, Lane AM, et al. Proton beam irradiation using a light-field technique for the treatment of choroidal hemangiomas. *Ophthalmologica*. 2010;224(4):209-16.

Chang JY, Verma V, Li M, et al. Proton beam radiotherapy and concurrent chemotherapy for unresectable stage iii non-small cell lung cancer: final results of a phase 2 study. *JAMA Oncol.* 2017 Aug 10; 3(8):e172032.

Chang JY, Kestin LL, Barriger RB, et al. ACR Appropriateness Criteria® nonsurgical treatment for locally advanced non-small-cell lung cancer: good performance status/definitive intent [online publication]. Reston (VA): American College of Radiology (ACR); 2014.

Chi A, Chen H, Wen S, et al. Comparison of particle beam therapy and stereotactic body radiotherapy for early stage non-small cell lung cancer: A systematic review and hypothesis-generating meta-analysis. *Radiother Oncol.* 2017 Jun;123(3):346-354.

Cuaron JJ, Chon B, Tsai H, et al. Early toxicity in patients treated with postoperative proton therapy for locally advanced breast cancer. *Int J Radiat Oncol Biol Phys.* 2015 Jun 1;92(2):284-91.

DeCesaris CM, Rice SR, Bentzen SM, et al. Quantification of acute skin toxicities in patients with breast cancer undergoing adjuvant proton versus photon radiation therapy: a single institutional experience. *Int J Radiat Oncol Biol Phys.* 2019 Aug 1;104(5):1084-1090.

Evans JR, Igwe C, Jackson TL, et al. Radiotherapy for neovascular age-related macular degeneration. *Cochrane Database Syst Rev.* 2020 Aug 26;8:CD004004.

Evans JR, Sivagnanavel V, Chong V. Radiotherapy for neovascular age-related macular degeneration. *Cochrane Database Syst Rev.* 2010 May 12;5:CD004004.

Frau E, Rumen F, Noel G, et al. Low-dose proton beam therapy for circumscribed choroidal hemangiomas. *Arch Ophthalmol.* 2004 Oct;122(10):1471-5.

Harada H, Fuji H, Ono A, et al. Dose escalation study of proton beam therapy with concurrent chemotherapy for stage III non-small cell lung cancer. *Cancer Sci.* 2016 Jul;107(7):1018-21.

Harsh GR, Thornton AF, Chapman PH, et al. Proton beam stereotactic radiosurgery of vestibular schwannomas. *Int J Radiat Oncol Biol Phys.* 2002;54(1):35-44.

Hartsell WF, Kapur R, Hartsell SO, et al. Feasibility of proton beam therapy for ocular melanoma using a novel 3d treatment planning technique. *Int J Radiat Oncol Biol Phys.* 2016 May 1;95(1):353-9.

Hattangadi JA, Chapman PH, Bussi re MR, et al. Planned two-fraction proton beam stereotactic radiosurgery for high-risk inoperable cerebral arteriovenous malformations. *Int J Radiat Oncol Biol Phys.* 2012 Jun 1;83(2):533-41.

Hattangadi-Gluth JA, Chapman PH, Kim D, et al. Single-fraction proton beam stereotactic radiosurgery for cerebral arteriovenous malformations. *Int J Radiat Oncol Biol Phys.* 2014 Jun 1;89(2):338-46.

Hayes, Inc. Hayes Evidence Analysis Research Brief. Proton Beam Therapy for Head and Neck Cancer. Lansdale, PA: Hayes, Inc. October 2019.

Hayes, Inc. Hayes Health Technology Assessment. Proton Beam Therapy for Non-Small Cell Lung Cancer. Lansdale, PA: Hayes, Inc. March 2020.

Holliday EB, Frank SJ. Proton radiation therapy for head and neck cancer: a review of the clinical experience to date. *Int J Radiat Oncol Biol Phys.* 2014 Jun 1;89(2):292-302.

Hong TS, Ryan DP, Blaszkowsky LS, et al. Phase I study of preoperative short-course chemoradiation with proton beam therapy and capecitabine for resectable pancreatic ductal adenocarcinoma of the head. *Int J Radiat Oncol Biol Phys.* 2011 Jan 1;79(1):151-7.

Hong TS, Ryan DP, Borger DR, et al. A phase 1/2 and biomarker study of preoperative short course chemoradiation with proton beam therapy and capecitabine followed by early surgery for resectable pancreatic ductal adenocarcinoma. *Int J Radiat Oncol Biol Phys.* 2014 Jul 15;89(4):830-8.

Hoppe BS, Flampouri S, Lynch J, et al. Improving the therapeutic ratio in hodgkin lymphoma through the use of proton therapy. *Oncology (Williston Park).* 2012 May;26(5):456-9, 462-5.

Horn S, Fournier-Bidoz N, Pernin V, et al. Comparison of passive-beam proton therapy, helical tomotherapy and 3D conformal radiation therapy in Hodgkin's lymphoma female patients receiving involved-field or involved site radiation therapy. *Cancer Radiother.* 2016 Apr;20(2):98-103.

Kabolizadeh P, Chen YL, Liebsch N, et al. Updated outcome and analysis of tumor response in mobile spine and sacral chordoma treated with definitive high-dose photon/proton radiation therapy. *Int J Radiat Oncol Biol Phys.* 2017 Feb 1;97(2):254-262.

Kagei K, Tokuyue K, Okumura T, et al. Long-term results of proton beam therapy for carcinoma of the uterine cervix. *Int J Radiat Oncol Biol Phys.* 2003;55(5):1265-1271.

Kim TH, Lee WJ, Woo SM, et al. Effectiveness and safety of simultaneous integrated boost-proton beam therapy for localized pancreatic cancer. *Technol Cancer Res Treat.* 2018 Jan 1;17:1533033818783879.

König L, Bougatf N, Hörner-Rieber J, et al. Consolidative mediastinal irradiation of malignant lymphoma using active scanning proton beams: clinical outcome and dosimetric comparison. *Strahlenther Onkol.* 2019 Jul;195(7):677-687.

Liao Z, Lee JJ, Komaki R, et al. Bayesian adaptive randomization trial of passive scattering proton therapy and intensity-modulated photon radiotherapy for locally advanced non-small-cell lung cancer. *J Clin Oncol.* 2018 Jan 2;JCO2017740720.

Lin SH, Merrell KW, Shen J, et al. Multi-institutional analysis of radiation modality use and postoperative outcomes of neoadjuvant chemoradiation for esophageal cancer. *Radiother Oncol.* 2017 Jun;123(3):376-381.

Miyayama N, Akaza H, Okumura T, et al. A bladder preservation regimen using intra-arterial chemotherapy and radiotherapy for invasive bladder cancer: a prospective study. *Int J Urol.* 2000;7(2):41-48.

Mizumoto M, Sugahara S, Nakayama H, et al. Clinical results of proton-beam therapy for locoregionally advanced esophageal cancer. *Strahlenther Onkol.* 2010 Sep;186(9):482-8.

Mizumoto M, Sugahara S, Okumura T, et al. Hyperfractionated concomitant boost proton beam therapy for esophageal carcinoma. *Int J Radiat Oncol Biol Phys.* 2011 Nov 15;81(4):e601-6.

Murphy ES, Suh JH. Radiotherapy for vestibular schwannomas: a critical review. *Int J Radiat Oncol Biol Phys.* 2011 Mar 15;79(4):985-97.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. B-cell Lymphoma. V1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Bladder Cancer. V6.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Bone Cancer. v1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Breast Cancer. V6.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Central Nervous System Cancers. V3.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Cervical Cancer. V1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Esophageal and Esophagogastric Junction Cancers. V4.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Gastric Cancer. v2.2019.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Head and Neck Cancers. v3.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Hodgkin Lymphoma. v2.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Non-Small Cell Lung Cancer. V8.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer. V1.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Pancreatic Adenocarcinoma. V1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Primary Cutaneous B-cell Lymphoma. V1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Small Cell Lung Cancer. V1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. T-cell Lymphoma. V1.2021.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Uterine Neoplasms. V1.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Uveal Melanoma. V2.2020.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Vulvar Cancer. v2.2021.

Noel G, Habrand JL, Mammar H, et al. Highly conformal therapy using proton component in the management of meningiomas. Preliminary experience of the Centre de Protontherapie d'Orsay. *Strahlenther Onkol.* 2002 Sep;178(9):480-5.

Oshiro Y, Okumura T, Kurishima K, et al. High-dose concurrent chemo-proton therapy for Stage III NSCLC: preliminary results of a Phase II study. *J Radiat Res.* 2014 Sep;55(5):959-65.

Petr J, Platzek I, Hofheinz F, et al. Photon vs. proton radiochemotherapy: Effects on brain tissue volume and perfusion. *Radiother Oncol.* 2018 Jul;128(1):121-127.

Pijls-Johannesma M, Grutters JP, Verhaegen F, et al. Do we have enough evidence to implement particle therapy as standard treatment in lung cancer? A systematic literature review. *Oncologist.* 2010;15(1):93-103.

Sachsman S, Flampouri S, Li Z, et al. Proton therapy in the management of non-Hodgkin lymphoma. *Leuk Lymphoma.* 2015;56(9):2608-12.

Sanda MG, Chen RC, Crispino T, et al. Clinically localized prostate cancer: AUA/ASTRO/SUO Guideline. Presentations from the 2017 AUA Annual Meeting. [http://www.auanet.org/guidelines/clinically-localized-prostate-cancer-new-\(aua/astro/suo-guideline-2017\)](http://www.auanet.org/guidelines/clinically-localized-prostate-cancer-new-(aua/astro/suo-guideline-2017)). Accessed February 17, 2021.

Sejpal S, Komaki R, Tsao A, et al. Early findings on toxicity of proton beam therapy with concurrent chemotherapy for non-small cell lung cancer. *Cancer.* 2011 Jul 1;117(13):3004-13.

Takaoka EI, Miyazaki J, Ishikawa H, et al. Long-term single-institute experience with trimodal bladder-preserving therapy with proton beam therapy for muscle-invasive bladder cancer. *Jpn J Clin Oncol.* 2017 Jan;47(1):67-73.

Terashima K, Demizu Y, Hashimoto N, et al. A phase I/II study of gemcitabine-concurrent proton radiotherapy for locally advanced pancreatic cancer without distant metastasis. *Radiother Oncol.* 2012 Apr;103(1):25-31.

Verma V, Iftekaruddin Z, Badar N, et al. Proton beam radiotherapy as part of comprehensive regional nodal irradiation for locally advanced breast cancer. *Radiother Oncol.* 2017 May;123(2):294-298.

Verma V, Lin SH, Simone CB 2nd, et al. Clinical outcomes and toxicities of proton radiotherapy for gastrointestinal neoplasms: a systematic review. *J Gastrointest Oncol.* 2016b Aug;7(4):644-64.

Verma V, Shah C, Mehta MP. Clinical outcomes and toxicity of proton radiotherapy for breast cancer. *Clin Breast Cancer.* 2016a Jun;16(3):145-54.

Wang J, Wei C, Tucker SL, et al. Predictors of postoperative complications after trimodality therapy for esophageal cancer. *Int J Radiat Oncol Biol Phys.* 2013a Aug 1;86(5):885-91.

Xi M, Xu C, Liao Z, et al. Comparative outcomes after definitive chemoradiotherapy using proton beam therapy versus intensity modulated radiation therapy for esophageal cancer: a retrospective, single-institutional analysis. *Int J Radiat Oncol Biol Phys.* 2017 Nov 1;99(3):667-676.

Zambarakji HJ, Lane AM, Ezra E, et al. Proton beam irradiation for neovascular age-related macular degeneration. *Ophthalmology.* 2006;113(11):2012-9.

Zenda S, Akimoto T, Mizumoto M, et al. Phase II study of proton beam therapy as a nonsurgical approach for mucosal melanoma of the nasal cavity or para-nasal sinuses. *Radiother Oncol.* 2016 Feb;118(2):267-71.

Policy History/Revision Information

Date	Summary of Changes
07/01/2021	<p data-bbox="337 237 592 268">Coverage Rationale</p> <ul data-bbox="337 275 1096 306" style="list-style-type: none"><li data-bbox="337 275 1096 306">• Replaced reference to “InterQual® 2020” with “InterQual® 2021” <p data-bbox="337 312 568 344">Applicable Codes</p> <ul data-bbox="337 350 1177 382" style="list-style-type: none"><li data-bbox="337 350 1177 382">• Added ICD-10 diagnosis codes C11.1, C11.2, C11.3, C11.8, and C11.9 <p data-bbox="337 388 641 420">Supporting Information</p> <ul data-bbox="337 426 1396 510" style="list-style-type: none"><li data-bbox="337 426 1396 457">• Updated <i>Clinical Evidence</i> and <i>References</i> sections to reflect the most current information<li data-bbox="337 457 641 489">• Removed <i>CMS</i> section<li data-bbox="337 489 885 520">• Archived previous policy version CS105IN.01

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.