

Minimally Invasive Procedures for Gastric and Esophageal Diseases (for Kentucky Only)

Policy Number: CS079KY.11

Effective Date: July 1, 2025

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	2
Applicable Codes	2
Description of Services	2
Clinical Evidence	4
U.S. Food and Drug Administration	21
References	22
Policy History/Revision Information	27
Instructions for Use	27

Related Policy

- [Bariatric Surgery \(for Kentucky Only\)](#)

Application

This Medical Policy only applies to the state of Kentucky.

Coverage Rationale

Note: This policy only applies to individuals 18 years of age and older.

For medical necessity clinical coverage criteria for minimally invasive procedures for GERD and Achalasia, refer to the InterQual® CP: Procedures, Antireflux Surgery or Hiatal Hernia Repair.

[Click here to view the InterQual® criteria.](#)

The per oral endoscopic myotomy (POEM) procedure is proven and medically necessary for treating individuals with [Achalasia](#) or [Diffuse Esophageal Spasm](#).

Per oral endoscopic myotomy (POEM) is unproven and not medically necessary for all other indications (e.g., Zenker's diverticula) due to insufficient evidence of efficacy.

Gastric per oral endoscopic myotomy (G-POEM) is unproven and not medically necessary for the treatment of [Gastroparesis](#).

The following are unproven and not medically necessary for treating [Gastroesophageal Reflux Disease](#) (GERD) due to insufficient evidence of efficacy:

- [Endoscopic therapies](#) other than Transoral Incisionless Fundoplication
- [Injection or implantation techniques](#)
- [LINX Reflux Management System](#)

Functional lumen imaging probe technology is unproven and not medically necessary for diagnosing Achalasia.

Endoluminal therapy with GERDx™ is investigational, unproven, and not medically necessary for treating GERD as it has not received U.S. Food and Drug Administration (FDA) approval.

Refer to the Medical Policy titled [Bariatric Surgery \(for Kentucky Only\)](#) for information regarding endoscopic therapies for the treatment of obesity.

Definitions

Achalasia: A primary esophageal motor disorder of unknown etiology characterized by degeneration of the myenteric plexus, which results in impaired relaxation of the esophagogastric junction (EGJ), along with the loss of organized peristalsis in the esophageal body [American Society of Gastrointestinal Endoscopy (ASGE)].

Diffuse Esophageal Spasm: Rare esophageal motility disorder characterized by simultaneous, uncoordinated, or rapidly propagated contractions that are of normal amplitude and accompanied by dysphagia (National Library of Medicine).

Gastroesophageal Reflux Disease: A condition where the lower esophageal sphincter (LES) relaxes too often or weakens which allows stomach acid to flow backward (or reflux) into the esophagus [American College of Gastroenterology (ACG)].

Gastroparesis: A chronic disorder which means delayed stomach emptying without a blockage (ACG).

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
43210	Esophagogastroduodenoscopy, flexible, transoral; with esophagogastric fundoplasty, partial or complete, includes duodenoscopy when performed
43257	Esophagogastroduodenoscopy, flexible, transoral; with delivery of thermal energy to the muscle of lower esophageal sphincter and/or gastric cardia, for treatment of gastroesophageal reflux disease
43284	Laparoscopy, surgical, esophageal sphincter augmentation procedure, placement of sphincter augmentation device (i.e., magnetic band), including cruroplasty when performed
43289	Unlisted laparoscopy procedure, esophagus
43497	Lower esophageal myotomy, transoral (i.e., peroral endoscopic myotomy [POEM])
43499	Unlisted procedure, esophagus
43999	Unlisted procedure, stomach

CPT® is a registered trademark of the American Medical Association

Description of Services

Gastroesophageal Reflux Disease (GERD) is a condition that is characterized by either a weak or dysfunctional lower esophageal sphincter (LES) that results in partially digested food from the stomach to flow back into the esophagus, a process known as reflux. Persistent GERD may lead to esophageal damage or other serious conditions, such as severe esophagitis, strictures, Barrett's metaplasia, and adenocarcinoma of the esophagus.

Initial treatment of GERD usually involves over the counter (OTC) antacids, OTC histamine-2-receptor antagonists (H₂RAs; also called H₂ blockers), and proton pump inhibitors (PPI). Daily use of proton pump inhibitors (PPI) is generally effective in the treatment of most patients with GERD; however, up to 40% have persisting symptoms (Weitzendorfer et al., 2018). For individuals who wish to discontinue use of these medications due to concern of long-term side effects, or for individuals whose GERD is refractory to pharmacologic treatment, an open or laparoscopic Nissen fundoplication may be considered. However, some individuals may not be suitable candidates given the invasiveness and risks associated with surgery. As a result, minimally invasive procedures, including endoscopic or endoluminal therapies and laparoscopic approaches, have been proposed as alternative treatment methods to improve the function of the LES, with the objective of eliminating symptoms, healing esophagitis, preventing recurrence of symptoms or progression of disease, and reducing the need for lifelong pharmacologic therapy.

Minimally invasive approaches proposed in the treatment of GERD, include the following:

- Radiofrequency energy: The Stretta procedure administers radiofrequency (RF) energy via endoscopic needles placed in the tissues surrounding the lower esophageal sphincter. The RF energy heats this neighboring tissue, creating thermal lesions. Submucosal scarring forms as the lesions heal, causing shrinkage and tightening around the LES.
- Endoscopic plication or suturing:
 - The NDO Endoscopic Plication System, also known as the NDO Plicator System, places a full-thickness transmural plication near the gastroesophageal junction under direct endoscopic visualization.
 - EsophyX is an endoluminal therapeutic option that uses a trans-oral and fastener deploying device. It is inserted orally within a thin, flexible tube and deployed inside the stomach to create a full thickness plication of the stomach fundus at the GE junction, thereby resembling an endoscopic fundoplication. The current TIF 2.0 technique (the initial TIF 1.0 technique is no longer recommended) generates a physiological valve via fasteners placed on the far posterior and anterior sides of the lesser curvature, with additional fasteners placed 1-3 cm proximal to the GE junction (Hayes, 2023).
 - GERDx™ (G-SURG) is an endoscopic full-thickness plication device that uses hydraulic elements for controlling.
 - The Medigus Ultrasonic Surgical Endostapler (MUSE™ system, Medigus) is an endoscopic stapling device for transoral partial fundoplication. According to the manufacturer's website, as the MUSE system contains the surgical stapler, microvisual, and ultrasonic capabilities, it allows a single physician to complete the procedure.
- Injection or implantation techniques include the following:
 - The Plexiglas® [polymethylmethacrylate (PMMA)] procedure involves injection of an inert polymer material into the submucosa of the proximal LES zone to provide bulking support to the sphincter and decrease transient relaxation of the lower esophageal sphincter (tLESRs).
 - Another bulking agent, pyrolytic carbon-coated beads (Durasphere®), is being evaluated for treatment of GERD.
 - The LINX™ Reflux Management System is an implant that consists of a ring that fits around the esophagus and is intended to prevent reflux of bile and acid from the stomach into the esophagus. According to the company website, the LINX system is a small flexible band of interlinked titanium beads with magnetic cores. The magnetic attraction is intended to help the LES resist opening to gastric pressures, preventing reflux from the stomach into the esophagus. A surgeon uses a laparoscopic incision to implant the device around the patient's esophagus just above the stomach while the patient is under general anesthesia.

Achalasia is a condition that affects the esophagus. It is a relatively rare cause of dysphagia manifested by esophageal aperistalsis and failure of relaxation of a hypertensive lower esophageal sphincter (LES) (Kohn 2019). Current treatment options include pharmacological, endoscopic, and surgical.

- The per oral endoscopic myotomy (POEM) procedure is a technique that involves guiding an endoscope through the esophagus, making an incision in the mucosa, creating a submucosal tunnel for access to the lower esophagus and gastroesophageal junction, and cutting the muscle fibers in the lower esophagus and proximal stomach. Internal incisions are closed with clips after myotomy is complete. POEM is an intricate endoscopic procedure that requires advanced endoscopic skills, knowledge of surgical anatomy, and expertise in submucosal endoscopy and management of adverse events (Khasab et al., 2020).

Diffuse Esophageal Spasm (also known as distal esophageal spasm) is a condition that leads to premature and rapidly produced contractions in the distal esophagus. Most patients present with difficulty swallowing and often have a sensation of foods stuck in their esophagus. Distal esophageal spasm is distinguished from other esophageal motility disorders that are associated with dysphagia by esophageal manometry testing.

Endoluminal functional lumen imaging probe (FLIP) technology is used for esophageal function testing, such as testing for Achalasia or Diffuse Esophageal Spasm. FLIP is performed alone or in conjunction with other diagnostic tests; EndoFlip™ is one such device. A catheter surrounded by a balloon filled with liquid extends from the esophagus into the stomach or from the stomach into the small intestine. The device then applies pressure while providing continuous diameter measurements along its length, giving information about esophageal and/or gastric function (Hayes, 2022).

Gastroparesis is a chronic condition which delays gastric emptying in the absence of an obstruction. Individuals often experience abdominal pain including nausea, vomiting, and bloating. It is more common in women and individuals with diabetes. Treatment options include dietary modifications, pharmacologic therapy, medication changes, and surgical treatments such as pyloroplasty.

Per Oral Endoscopic Myotomy (POEM)

In a health technology assessment by Hayes (2023), POEM has a “potential but unproven benefit” as an alternative to laparoscopic Heller myotomy in patients with esophageal achalasia. The authors of the report conclude that the available low-quality evidence suggested the POEM procedure is generally safe and may achieve at least similar results to both laparoscopic Heller myotomy (LHM) and pneumatic dilation (PD) for most efficacy outcomes. The body of evidence on POEM vs LHM was of moderate size including 16 studies, whereas evidence on POEM versus PD was presented in only four studies. It is suggested additional studies of fair to good quality are needed to reveal optimal treatment protocols and provide information for longer-term outcomes.

North and Tewari (2024) conducted a systematic review in accordance with PRISMA guidelines, incorporating the Werner study cited below, to compare per oral endoscopic myotomy (POEM) to laparoscopic Heller myotomy (LHM) and pneumatic dilation (PD) for treating esophageal achalasia (EA). The review included 31 studies and the authors found that POEM significantly improves achalasia symptoms over an extended period, particularly benefitting type III achalasia patients. Additionally, they noted POEM also shows a lower recurrence rate requiring retreatment compared to PD, suggesting it as a safer and more effective long-term treatment option. When compared to LHM, the authors stated POEM demonstrates similar patient outcomes in symptom relief and recurrence, albeit with a higher incidence of gastro-esophageal reflux disease (GORD). However, this can be effectively controlled with medical management over the long term. The authors concluded overall; the findings support POEM as a viable first-line treatment for achalasia.

Saleh et al. (2023) evaluated the efficacy of POEM versus pneumatic dilation (PD) for patients with persistent or recurrent symptoms following a laparoscopic Heller myotomy (LHM). Ninety participants were recruited; 45 participants were randomly assigned to receive POEM and the other 45 were assigned to receive PD. Inclusion criteria consisted of patients aged 18-80 years with persistent or recurrent symptoms after LHM and an Eckardt symptom score > 3. For the PD group a series of dilations with Rigidflex balloons (Boston Scientific) was performed where the participant reviewed at least two dilations, with the last one occurring at a minimum of at least 35 mm. Those receiving POEM underwent general anesthesia. The primary outcome was treatment success after one-year follow-up, which was defined as an Eckardt score of ≤ 3 and no unscheduled re-treatment. Secondary outcomes included scores from the 36-Item Short Form Health Survey (SF-36) and the achalasia disease-specific quality of life questionnaire (ADSQoL). Symptoms and scores were assessed at three months and again at one year. The authors found the participants receiving POEM had greater treatment success at one-year than the patients treated with PD. The Eckardt score was found to be lower in the patients treated with POEM vs those treated with PD. At one-year follow-up, the authors saw a higher incidence of reflux esophagitis in the POEM group, but only 8.3% were assigned a grade C whereas 16.7% had grade C from the PD group. Serious adverse events were minimal and included microp perforation occurring in one participant after POEM and the other consisted of chronic severe reflux symptoms after PD; both were treated and the patients continued with the study. The author's concluded achalasia patients treated with POEM for persistent or recurrent symptoms after LHM resulted in a higher success rate than those treated with PD. Limitations included lack of long-term outcomes and lack of blinding.

In a single center randomized controlled trial (RCT), de Moura et al. (2022) compared POEM to laparoscopic myotomy and partial fundoplication (LM-PF) for efficacy and outcomes in the treatment of achalasia. Forty participants were randomized to undergo either POEM or LM-PF. Inclusion criteria consisted of patients ≥ 18 years of age with an achalasia diagnosis, dysphagia score ≥ 11 and Eckardt score > 3. Patients were followed up at one-, six- and 12 months which included Eckardt scores, esophagogastroduodenoscopy (EDG) procedure, timed barium esophagograms and completion of the Medical Outcomes Study 36-item Short-Form Health Survey. Success was defined with symptom improvement (≤ 3 -point reduction in the Eckardt score), a lower esophageal sphincter (LES) pressure < 15 mmHg, and a > 50% reduction in the height of the barium column at one minute. The authors found improvements in both groups which was supported by a decrease in the Eckardt score. It was concluded that POEM and LM-PF appear to be equally effective in controlling the symptoms of achalasia, shortening length of stay (LOS), and minimizing adverse events (AEs). Limitations included a sample size that may have been too small to detect clinically significant differences between groups and absence of pHmetry evaluation, which is the main method for GERD evaluation.

Huang et al. (2021) completed a systematic review and meta-analysis to evaluate the safety and efficacy of POEM in patients with achalasia and a previous Heller myotomy (HM). A search was conducted using PubMed, Embase, and the Cochrane Library. A total of nine observational studies involving 272 patients were found. Primary outcomes included clinical success as defined by pre- and post-op Eckardt scores, lower esophageal sphincter (LES) pressure and integrated relaxation pressure (IRP) scores; secondary outcome included safety assessment as defined by adverse events and incidence of postop GERD. All nine studies reported a significant reduction in the Eckardt score by 5.14 (95% CI, 4.19-6.09), with significant heterogeneity. Clinical success was achieved in 90% of the patients. LES pressure and IRP

were significantly lowered by 12.01 mm Hg and 10.02 mm Hg, respectively. Adverse events (AEs) were reported in six studies with mucosal injury as the most common, and this occurred in 11 patients. Based on the analysis, the authors concluded that POEM is a safe and effective treatment for patients with achalasia; this was supported by the favorable Eckardt scores and manometry parameters. Limitations included lack of comparison to other approaches, non-randomization, considerable heterogeneity across all outcome measures, and short-term follow-up. Additional prospective, controlled studies with long-term follow-up are warranted to confirm these findings.

Chandan et al. (2020) performed a systematic review and meta-analysis to evaluate the efficacy of POEM in patients with spastic esophageal disorders (SED) and if variation in total myotomy length or prior endoscopic treatment had any impact on the clinical success. A comprehensive literature search in PubMed, EMBASE, Google-Scholar, Scopus, and Cochrane Review retrieved nine studies which included 210 patients; of the nine studies, three studies were prospective and the other six were retrospective. Several outcomes were assessed and clinical success was defined as achieving an Eckardt score ≤ 3 post-intervention. The overall clinical success rate was documented at 89.6% with low heterogeneity. Symptomatic reflux was also analyzed and all but one study reported that patients with reflux responded with proton pump inhibitor therapy. Follow-up periods ranged from 2.7 months to 27 months. Other adverse events included chest/epigastric pain that required hospitalization, esophageal leak, pneumothorax and post-op pain. The authors concluded that, while POEM is safe and effective for SED, total myotomy length and prior endoscopic or medical treatments had no effect on its clinical success. Limitations included lack of comparison to another approach, retrospective design, and all studies were performed in tertiary-care centers thereby not giving a true representation of the general population. (Khashab [2018] is included in this systematic review).

In a prospective multicenter randomized open label trial, Werner et al. (2019) compared POEM to laparoscopic Heller's myotomy (LHM) plus fundoplication in 221 patients with achalasia using a design to demonstrate non-inferiority. The patients were randomly assigned in a 1:1 ratio to undergo either POEM or LHM plus fundoplication. The POEM procedure was performed by a physician with formal POEM training including esophageal interventions such as endoscopic mucosal resection and submucosal dissection; LHM was performed according to current standards. Clinical data was collected at three, six, 12, and 24 months; patient assessment was performed with phone calls, mail, and follow-up appointments. The Eckardt symptom score was the validated questionnaire used which identified success with a score of three or less by the two-year follow up appointment. Clinical success at the two-years' follow-up was observed in 83.0% of patients in the POEM group and 81.7% of patients in the LHM group (difference, 1.4 percentage points; 95% confidence interval [CI], -8.7 to 11.4; $p = 0.007$ for noninferiority). Limitations included lack of obtaining appropriate consent from patients, lack of blinding, and surgeon experience was superior for HLM versus POEM. The author's concluded POEM was non-inferior to LHM in controlling symptoms of achalasia at two years with less adverse events; it was noted the patients with the POEM procedure were more common to experience gastroesophageal reflux than the patients who underwent HLM.

In a systematic review, which did not include the Werner study cited above, Li et al. (2019) investigated the long-term efficacy and safety of POEM with follow-up period over two years. Ten eligible studies met the inclusion criteria and were published between January 2015 and November 2017. A total of 372 patients successfully underwent POEM with one failure due to serious inflammation and adhesion of the esophagus. The mean follow-up period was 30 months. The mean preoperative and postoperative Eckhart scores decreased from 7.4 to 1.4, respectively. The authors found POEM to be effective and safe for the treatment of achalasia during the two years long-term follow-up duration. It was concluded further multi-center studies with randomization comparing POEM with other treatment modalities are warranted for the future. Limitations included small sample size for most of the studies and lack of comparison to other approaches.

He et al. (2019) (not included in the systematic reviews cited above) collected prospective data in a case series of 115 patients to evaluate the long-term efficacy of POEM for patients with achalasia. The Eckardt scoring system was used and success was found in 91.3% of the patients. Twenty-one patients were found to have symptoms of reflux during the two-year follow-up. The authors concluded that POEM was safe and effective for treating achalasia with favorable long-term outcomes. The findings are limited by lack of comparison group.

In a 2018 retrospective multicenter study, Khashab et al. assessed the technical success, clinical response, and adverse events of POEM in 50 patients with non-achalasia esophageal motility disorders such as esophagogastric junction outflow obstruction (EGJO), diffuse esophageal spasm (DES), and jackhammer esophagus (JE). Patients diagnosed with achalasia were excluded. Just over half of the patients were treatment naïve. The results showed that POEM was successful in all 50 patients, and nine adverse events (AEs) were reported: 55.6% were rated as mild and 44.4% as moderate with no severe events. At the median follow up time of 242 days, 42 patients achieved clinical success and the majority had complete or almost complete resolution of symptoms. Reflux symptoms developed in 22.2% of patients, all of whom were successfully treated with proton pump inhibitors. The authors concluded that POEM is safe and effective for the management of non- achalasia esophageal motility disorders and randomized trials are needed to confirm these findings. The findings are limited by lack of comparison group.

Khan et al. (2017) conducted a systematic review and meta-analysis of the published literature regarding the efficacy and safety of per-oral endoscopic myotomy (POEM) for the treatment of all spastic esophageal disorders (SEDs). Included were ninety-eight full studies of five or more patients that reported clinical success and post procedure adverse events and included eight observational studies that included follow up ranging from three months to three years were included in the meta-analysis. Three studies were prospective and the remaining five were retrospective. The total number of patients was 179, with the following diagnoses: 116 had type III achalasia, 37 had jackhammer esophagus, 18 had diffuse esophageal spasm, and eight had hypertensive non-relaxing lower esophageal sphincter. The results showed a weighted mean pooled rate (WPR) of success of POEM for type III achalasia at 87%, jackhammer esophagus was 72%, and diffuse esophageal spasm at 88%. The WPR of success of POEM for all SEDs was 87%. All studies reported adverse events and showed a WPR of 11% for type III achalasia, 16% for jackhammer esophagus and 14% for diffuse esophageal spasm. The authors concluded that POEM is a highly effective and safe treatment modality for treating SEDs, and larger prospective studies are required to validate these results. The findings are however limited by lack of comparison group.

Marano et al. (2016) performed a systematic review and meta-analysis to investigate the efficacy and safety of POEM compared with laparoscopic Heller's myotomy (LHM) for the treatment of achalasia. The search produced 11 studies. The total number of included patients was 486 (196 in POEM group and 290 in LHM group) ranging from eight to 180 patients per study. The Eckardt score was used in five of the studies which showed non-statistically significant difference between POEM and LHM favoring POEM. The review and analysis identified some limitations including high heterogeneity rate, no randomization of patients and significant publication bias. Furthermore, all selected studies did not report follow-up results past one year. The authors concluded additional high-quality clinical trials with randomization and long-term evaluation comparing POEM with other standard procedures are needed.

Zenker's Peroral Endoscopic Myotomy (Z-POEM)

Peroral endoscopic myotomy (POEM) is a novel technique in the treatment of Zenker's diverticulum (ZD). There is insufficient quality evidence in the published clinical literature involving large well-designed studies with standardized post-procedure protocols. Long term monitoring and sufficient evidence to confirm safety and efficacy are insufficient.

Al Ghamdi et al. (2022) conducted a multi-center, international, retrospective cohort study comparing the effectiveness of flexible and rigid endoscopic septotomy with Zenker's peroral endoscopic myotomy (Z-POEM) for treating Zenker's diverticulum. The study included 245 participants from 12 centers. Primary outcomes assessed were clinical success (defined as a decrease in Dakar and Bennett dysphagia score to ≤ 1), clinical failure and clinical recurrence. Secondary outcomes included technical success and incidence/severity of adverse events. Z-POEM was the most frequently used method ($n = 119$), followed by flexible ($n = 86$) and rigid ($n = 40$) endoscopic septotomy. Baseline participant characteristics were generally balanced across the three groups. At the initial follow-up, clinical success rates were comparable among the groups; 92.7% for Z-POEM, 89.2% for rigid septotomy, and 86.7% for flexible septotomy ($p = 0.26$). Symptom recurrence was observed in 24 participants (15 Z-POEM during a mean follow-up of 282.04 [SD 300.48] days, six flexible, three rigid [$p = 0.47$]). Adverse events were more frequent in the rigid septotomy group (30.0%) compared to the Z-POEM (16.8%) and flexible septotomy (2.3%) groups ($p < 0.05$). The authors concluded that all three techniques are effective for treating symptomatic Zenker's diverticulum, although Z-POEM had a higher recurrence rate. They also noted that flexible endoscopic septotomy had a shorter procedure time, similar clinical success rate, and fewer adverse events compared to Z-POEM and rigid endoscopic septotomy. The authors recommended prospective studies with longer-term follow-up to provide more conclusive evidence regarding the outcomes of each approach. Limitations of this study included its multicenter nature, which introduced variability in techniques and follow-up protocols across centers, potential underreporting of adverse events, and relatively small sample sizes in some groups, potentially limiting the ability to detect significant differences. Yang et al. (2020) is included in this review.

Zhang et al. (2022) conducted a systematic review which evaluated the safety and efficacy of Z-POEM for Zenker's diverticulum (ZD) and compared the feasibility and effectiveness of Z-POEM with that of a flexible endoscopic septotomy (FES). A search was conducted using PubMed, EMBASE, Web of Science, and Cochrane Library databases which returned eleven studies (three prospective and eight retrospective) for analysis. Eligibility criteria for article search included individuals with symptomatic ZD, completed Z-POEM procedure, a control group, technical and clinical success rates, and adverse events. Only four studies reported before and after symptom score changes for the Z-POEM procedure. The authors found no significant differences between Z-POEM and FES when it came to procedure time, clinical recurrence, or adverse events; future high-quality comparative studies are required to further validate these findings. Limitations included the type of studies involved, mostly observational studies, potential duplication of study participants amongst studies assessed, lack of objective indicators and publication bias.

Budnicka et al. (2021, included in Zhang (2022) systematic review above) conducted a multicenter retrospective case series aimed at analyzing the feasibility of POEM for Zenker's diverticula. Twenty-two patients with various degrees of dysphagia diagnosed with symptomatic ZD were included. Primary outcomes were the rate of technical success and the

procedure's clinical success. These were defined by completion of all procedural steps and resolution of dysphagia or resolution of symptoms. POEM was successful in all 22 patients; no severe or fatal adverse outcomes were reported. Clinical success was achieved in twenty patients; two patients continued with persistent dysphagia. The authors concluded Z-POEM as a viable option for treatment in relieving dysphagia and other related symptoms. However, limitations included retrospective design, small sample size, lack of comparison group, and short-term outcomes. Additional future studies should include comparative studies with long term efficacy.

A retrospective multicenter case series on the use of POEM in the management of Zenker's diverticulum was conducted by Yang et al. (2020). A total of 75 participants from ten different international centers who underwent Z-POEM between January 2014 and November 2018 were included. Diagnoses of ZD was confirmed by endoscopy along with a dysphagia score from Dakkak and Bennett scoring system (0 = no dysphagia; 1 = dysphagia to solids; 2 = dysphagia to semisolids; 3 = dysphagia to liquids; 4 = complete dysphagia). Patients scores were obtained pre- and post-procedure. Success was defined as complete or near resolution of dysphagia; this was confirmed by the Dakkak and Bennett scores. Clinical success was achieved in 69 of the 75 patients; these patients had a decrease in their dysphagia score from 1.96 to .25. Only eight patients had follow-up for two years; thirty-one patients had a 12-month follow-up and fourteen had 18-months. The author's conclusions suggest that Z-POEM is safe and feasible for treatment of symptomatic ZD. However, limitations such as variations and restrictions in follow-up and lack of standardized management across multiple centers suggest additional comparative studies with long-term outcomes are needed.

Albers et al. (2016) conducted a systematic review analyzing endoscopic versus surgical treatment of Zenker's diverticulum. Out of 357 articles, 11 studies met the inclusion criteria, all cohort studies. Common endoscopic treatments included stapling of the diverticulotomy, CO2 laser and harmonic scalpel. Surgical approaches included cricopharyngeal myotomy and suspension, inversion, or excision of pouch, myotomy only and Dolman's procedure with pouch excision only. Meta-analysis revealed a significant reduction in the risk of recurrence of symptoms with use of the surgical approach compared to endoscopic treatment. However, for complications, it was shown fewer occurred with endoscopic treatment versus that of the surgical approach. The authors found when compared with a surgical approach, the endoscopic approach appears to result in shorter facility stays, earlier diet introduction and lower rates of complications, but demonstrates a higher rate of recurrence in symptoms. Limitations included studies with only retrospective cohorts (no RCTs comparing the techniques) and a large loss to follow-up.

Gastric Peroral Endoscopic Myotomy (G-POEM)

G-POEM is an innovative technique for the treatment of severe gastroparesis. There is insufficient quality evidence in the published clinical literature to support the safety and efficacy of this technique along with long-term effectiveness. There is significant patient heterogeneity regarding the etiology of gastroparesis (e.g., diabetic, idiopathic, medication induced, or postsurgical) as well as variability in the definition of clinical success. The generalizability of the positive findings are questionable as a large portion of the studies are performed at tertiary care centers with experienced interventional endoscopists. Future studies are essential for identifying the subgroup of patients which benefit from the intervention. Several clinical trials are in progress for the G-POEM; information can be found at <https://www.clinicaltrials.gov>.

Aziz et al. (2023) conducted a systematic review and meta-analysis comparing gastric per-oral endoscopic myotomy (G-POEM) with surgical pyloromyotomy for treating refractory gastroparesis. The meta-analysis, following Preferred Reporting items for Systematic Review and Meta-Analysis (PRISMA) guidelines, included a comprehensive search of MEDLINE, Embase, Web of Science, KCI – Koran Journal index, Global Index Medicus, and Cochrane databases. The authors identified four studies comprising 385 patients (216 in the G-POEM group and 169 in the surgical group), encompassing case-control, cohort, and randomized controlled trials (RCTs). Symptoms of gastroparesis were assessed using the Gastroparesis Cardinal Symptom Index (GCSI) score. Clinical success was defined as improvement in gastric emptying study (GES) and/or GCSI score according to study-specific criteria during follow-up. The meta-analysis found no significant difference in post-procedure GCSI scores (MD: -0.33, $p = 0.39$) or in reduction of GCSI scores from pre-to post-operatively (MD: 0.27, $p = 0.55$) between G-POEM and surgical pyloromyotomy. The authors suggested that G-POEM shows promise as a potentially cost-effective approach for managing refractory gastroparesis compared to surgery. They emphasized the need for future RCTs directly comparing different surgical techniques to better understand their relative benefits and efficacy. Limitations of this study included the scarcity of RCT's and a relatively small number of studies with fewer participants in each intervention group. They noted challenges in accounting for outcomes based on gastroparesis etiology, surgical approach (open, laparoscopic, robotic), and prior treatments like medications or botulinum toxin injections. Furthermore, the included studies predominantly involved participants from advanced tertiary care centers, raising concerns about generalizability of the findings. Inconsistent follow-up protocols across studies also hindered comprehensive assessment of important outcomes such as duration and timing of symptom improvement, GES results, and changes in GCSI scores.

In a systematic review, Stojilkovic et al. (2023) aimed to evaluate the long-term clinical effectiveness and safety of Gastric per-oral endoscopic myotomy (G-POEM), focusing on outcomes such as the Gastroparesis Cardinal Symptom Index (GCSI) score, adverse events, and hospital stay. They reviewed eleven studies involving 900 individuals, alongside references by Labonde et al. (2022) and Abdelfatah et al. (2021) as noted below. Due to variability in data definitions and reporting among studies, the authors applied a uniform clinical criterion: a one-point reduction in GCSI score. They found that clinical success, defined as a GCSI decrease of at least one point from baseline, was achieved in 92.8% of patients (662 out of 713) at one-year follow-up, 91.5% (421 out of 460) at two years, 100% (270 out of 270) at three years, and 100% (102 out of 102) at four years. Adverse events were reported in 62 out of 835 patients across nine studies, with common occurrences including bleeding and mucosal tears. Long-term adverse events were not extensively documented, and individual symptom assessments were limited due to varying inclusion across studies. The authors concluded that G-POEM is a safe and effective treatment option for patients with refractory gastroparesis, noting sustained symptom improvement up to four years post-procedure. They recommended future research to explore how gastroparesis etiology may influence treatment outcomes. Limitations of the review include potential bias from patient dropout during follow-up, which could skew results if dissatisfied patients were more likely to withdraw. Additionally, the review's reliance on two authors may have introduced bias in the selection of studies for inclusion or exclusion.

Martinek et al. (2022) conducted an RCT on forty-one patients with severe gastroparesis which compared G-POEM to a sham procedure. Participant eligibility included individuals over the age of 18 years with severe or refractory gastroparesis greater than six months. Severe gastroparesis was identified as a score > 2.3 on the Gastroparesis Cardinal Symptom Index (GCSI). Primary outcome measured was treatment success which was defined by a decrease of at least 50% in the total GCSI. Twenty-one participants were randomized to the treatment group where G-POEM was performed. Twenty individuals were included in the sham group which included upper GI endoscopy lasting at least 40 minutes; these participants were offered the G-POEM procedure if their symptoms were not relieved with the sham treatment. Clinical data was collected at three- and six-months. The authors found 15 of 21 patients in the G-POEM treatment group and four of 20 patients in the control group had success at the six-month follow-up. Twelve of fifteen patients from the control group were offered and agreed to have G-POEM following their sham treatment; nine of these participants achieved success. Ten serious adverse events (SAEs) occurred and three were found to be directly related to G-POEM. One participant developed a gastric ulcer near the pylorus, another developed a mucosal injury during the G-POEM and the third developed moderate dumping syndrome after three months with a need for hospitalization. The authors concluded the results demonstrated that G-POEM was beneficial in a substantial proportion of patients with severe and refractory gastroparesis (GP), however limitations included small sample size, lack of long-term outcomes, unclear masking to group assignment, and inability to accurately assess the relationship between the change in gastric emptying and symptomatic improvement. Future research should include larger sample sizes to assess the balance of benefit and harm, and an emphasis on long-term results.

Chung et al. (2022) evaluated the efficacy and safety of G-POEM for eleven patients with refractory gastroparesis; nine patients had diabetes, one had systemic lupus erythematosus and the other was idiopathic. Refractory gastroparesis was defined as persistent symptoms which failed lifestyle modification, failed medical treatment, contained adverse event(s) (AE) from medications or had repeated admission for nutritional support. Clinical response was defined as a decrease of > 25% in at least two subscales of cardinal symptoms of Gastroparesis Cardinal Symptom Index (GCSI) and improvement of gastric emptying scintigraphy (GES). While the complication rate in this study appeared higher than in other studies, the authors noted these were minor and resolved with conservative treatment. The authors found G-POEM was an efficient and safe procedure. Limitations include small sample size and no comparison groups; future studies are warranted and should include shams along with long-term outcomes.

Baret et al. (2022) assessed the rate of severe adverse events (AE) for patients that underwent G-POEM for refractory gastroparesis. Two hundred and seventeen patients (81 males, 136 females) from five French centers were included for analysis. G-POEM was performed by interventional endoscopists with a high level of expertise in performing the procedure. Postoperatively, the patients were monitored daily for pain and fever for one to four days before discharge. Follow-up occurred between three and six months and again at one year. Early postop complications included significant pain requiring analgesics, bleeding and one peripyloric abscess; late postop complications included melena with blood loss and three cases of dumping syndrome. A total of four patients were re-hospitalized, but no surgical intervention was required. The authors found less than 0.5% of serious AEs and thus confirmed G-POEM a safe procedure, however the data should be confirmed by larger prospective studies. The findings are limited by lack of comparison group and retrospective design.

Labonde et al. (2022) analyzed 46 patients who underwent G-POEM for refractory gastroparesis. These patients had persistent symptoms despite dietary control and prokinetic treatment for six months. Pre-procedural data obtained included Gastroparesis Cardinal Symptom Index (GCSI), patient assessment of Upper Gastrointestinal Disorders Quality of Life, and Gastrointestinal Quality of Life Index scores. Follow-up occurred in person or by phone at one, three, and six

months and then every six months thereafter. Gastric emptying scintigraphy (GES) was performed at three months after G-POEM. The primary endpoint was the assessment of clinical success at three years which included a decrease of at least one point in the GCSI score. The authors found the GCSI score decreased from 3.33 to 1.80 with improvement in nausea, satiety and bloating. The authors felt these improvements were significant and remained stable through three years. Limitations included study design, lack of comparison group, potential selection bias, and small sample size. Future large multicenter studies with comparison groups are necessary.

Mohan et al. (2020) conducted a systematic review and meta-analysis to evaluate the safety, efficacy, and predictive factors of G-POEM in the treatment of refractory gastroparesis. A search was conducted using PubMed, Embase, SCOPUS, and Web of Science. The authors included studies that evaluated the clinical outcomes of G-POEM and separate studies that evaluated the outcomes of surgical pyloroplasty in patients with medically resistant gastroparesis. Outcomes assessed included gastric emptying scintigraphy (GES) and scores from Gastroparesis Cardinal Symptom Index (GCSI). A total of 17 articles with 707 patients were included for review. The etiology for gastroparesis included idiopathic, diabetes, post-surgical and various unclassified causes. The authors found 76% of patients that had G-POEM had a reduction in clinical symptoms along with improved gastric emptying when compared to the pyloroplasty procedure. After having the G-POEM procedure, the mean GCSI score improved to 1.8, down from 3.4; the 4-h GES score improved from 49.9 to 20.6. Limitations included studies performed in tertiary-care referral centers which did not give a good representation of the population or community, subjective scoring of patient's symptoms and lack of long-term outcomes. The findings are limited by lack of direct comparison to contemporary controls within studies. While the authors concluded that G-POEM appears to be just as effective as surgical pyloroplasty, future studies are warranted.

In a retrospective case series, long-term outcomes of G-POEM for patients with refractory gastroparesis were investigated by Abdelfatah et al. (2021). The authors found out of 90 patients, 73 exhibited a positive outcome with symptom resolution and 17 were unsuccessful. Success was identified as a decrease of at least one point in the average total Gastroparesis Cardinal Symptom Index (GCSI) score with more than a 25% decrease in at least two subscales of cardinal symptoms. The successful patients also experienced a decrease in ER visits and hospitalizations. The authors concluded the long-term results of G-POEM demonstrate significant and sustainable improvement in the quality of the patient's life. Limitations included lack of comparison group, retrospective design, loss of patient follow-up and subjective patient-reported data.

Zhang et al. (2019) conducted a systematic review on the safety and efficacy of G-POEM for gastroparesis. A search was conducted using PubMed, Embase, Cochrane Library and Web of Science databases. A total of 14 trials (nine case series, four cohort and one case control study) with 276 individuals were included in the review. The etiology of gastroparesis varied amongst the studies but included idiopathic, diabetic and post-surgical gastroparesis. The main outcome was clinical safety and efficacy of the procedure. The Gastric emptying scintigraphy (GES) normalization and symptom improvement were considered the two main indicators for clinical efficacy. The follow up period varied amongst the studies, but an improvement in the Gastroparesis Cardinal Symptom Index (GCSI) score was seen in all of them. Most adverse events such as capnoperitoneum, pain and minor bleeding were considered mild and manageable. Amongst all the studies, only four patients from three different studies developed recurrent gastroparesis post procedure. The authors concluded G-POEM was an effective treatment for gastroparesis. Limitations included retrospective design of studies, lack of control groups, small sample sizes and lack of standardized evaluation for clinical improvement.

Endoscopic Therapies

Radiofrequency Energy (Stretta System)

Currently there is insufficient evidence regarding the safety and effectiveness of radiofrequency energy for gastroesophageal conditions. Studies are limited by small sample size, single center locations, and short-term follow-up. Meta analyses were limited by lack of control groups in a large portion of the included studies. Its role must be better defined in robust, well-designed clinical trials with long-term results.

A Hayes Health Technology Assessment (2023) evaluated the effectiveness and safety of Stretta radiofrequency (RF) to treat GERD. Fourteen articles found an overall low-quality body of evidence that suggests Stretta RF therapy may be safe and improve GERD symptoms for patients and quality of life for up to five years, however the evidence also reflects substantial uncertainty regarding its effectiveness when compared with laparoscopic fundoplication (LF), which is considered the surgical standard care. The overall conclusion of the report is that there is potential but unproven benefit for this technology.

In a 2020 randomized, double blind, sham controlled multi-center study, Zerbib et al. assessed the efficacy of esophageal radiofrequency (Stretta® system, Mederi Therapeutics) in sixty-two patients with moderate to severe gastro-esophageal reflux disease at least three times a week and refractory to proton pump inhibitors (PPIs). Completed questionnaires

consisting of the Gastrointestinal Symptoms Rating Scale (GSRS) and the Quality of Life in Reflux and Dyspepsia (QOLRAD) were collected. Patients were then randomized to receive either esophageal radiofrequency, or a sham procedure performed by a physician who would not be involved in follow up to maintain double blinding. All patients were instructed to take a double dose of PPIs after the procedure, and follow-up visits were planned at weeks four, eight, 12, 18, 24 and end of study at 48 weeks post-procedure to assess symptom relief, PPI use and any side effects. The intake of antacids as well as the presence of other digestive symptoms were also assessed. At each visit, if symptoms were adequately controlled, patients were instructed to decrease PPI from a double to a single dose, and as improvement continued, to “on demand” use. For patients who were having success, at week 24, an upper gastrointestinal endoscopy was performed (for therapeutic failures, the patients were offered an open esophageal radiofrequency procedure with the same follow-up). Five patients were lost to follow-up, and one withdrew his consent to participate, resulting in 26 patients being treated, and 30 patients treated with the sham procedure. The results showed that there was no significant difference between the treatment and sham groups at weeks 24 and 48 regarding days without heartburn, days without any other digestive symptoms, PPIs and antacids intake, and the number of patients not taking PPIs. There were no procedure related safety issues. The authors concluded that esophageal radiofrequency is a relatively invasive procedure for a benign disorder and did not demonstrate efficacy for the treatment of GERD refractory to PPIs.

Viswanath et al. (2019) reported a prospective case series of 50 patients who underwent endoscopic antireflux radiofrequency treatment (Stretta) for refractory GERD. Assessment involved the use of the Gastro-esophageal Reflux Disease-Health-Related Quality of Life (GERD-HRQL) questionnaire, which evaluated symptoms and PPI dependency, before and after treatment. Median follow-up post treatment was 771 days. The average GERD-HRQL score improved from 46.2/75 (± 14.2) before Stretta treatment to 15.2/75 (± 17.3) after Stretta treatment. The authors concluded that in select patients with GORD, Stretta improves quality of life and decreases PPI dependency, and is a viable option for patients who are unwilling or unable to undergo surgery. They also concluded that randomized controlled trials with larger patient populations are needed to further assess Stretta. Limitations of this study include lack of concurrent comparison group, its small numbers, and that the pre-Stretta assessments were carried out by a variety of teams thus the potential for inconsistencies.

In another case series, Noar et al. (2017) prospectively assessed and compared patient-reported outcomes in 18 patients refractive to laparoscopic Nissen fundoplication (LNF) and 81 patients with GERD refractory to medical management that all underwent Stretta during ten-year follow-up. Patient-reported outcomes measured were GERD-HRQL (health-related quality of life), patient satisfaction scores, and daily medication requirements. The refractory LNF subset demonstrated median improvements in GERD-HRQL, satisfaction, and medication use at all follow-up time points \geq six months to ten years, which was significant from a baseline of both on- and off-medications ($p < 0.05$). Specifically, at ten years, median GERD-HRQL decreased from 36 to seven ($p < 0.001$), satisfaction increased from one to four ($p < 0.001$), and medication score decreased from seven to six ($p = 0.040$). Nine patients decreased medication use by half at ten years. No significant differences existed between refractory LNF and standard refractory GERD subsets at any follow-up time point \geq six months to ten years ($p > 0.05$) after Stretta. At ten years, no significant differences were noted between refractory LNF and standard Stretta subsets regarding medication use ($p = 0.088$), patient satisfaction ($p = 0.573$), and GERD-HRQL ($p = 0.075$). Stretta procedures were completed without difficulty or significant intraoperative or long-term adverse events. The authors concluded that within a small series of patients with refractory LNF, Stretta resulted in sustained improvement over ten years with equivalent outcomes to non-LNF standard Stretta patients. Study limitations include lack of concurrent comparison group, non-randomization, and small patient population.

Kalapala et al. (2017) reported short outcomes (three months) from a prospective randomized study comparing the Stretta treatment with controls receiving proton pump inhibitors (PPIs). Patients ($n = 20$) with symptoms of heartburn, regurgitation, abnormal esophageal acid exposure ($\geq 4\%$), and endoscopically confirmed esophagitis were included into the study. The primary measure was improvement in quality of life (QOL) and decrease in the frequency and severity of GERD symptoms. The mean age of the patients was 39 (± 15) years and controls were 34 (± 11) years. Three months after Stretta, 80% reported improvement in QOL compared to 40% in the control group. At the end of three months, significant ($p < 0.05$) improvement in GERD symptom score for heartburn, regurgitation, chest pain, and cough compared with the control group was observed. After Stretta treatment, 60% of the patients were free of PPIs whereas there was no change in the control group. Almost 80% of the patients on Stretta treatment were satisfied with the treatment compared to 30% of the patients in the control group. The study was limited by the small sample size and short follow-up, therefore, randomized controlled trials with larger patient populations and longer follow-up periods are needed to further assess Stretta.

Fass et al. (2017) conducted a systematic review and meta-analysis of randomized controlled and cohort studies to determine the efficacy of the Stretta procedure in treating patients with GERD. Twenty-eight studies (four RCTs, 23 cohort studies, and one registry) representing 2,468 unique patients using Stretta were included in the meta-analysis. The (unweighted) mean follow-up time for the 28 studies was 25.4 (14.0, 36.7) months. The pooled results showed that the

Stretta reduced (improved) the health-related quality of life score by -14.6 (-16.48, -12.73) ($p < 0.001$). Stretta also reduced (improved) the pooled heartburn standardized score by -1.53 (-1.97, -1.09) ($p < 0.001$). After Stretta treatment, only 49% of the patients using proton pump inhibitors (PPIs) at baseline required PPIs at follow-up ($p < 0.001$). The Stretta treatment reduced the incidence of erosive esophagitis by 24% ($p < 0.001$) and reduced esophageal acid exposure by a mean of -3.01 (-3.72, -2.30) ($p < 0.001$). Lower esophageal sphincter (LES) basal pressure was increased post Stretta therapy by a mean of 1.73 (-0.29, 3.74) mmHg ($p = \text{NS}$). The authors concluded that the Stretta procedure significantly improves subjective and objective clinical endpoints, except LES basal pressure, and therefore should be considered as a viable alternative in managing GERD. The findings are however primarily explained by the included observational studies with overlap in some outcomes effect size between control and intervention groups of the included RCTs. Longer-term outcomes are needed to further evaluate the Stretta procedure. (Aziz et al. [2010], Coron et al. [2008], Dugher et al. [2014] and Noar et al. [2014] which were previously cited in this policy are included in this meta-analysis).

Arts et al. (2012) conducted a small double-blind randomized cross-over study of Stretta and sham treatment (included in the Fass et al. [2017] systematic review above). Patients underwent two upper gastrointestinal endoscopies with three months interval, during which active or sham Stretta treatment was performed in a randomized double-blind manner. In all, 22 GERD patients (17 females, mean age 47 ± 12 years) participated in the study; 11 in each group. Initial sham treatment did not affect any of the parameters studied. Three months after initial Stretta procedure, no changes were observed in esophageal acid exposure and lower esophageal sphincter (LES) pressure. In contrast, symptom score was significantly improved, and gastro-esophageal junction (GEJ) compliance was significantly decreased. Administration of sildenafil, an esophageal smooth muscle relaxant, normalized GEJ compliance to pre-Stretta level, arguing against GEJ fibrosis as the underlying mechanism. The authors concluded that Stretta improved GERD symptoms and decreased GEJ compliance. According to the authors, the limitation of this study was reflux evaluation did not include impedance monitoring. The study was also limited by a small patient population, short follow-up, and lack of comparison to other surgical alternatives.

Endoscopic Plication or Suturing

There is a lack of quality evidence to support the use of endoscopic plication or suturing for GERD. Studies are limited by definition of proper PPI use and evaluation prior to determining candidacy for procedures. Studies are limited by small sample size with lack of randomization, lack of control groups, and short duration of follow-up. Post hoc analysis introduces bias to outcomes. Additional studies are needed to support the safety and efficacy of these techniques with long-term effectiveness.

Testoni et al. (2021) conducted a systematic review and meta-analysis on long-term outcomes of transoral incisionless fundoplication (TIF) for patients with GERD. A search of publications through May of 2020 returned eight articles with long term outcomes of greater than three years for analysis. Outcomes evaluated in the analysis included overall patient satisfaction, daily proton pump inhibitor (PPI) consumption, GERD-health related quality of life (GERD-HRQL) scores, normalization of heartburn and regurgitation scores before and after the TIF procedure. The authors found TIF resulted in long-term patient satisfaction with reduction in PPI use in approximately 75% of the patients over a five-year period. At the ten-year mark, about two-thirds of the patients were satisfied. The findings are however limited by lack of comparison group for most of the included studies.

EndoCinch

Schwartz et al. (2013) evaluated the long-term effects of EndoCinch as a treatment option for GERD. Sixty patients were randomized into three different groups: Endocinch treatment, sham treatment, and observation. Inclusion criteria consisted of individuals with persistent heartburn and/or regurgitation, unwillingness to take lifelong medications, and esophageal pH results compatible with a GERD diagnosis. Baseline questionnaire utilizing a six-point numeric scale measuring heartburn and regurgitation frequency and a four-point numeric scale measuring severity was completed prior to study and again at three months; additional reassessment took place at six- and 12-months and then yearly thereafter. Quality of life assessments were done using the 20-item Short-Form Health Survey (SF-20). Endoscopic suturing was carried out with the Endocinch suturing device (BARD Endoscopic Technologies) which created three plications. During the first year any individuals with failure were offered one or two additional plications; three patients were lost to follow-up in the first year. Between 12 and 48 months, eight patients underwent antireflux surgery and one patient received an alternative endoscopic treatment. Four questionnaires were either incomplete or missing and therefore not used. In the end 43 participants were completely analyzed. At the end of follow-up, while the Endocinch procedure looked to improve GERD symptoms, decrease use of medications and increase the quality of life for approximately half the patients, 80% of those still required PPI management. The authors' concluded Endocinch in the long term was not beneficial. Limitations included small sample size and large loss to follow-up.

Endoscopic Plicator or Suturing

Niu et al. (2024) conducted a comprehensive systematic review and meta-analysis to assess the efficacy, safety, and long-term results of Endoscopic Fundoplication (EFTP) in treating GERD. Their analysis encompassed 13 studies, comprising of four RCTs and nine prospective cohort studies, with a total of 429 individuals undergoing EFTP and 146 individuals undergoing sham surgery for GERD treatment. Key studies included in this review were Kalapala et al. (2022), Antoniou et al. (2012), and Rothstein et al. (2006) cited below. Primary outcomes focused on improvements in GERD-related quality of life (GERD-HRQL) scores (> 50% improvement), cessation of proton pump inhibitor (PPI) use, and the need for laparoscopic fundoplication post-EFTP. Secondary outcomes included changes in GERD-HRQL scores, DeMeester scores, esophageal acid exposure time, total reflux events, Gastrointestinal Quality of Life Index (GIQLI), procedure duration, and adverse events associated with the procedure. The authors review of prospective studies indicated significant improvement in symptoms following EFTP, with 63% of patients achieving > 50% improvement in GERD-HRQL scores. Moreover, there was a substantial reduction in PPI dependency, with a cessation rate of 65%. EFTP also led to improvements in esophageal pH levels and DeMeester scores, demonstrating an average weighted mean difference (WMD) of -16.44. Additionally, there was a decrease in total reflux events and a low incidence of severe adverse events reported. Based on these findings, the authors suggest that EFTP could be a promising alternative for patients who do not respond adequately to conventional therapies. They recommend further research comparing EFTP with other minimally invasive GERD treatments, such as transoral incisionless fundoplication (TIF). They also emphasize the importance of long-term follow-up to assess the durability of treatment effects and to identify potential late-onset complications or symptom recurrence. Standardized follow-up protocols across studies are recommended to enhance comparability. The study has several limitations, including the need for long-term data on the durability of EFTP as a GERD treatment. Previous research on surgical GERD treatments has indicated significant rates of symptom recurrence after five years. Furthermore, while the majority of the participants were indeed resistant to long-term PPI treatment, not all studies reviewed included participants who were on long-term PPI therapy or unresponsive to it before undergoing EFTP, which could affect generalizability.

Kalapala et al. (2022) conducted a randomized, double blinded sham-controlled trial to assess the safety and efficacy of endoscopic full-thickness fundoplication (EFTP) in patients with GERD that were dependent on proton pump inhibitor (PPI) therapy. Seventy participants were assigned to one of two groups: one received the endoscopic full-thickness fundoplication (EFTP) and the other the sham therapy. The sham procedure positioned the device 1 cm below the gastro-esophageal junction, but the sutures were not deployed like in the EFTP procedure. Patient follow-up was completed at three, six, and 12 months along with telephone calls made every two to four weeks. In the EFTP group, 65.7% of patients obtained a 50% or more reduction in GERD-HRQL score compared with only 2.9% in the sham group. The PPI dependence at 12 months in the sham group was significantly higher than that of the EFTP group. The authors concluded that EFTP appears to be a new promising alternative to surgery for patients that may not want to continue with long-term PPI therapy, however larger trials with longer follow-up periods are required to confirm the benefits.

de Moura et al. (2018) evaluated long-term results of 47 patients non-responsive to proton pump inhibitors (PPIs) who underwent endoluminal plication (n = 26) or polymer injection (n = 21) for the treatment of GERD as part of a case series. The number of patients with no response to endoscopic treatment with reintroduction of PPIs increased in time for both techniques. There was symptomatic improvement up to 12 months, with progressive loss of this trending up to 60 months for both procedures. Health-related quality of life score (GERD-HRQL) demonstrated total response in both procedures at one, three, six, and 12 months. The 60-month analysis showed an increased number of patients with no response in both groups. The quality-of-life assessment (SF-36) showed benefit in polymer injection up to three months and showed a higher rate of complications. There were no deaths. There was healing of esophagitis at three months in 45% of patients in polymer injection and 40% in endoluminal plication. There was no improvement in manometric or pH findings. The authors concluded that endoscopic therapies were ineffective in controlling GERD in the long term. Limitations included lack of randomization and lack of uniform objective data analysis.

In an RCT, Antoniou et al. (2012) evaluated the effectiveness of endoscopic plication and laparoscopic fundoplication in terms of QOL and symptom control. A total of 60 patients with documented GERD were randomly assigned to undergo either endoscopic plication or laparoscopic fundoplication. QOL scores and symptom grading were recorded before treatment and at three- and 12-months of follow-up. Twenty-nine patients from the endoscopic group and 27 patients from the operative group were available at follow-up. QOL scores showed a substantial and similar increase for both groups after treatment. Symptoms of heartburn, regurgitation, and asthma were significantly improved in the endoscopic group, whereas laparoscopic fundoplication was more effective in controlling symptoms of heartburn and regurgitation compared to the endoscopic procedure. The authors concluded that endoscopic plication and laparoscopic fundoplication resulted in significant symptom improvement with similar QOL scores in a selected patient population with GERD, whereas operative treatment was more effective in the relief of heartburn and regurgitation at the expense of higher short-term dysphagia rates. Small sample size and lack of long-term follow-up limit the validity of these conclusions.

In a randomized, single-blind, prospective, multicenter trial by Rothstein et al. (2006), 159 patients were selected to either undergo endoscopic full-thickness restructuring of the gastric cardia with transmural suture (n = 78) or a sham procedure (n = 81) to determine the effectiveness of endoscopic full-thickness plication for the treatment of GERD. Group assignments were revealed following the three-month evaluation. By intention-to-treat analysis, at three months, the proportion of patients achieving $\geq 50\%$ improvement in GERD-HRQL score was significantly greater in the active group compared with the sham group. Complete cessation of PPI therapy was higher among patients in the active group than in the sham group. However, the median percent time that pH < 4 was not differently improved between the active and sham group. Between-group analysis revealed the active therapy was superior to sham treatment in improving the median percent time that the pH value was < 4. The authors concluded that endoscopic full-thickness plication was effective in reducing GERD symptoms and PPI use compared with a sham procedure. Additional studies are needed to evaluate the durability of endoscopic full-thickness plication for the treatment of GERD, as this study is limited by a relatively short follow-up.

GERDx™

Weitzendorfer et al. (2018) assessed the clinical safety and efficiency of the GERDx™ device by evaluating clinical parameters, reflux symptom scores, and quality of life (QoL) in a case series. Patients (n = 40) with at least one typical reflux symptom despite treatment with a PPI for > six months, pathologic esophageal acid exposure, hiatal hernia of size < 2 cm, and endoscopic Hill grade II-III were included. Evaluation of Gastrointestinal Quality of Life Index (GIQLI), symptom scores, esophageal manometry, and impedance-pH-monitoring were performed at baseline and at three months after surgery. Four out of forty patients experienced postoperative complications requiring intervention. Seven of forty patients were subjected to laparoscopic fundoplication three months after endoscopic plication due to persistent symptoms and were lost to further follow-up. Thirty out of forty patients were available at three-month follow-up. There was an improvement of the GIQLI score, from a mean of 92.45 ± 18.47 to 112.03 ± 13.11 ($p < 0.001$). The general reflux-specific score increased from a mean of 49.84 ± 24.83 to 23.93 ± 15.63 ($p < 0.001$), and the DeMeester score from a mean of 46.48 ± 30.83 to 20.03 ± 23.62 ($p < 0.001$). There was no significant change in manometric data after intervention. Three of thirty patients continued daily antireflux medication. The authors concluded that endoscopic plication with the GERDx™ device reduced distal acid exposure of the esophagus, reflux-related symptoms, and improved GIQLI scores with minimal side effects in a selected cohort of patients and may be a safe alternative in the treatment of GERD. Randomized clinical trials with larger patient populations and longer follow-up periods are needed to further assess GERDx.

MUSE™

Testoni et al. (2022) assessed the effects of transoral incisionless fundoplication (TIF) by MUSE on 46 patients. Inclusion criteria consisted of individuals > 18 and < 70 years of age, chronic GERD-related symptoms, endoscopic findings of GERD or Barrett's esophagus, body mass index < 40 kg/m², seeking alternative treatment and available for long-term follow-up. Symptoms and daily PPI consumption were assessed prior to the TIF-MUSE procedure, at six- and 12- months, and then yearly thereafter for at least three years. All patients completed the GERD-Health-Related Quality of Life (GERD-HRQL) and Reflux Symptom Index (RSI) questionnaires as tools to assess symptom severity of GERD. The authors found the GERD-HRQL and RSI scores were reduced by almost 50% in approximately 75% of the patients at the 6-month follow-up and decreased to 67.7% of patients at three years. Two severe adverse events occurred "as a consequence of delayed esophageal and intra-procedural gastric perforation at the stapler site." The authors concluded the MUSE™ device proved to be effective in about two-thirds of patients, but that severe complications requiring surgery occurred in two cases. Limitations included small sample size, lack of control group and inability to draw conclusions for patients with more severe degrees of esophagitis or patients with hypersensitive esophagus.

Kim et al. (2016) reported in a case series long-term outcomes from the Zacherl et al. (2015) MUSE study using the Medigus Ultrasonic Surgical Endostapler (MUSE™). Efficacy and safety data for 37 patients were analyzed at baseline, six months, and four years post-procedure. In one center (IU), efficacy and safety data were evaluated at baseline, six months post-procedure, and then annually up to four years. No new complications were reported in their long-term analysis. The proportions of patients who remained off daily PPI were 83.8% (31/37) at six months and 69.4% (25/36) at four years post-procedure. GERD-Health Related Quality of Life (HRQL) scores (off PPI) were significantly decreased from baseline to six months and four years post-procedure. The daily dosage of GERD medications, measured as omeprazole equivalents (mean \pm SD, mg), decreased from 66.1 ± 33.2 at baseline to 10.8 ± 15.9 at six months and 12.8 ± 19.4 at four years post-procedure (< 0.01). The authors concluded that the MUSE™ stapling device appears to be safe and effective in improving symptom scores as well as reducing PPI use in patients with GERD and that the results appeared to be equal to or better than those of the other devices for endoluminal GERD therapy. Future studies with larger patient series, sham control group, and greater number of staples are awaited to further evaluate MUSE. Findings are limited by lack of comparison group.

Zacherl et al. (2015) reported six-month outcomes from a multi-center prospective case series using the MUSE™ for the treatment of GERD (n = 69; three lost to follow-up). Six months after the procedure, the GERD-HRQL score improved by > 50% off PPI in 73% (48/66) of patients (95% CI 60-83%). Forty-two patients (64.6%) were no longer using daily PPI medication. Of the 23 patients who continued to take PPI following the procedure, 13 (56.5%) reported a ≥ 50% reduction in dose. The mean percent of total time with esophageal pH < 4.0 decreased from baseline to six months (p < 0.001). Common adverse events were peri-operative chest discomfort and sore throat. Two severe adverse events requiring intervention occurred in the first 24 subjects, no further esophageal injury or leaks were reported in the remaining 48 enrolled subjects. Early experience with the device necessitated procedure and device changes to improve safety, with improved results in the later portion of the study. Continued assessment of durability and safety are ongoing in a three-year follow-up study of this patient group. Findings are limited by lack of comparison group.

Other clinical trials regarding endoscopic plicator or suturing are limited to observational case series that do not allow for conclusions about durability and long-term effectiveness (Birk et al., 2009; von Renteln et al., 2009).

Polymer Injection and Implantation Techniques

Plexiglas and Durasphere

The available evidence for plexiglas and Durasphere techniques for gastroesophageal conditions is insufficient to consider the procedure proven to be effective and safe; additional randomized studies are warranted.

In a small case series, Ganz et al. (2009) assessed the long-term safety and effectiveness of Durasphere (Carbon Medical Technologies), an injectable bulking agent, in the treatment of mild to moderate GERD. Nine patients completed the 12-month trial. There were no adverse events. The procedure was well tolerated with minimal patient discomfort and no dysphagia. At 12 months, 70% of patients discontinued all antacid medication completely and 90% of patients reduced proton pump inhibitor (PPI) use by greater than 50%. There were no reports of esophagitis (at 12 months), erosion, ulceration, or sloughing of material at any injection site. The Durasphere material did not appear to migrate. The authors concluded that Durasphere appears to be a promising new injectable bulking agent for the treatment of mild to moderate GERD, with demonstrable efficacy and no significant adverse events in a small cohort of patients. Study limitations include lack of control group and small number of subjects.

Chen et al. (2009) conducted a systematic review that included 33 studies examining seven endoscopic procedures (Stretta procedure, Bard EndoCinch, Wilson-Cook Endoscopic Suturing Device, NDO Plicator, Enteryx, Gatekeeper Reflux Repair System and Plexiglas). Of the three procedures that were compared with sham controls (Stretta procedure, Bard EndoCinch and Enteryx), patient outcomes in the treatment group were either as good as, or significantly better than, those of control patients in terms of heartburn symptoms, quality of life (QOL), and medication usage. However, for the two procedures that were compared with the laparoscopic fundoplication (Stretta) procedure and the Bard EndoCinch device, outcomes for patients in the endoscopic group were conflicting. Some patients in the endoscopic group experienced comparable outcomes as patients undergoing the laparoscopic approach, while others experienced inferior outcomes. The authors concluded that there is insufficient evidence to determine the safety and efficacy of endoscopic procedures for GERD, particularly over the long term (Chen et al., 2009).

LINX Reflux Management System

There is insufficient evidence to conclude LINX is effective and safe on the long-term for GERD treatment. While short term results are promising, there is limited data on long-term effectiveness and safety of the LINX device. The follow-up periods in most studies may be insufficient to fully capture long-term complications or device durability. Additional research involving larger, randomized control trials with long-term outcomes is needed to establish its safety and efficacy, in the context of other mechanical approaches to GERD treatment that have shown benefits on the short-term but not on the long-term.

Valinoti et al. (2024) conducted a systematic review and meta-analysis in accordance with the PRISMA guidelines to evaluate the effectiveness and safety of magnetic sphincter augmentation (MSA) in patients with gastroesophageal reflux disease (GERD). The review included 22 studies with a total of 4,663 MSA patients. The author's findings indicated that the MSA device is highly effective, providing symptom relief and improved quality of life for most patients. However, postoperative dysphagia and the need for endoscopic dilation are relatively common, and while the risk of esophageal erosion is low, it significantly increases over time. The authors recommend further studies with objective assessments and longer follow-up periods. They also emphasize the importance of standardized diagnostic methods for more objective and comparable outcome assessments. They noted the study has several limitations. Firstly, there were methodological design discrepancies among the included studies. Secondly, the majority of the studies had a short follow-up period. Thirdly, there was significant statistical heterogeneity in many of the outcomes assessed. Lastly, few studies evaluated patients using postoperative pH monitoring. (Ganz [2016], Louie [2019], and Ferrari [2020] included below).

An ECRI clinical assessment on the LINX® Reflux Management System for treating GERD identified a review of evidence from 2017 through 2023 that included three systematic reviews, one randomized control trial, one patient registry study and two economic studies. The evidence appears to indicate that LINX may be safe and as effective as laparoscopic Nissen fundoplication, however the studies had a high risk of bias and not enough patients were included to confirm the conclusions. It was concluded larger multicenter RCTs and longer follow-up with comparisons of LINX with other GERD devices would be useful to determine long-term safety and efficacy (ECRI 2023).

A Hayes Health Technology Assessment (2022) evaluated the effectiveness and safety of magnetic sphincter augmentation (MSA) with the LINX Reflux Management System for the treatment of GERD. One RCT compared MSA with twice-daily PPI treatment and the other five nonrandomized cohort studies compared MSA with laparoscopic fundoplication (LF). While the low-quality body of evidence suggests that MSA may improve the patient's quality of life and their GERD symptoms along with a reduction in proton pump inhibitor (PPI) use, there remains uncertainty about the long-term safety and efficacy of this device. The report overall conclusion is that there is potential but unproven benefit to the LINX system.

Zhuang et al. (2021, included in the Hayes report above) performed a systematic review and meta-analysis to determine the efficacy and safety of magnetic sphincter augmentation (MSA) in the management of refractory GERD as well as comparing MSA efficacy to proton pump inhibitor (PPI) or laparoscopic Nissen fundoplication (LNF). Ten single-arm studies, one randomized controlled trial and three cohort studies involving 1138 participants were included. Post-MSA PPI withdrawal, significant GERD-health-related quality of life (GERD-HRQL) improvement and acid exposure time (AET) normalization were achieved in 87.0%, 88.0% and 75.0% of the patients, individually. The incidence of postoperative dysphagia was 29% and endoscopic dilation was required in 7.4% of patients undergoing MSA. MSA showed a better efficacy in symptom control than PPI (PPI cessation: 91% vs 0%; GERD-HRQL improvement: 81% vs 8%) and similar effectiveness but a lower risk of gas-bloat syndrome (risk ratio [RR] 0.69, 95% confidence interval [CI] 0.51-0.93, $p = 0.01$) and better reserved ability to belch (RR 1.48, 95% CI 0.76-2.86, $p = 0.25$) compared with LNF. Study limitations included the following: limited number of clinical studies; only three studies were included in the comparative results between MSA and LNF; and potential for selection bias that could have led to an overestimation of efficacy in MSA since the patient selection had less severe GERD. The authors concluded that MSA was an effective and safe therapy for GERD for patients with PPI-refractory symptoms and pathological reflux. There was only one randomized comparative trial that presented an advantage over a double dose of PPI. Therefore, there is a need for additional randomized trials that compare the efficacy of MSA with other therapies (Bonavina [2010], Lipham [2012], Ganz [2103 and 2016], Saino [2015] and Warren [2016] previously cited in this policy are included in this meta-analysis).

Chandan et al. (2021) conducted a systematic review and meta-analysis on patients undergoing treatment for refractory GERD and compared the efficacy of magnetic sphincter augmentation (MSA) with that of transoral incisionless fundoplication (TIF2). Twenty-four articles were included in the analysis which consisted of 1,074 patients that underwent MSA, and 868 patients underwent TIF2. In the MSA cohort, six studies were prospective, and three studies were retrospective; in the TIF cohort, eleven studies were prospective and four were retrospective. The authors found the clinical success rate, demonstrated by improvement of scores in GERD-health-related quality of life (GERD HRQL), was 80% for MSA and 77% for TIF. It was concluded that both procedures have a similar efficacy, but MSA seem to outperform TIF2. Overall, 91.3% of the MSA patients were able to discontinue PPI therapy compared to 63.8% of the TIF2 patients. Limitations included lack of long-term data.

Bell et al. (2020, included in the Hayes report above) compared the effects of magnetic sphincter augmentation (MSA) versus proton pump inhibitor (PPI) in a randomized trial. One hundred fifty-two patients with moderate to severe regurgitation symptoms across twenty-one U.S. clinical sites were randomized into two groups. Additional inclusion criteria for the participants were once daily PPIs for at least eight weeks, body mass index < 35 kg/m², abnormal pH testing (DeMeester score < 4), hiatal hernia < 3 cm by endoscopy and absence of Barrett's esophagus or Los Angeles Classification Grade C or D esophagitis. Participants were assessed at six and 12 months with the Reflux Disease Questionnaire (RDQ) and the GERD Health Related Quality of Life (GERD-HRQL) standard assessments along with specific questions concerning bloating, diarrhea, flatulence, and medication use. One group ($n = 102$) received PPI (20 mg of omeprazole twice daily) and the other group ($n = 50$) received laparoscopic MSA. The authors found that MSA controlled regurgitation in 96% of patients versus only 19% of patients receiving PPIs reported control of regurgitation. The regurgitation had been sustained over 12 months. The second portion of the study allowed eligible patients (39%) from the PPI group to crossover and receive the laparoscopic MSA if they had not demonstrated improvement with the twice daily medication. The authors concluded MSA is an effective surgical treatment option for patients with medically refractory regurgitative GERD. The study is limited by limited follow-up.

Ferrari et al. (2020) followed a cohort of 124 individuals who underwent laparoscopic implantation of a magnetic sphincter augmentation (MSA) device. The goal was to assess the long-term safety and efficacy of the Linx Reflux Management

System for six to 12 years. Prior to surgery, all patients completed a diagnostic assessment that included the GERD-health-related quality of life (GERD-HRQL) questionnaire, upper GI endoscopy, barium swallow study, ambulatory esophageal pH monitoring and esophageal manometry. Success was defined as $\geq 50\%$ improvement in the GERDHRQL total score and discontinuation of proton pump inhibitor (PPI) medication. During follow up, over a five-year period, eight patients (2.4%) required a single endoscopic pneumatic dilation due to persistent dysphagia. Thirty-one patients (9.2%) required removal of the device for various reasons; erosion and regurgitation were the top two reasons with six patients each. The average total GERD-HRQL score decreased from 19.9 (baseline) to 4.01. The authors found eighty-one percent of the patients had a successful clinical outcome and were able to discontinue their PPI use. Long-term results in thirty-two patients past ten years found zero dysphagia, seven individuals with occasional PPI use and only three with daily PPI use. The total overall patient satisfaction rate was 92.5%. The authors concluded MSA allows control of GERD symptoms and improvement in patient quality of life without significant safety issues. However, it was also concluded that additional RCTs could provide more definitive conclusions. Limitations included no comparison group and possible selection bias with large loss to follow up over time.

Schizas et al. (2020) conducted a systematic review to investigate the safety and efficacy of the LINX® Reflux Management System. After screening 614 articles, a total of 35 studies fit the criteria and were analyzed. According to the authors, although laparoscopic fundoplication (LF) and magnetic sphincter augmentation (MSA) both appear to be safe and effective procedures, MSA appears to have a few distinct advantages such as a less technical procedure, less bloating and superiority in the ability to vomit/belch, easily reversible and if it fails, LF is still a viable option after device removal. The authors' findings suggested that MSA with the LINX device is a safe procedure and has the potential to bridge the treatment gap between maxed out medical treatment and laparoscopic fundoplication. The authors also concluded that further studies with longer follow-up are needed (Asti [2016], Desart [2015], Reynolds [2015], and Bonavina [2008 and 2010] previously cited in this policy are included in this systematic review).

A prospective, multicenter, randomized control trial was conducted by Bell et al. (2019, included in the ECRI report) comparing magnetic sphincter augmentation (MSA) (n=50) to double-dose proton-pump inhibitor (PPI) therapy (omeprazole, 20 mg, twice a day) (n=102). The goal of the study was to compare the effect of the two treatments for elimination of moderate to severe regurgitation. As reported on a foregut symptom questionnaire, at six months, 89% of patients treated with MSA reported relief of regurgitation, with 81% reporting $\geq 50\%$ improvement in GERD-health-related quality of life scores. Ten percent of the PPI group reported relief of regurgitation with eight percent of the PPI group reporting $\geq 50\%$ improvement in GERD-health-related quality of life scores. However, 28 percent of MSA patients reported transient dysphagia, with 4% reporting ongoing dysphagia. The authors concluded that patients who continue to experience moderate to severe regurgitation despite PPI treatment should be considered for MSA. Randomized controlled trials with larger patient populations and long term follow up are needed to further assess the long-term safety and efficacy of MSA.

Louie et al. (2018, included in the Schizas et al. (2020) systematic review above) reported one-year results from a mandated post-approval multicenter, prospective case series of 200 patients with pathologic acid reflux confirmed by esophageal pH testing, who underwent magnetic sphincter augmentation (MSA). Predefined clinical outcomes were assessed at the annual visit including a validated, disease-specific questionnaire, esophagogastroduodenoscopy (EGD) and esophageal pH monitoring, and use of proton pump inhibitors. At one year, the mean total acid exposure time decreased from 10.0% at baseline to 3.6%, and 74.4% of patients had normal esophageal acid exposure time (% time pH $< 4 \leq 5.3\%$). GERD Health-Related Quality of Life scores improved from a median score of 26.0 at baseline to 4.0 at one year, with 84% of patients meeting the predefined success criteria of at least a 50% reduction in total GERD Health-Related Quality of Life score compared with baseline. The device removal rate at one year was 2.5%. There was a report of one erosion, and no serious adverse events were reported. Although the authors conclude that safety and effectiveness of MSA has been demonstrated outside of an investigational setting, study limitations include lack of contemporaneous comparison group receiving a different GERD treatment and relatively short follow-up period.

In a retrospective observational study, Warren et al. (2018, included in the Schizas et al. [2020] systematic review above) analyzed factors influencing the outcome of magnetic sphincter augmentation (MSA) for chronic GERD using data from a pivotal trial (n = 99) and the authors prospectively maintained esophageal database (n = 71). A priori outcomes were defined as excellent (GERD-HRQL < 5 , no PPI, no esophagitis), good (GERD-HRQL 6-15, no PPI, grade A esophagitis), fair GERD-health-related quality of life (GERD-HRQL 16 to 25, PPI use, grade B esophagitis), and poor (GERD-HRQL > 25 , PPI use, grade C/D esophagitis). Univariable and multivariable logistic regression analyses were performed to determine predictors of achieving an excellent/good outcome. A total of 170 patients underwent MSA with a median age of 53 years, (43-60) and a median BMI of 27 (IQR = 24-30). At baseline, 93.5% of patients experienced typical symptoms and 69% atypical symptoms. At univariable analysis, excellent/good outcomes were negatively impacted by BMI, preoperative lower esophageal sphincter (LES) residual pressure, Hill grade, and hiatal hernia. At multivariable analysis, BMI > 35 (OR = 0.05, 0.003-0.78, p = 0.03), structurally defective LES (OR = 0.37, 0.13-0.99, p = 0.05), and preoperative

LES residual pressure (OR = 0.89, 0.80-0.98, $p = 0.02$) were independent negative predictors of excellent/good outcome. The authors' conclusion is that MSA results in excellent/good outcomes in most patients but a higher BMI, structurally defective sphincter, and elevated LES residual pressure may prevent this goal. The authors' conclusion is that a higher BMI, structurally defective sphincter, and elevated LES residual pressure may prevent optimal treatment with MSA. The findings however do not provide evidence for the safety and efficacy of MSA compared to other therapeutic approaches.

Aiolfi et al. (2018, included in the Schizas et al. [2020] systematic review above) conducted a systematic review and meta-analysis of early results of magnetic sphincter augmentation (MSA) versus fundoplication for the treatment of GERD. Seven observational cohort studies, published between 2014 and 2017, matched the inclusion criteria. Overall, 1,211 patients, 686 MSA and 525 laparoscopic fundoplication (LF), were included. Postoperative morbidity ranged from 0 to 3% in the MSA group and from 0 to 7% in the LF group, and there was no mortality. Dysphagia requiring endoscopic dilatation occurred in 9.3% and 6.6% of patients respectively (OR = 1.56, 95% CI = 0.61-3.95, $p = 0.119$). The pooled OR of gas/bloat symptoms, ability to vomit, and ability to belch were 0.39 (95% CI 0.25-0.61; $p < 0.001$), 10.10 (95% CI 5.33-19.15; $p < 0.001$), and 5.53 (95% CI 3.73-8.19; $p < 0.001$), respectively. The postoperative GERD-health-related quality of life (GERD-HRQL) was similar ($p = 0.101$). The pooled OR of proton pump inhibitor (PPI) suspension, endoscopic dilation, and reoperation were similar in the two patients' groups ($p = 0.548$, $p = 0.119$, $p = 0.183$, respectively). The authors concluded that both anti-reflux procedures are safe and effective up to one-year follow-up. PPI suspension rate, dysphagia requiring endoscopic dilatation, and disease-related quality of life are similar in the two patient groups. MSA is associated with less gas/bloat symptoms and increased ability to vomit and belch. The findings are limited by inclusion of observational studies only and relatively short follow-up periods.

Alicuben et al. (2018, included in the Schizas et al. [2020] systematic review above) reported on the worldwide experience with erosion of the magnetic sphincter augmentation (MSA) device in a large case series. In total, 9,453 devices were placed and there were 29 reported cases of erosions. The median time to presentation of an erosion was 26 months with most occurring between one and four years after placement. The risk of erosion was 0.3% at four years after device implantation. Most patients experienced new-onset dysphagia prompting evaluation. Devices were successfully removed in all patients most commonly via an endoscopic removal of the eroded portion followed by a delayed laparoscopic removal of the remaining beads. At a median follow-up of 58 days post-removal, there were no complications and 24 patients have returned to baseline. Four patients reported ongoing mild dysphagia. The authors concluded that erosion of the LINX device is an important but rare complication to recognize that has been managed via minimally invasive approaches without long-term consequences. Continued monitoring and reporting of MSA erosion will provide longer-term experience.

In a systematic review and meta-analysis of the LINX[®] magnetic esophageal sphincter augmentation (MSA) versus Nissen fundoplication for gastroesophageal reflux disease, Skubleny et al. (2017) included randomized controlled trials, non-randomized comparison study and case series with greater than five patients. Five hundred and forty-seven titles were identified through primary search, and 197 titles or abstracts were screened after removing duplicates. Meta-analysis was performed on postoperative quality of life outcomes, procedural efficacy, and patient procedural satisfaction. Three primary studies identified a total of 688 patients, of whom 273 and 415 underwent Nissen fundoplication and MSA, respectively. MSA was statistically superior to laparoscopic Nissen fundoplication (LNF) in preserving patient's ability to belch (95.2 vs. 65.9%, $p < 0.00001$) and ability to emesis (93.5 vs. 49.5%, $p < 0.0001$). There was no statistically significant difference between MSA and LNF in gas/bloating (26.7 vs. 53.4%, $p = 0.06$), postoperative dysphagia (33.9 vs. 47.1%, $p = 0.43$) and proton pump inhibitor (PPI) elimination (81.4 vs 81.5%, $p = 0.68$). The authors' conclusion is that magnetic sphincter augmentation appears to be an effective treatment for GERD with short-term outcomes comparable to the more technically challenging and time-consuming Nissen fundoplication. The authors also concluded that long-term comparative outcome data past one year is needed in order to further understand the efficacy of magnetic sphincter augmentation.

Smith et al. (2017) reported that out of a total of 3283 procedures reviewed for magnetic sphincter augmentation device (MSAD), device removal occurred in 2.7% of cases. The most common causes of removal were dysphagia, continued reflux, and device erosion into the esophagus. Salvador et al. (2017), Parmar et al. (2017), and Lipham, et al. (2015), report similar findings.

Ganz et al. (2016, included in the Schizas et al. [2020], Zhuang et al. [2021] and Valinoti et al. [2024] systematic reviews above) reported in a case series the five-year follow-up evaluation of patients who received a magnetic sphincter augmentation (MSA) device for GERD. The original prospective study at 14 centers in the United States and the Netherlands was conducted on 100 adults with GERD for six months or more, who were partially responsive to daily proton pump inhibitors (PPIs) and had evidence of pathologic esophageal acid exposure. At baseline, the median GERD-health-related quality of life (GERD-HRQL) scores were 27 in patients not taking PPIs and 11 in patients on PPIs; five years after device placement this score decreased to four. All patients used PPIs at baseline; this value decreased to

15.3% at five years. Moderate or severe regurgitation occurred in 57% of subjects at baseline, but only 1.2% at five years. All patients reported the ability to belch and vomit if needed. Bothersome dysphagia was present in 5% at baseline and in 6% at five years. Bothersome gas-bloat was present in 52% at baseline and decreased to 8.3% at five years. The authors concluded that MSA provides significant and sustained control of reflux, with minimal side effects or complications, which in their opinion validates the long-term safety and efficacy of MSA for patients with GERD. The study is however limited by lack of comparison group.

Riegler et al. (2015, included in the Hayes report) evaluated using a retrospective cohort study design the evidence for magnetic sphincter augmentation device (MSAD) and laparoscopic fundoplication (LF) in clinical practice. Two hundred forty-nine patients (202 MSAD patients and 47 LF patients) had completed one-year follow-up. The LF group was older and had a greater frequency of large hiatal hernias and Barrett's esophagus than the MSAD group ($p < 0.001$). The median GERD-health related quality of life score improved from 20.0 to 3.0 after MSAD and 23.0 to 3.5 after LF. Moderate or severe regurgitation improved from 58.2 to 3.1% after MSAD and 60.0 to 13.0% after LF ($p = 0.014$). Discontinuation of proton pump inhibitors (PPIs) was achieved by 81.8% of patients after MSAD and 63.0% after LF ($p = 0.009$). Excessive gas and abdominal bloating were reported by 10.0% of patients after MSAD and 31.9% following LF ($p \leq 0.001$). Following MSAD, 91.3% of patients were able to vomit if needed, compared with 44.4% of those undergoing LF ($p < 0.001$). Reoperation rate was 4.0% following MSAD and 6.4% following LF. The authors conveyed that antireflux surgery should be individualized to the characteristics of each patient, taking into consideration anatomy and propensity and tolerance of side effects. They concluded that both MSAD and LF showed significant improvements in reflux control, with similar safety and reoperation rates. In their opinion, in the treatment continuum of antireflux surgery, MSAD should be considered as a first-line surgical option in appropriately selected patients without Barrett's esophagus or a large hiatal hernia in order to avoid unnecessary dissection and preserve the patient's native gastric anatomy. The study is however limited by lack of randomization.

Lipham et al. (2015) conducted a case series of antireflux surgery with a Magnetic Sphincter Augmentation Device (MSAD). The aim of the study was to examine the safety profile of the MSAD in the first 1,000 implanted patients. The author compiled data from multiple sources starting on July 1, 2013. The analysis included intra/perioperative complications, hospital readmissions, procedure-related interventions, reoperations, and device malfunctions leading to injury or inability to complete the procedure. The authors report that approximately 1,000 patients worldwide have been implanted with the MSAD, at 82 institutions with median implant duration of 274 days. They concluded that the safety analysis of the first 1,000 patients treated with MSAD for gastroesophageal reflux disease confirms the safety of this device and the implantation technique. The preliminary and positive results of this study are hampered by lack of an adequate comparator group.

Functional Lumen Imaging Probe (FLIP)

While the FLIP technology is used to enhance current procedures such as endoscopy, the evidence is insufficient for use in assessing esophageal and gastric disorders. The utility is yet to be defined for the diagnostic work-up, procedure guidance or post-procedure evaluation of clinical outcomes. Additional research involving larger, RCTs with long-term outcomes is needed to establish its safety and efficacy.

Rafeeqi et al. (2023) analyzed the utility of EndoFLIP in diagnosis and management of achalasia in pediatric patients. The authors theorized that the distensibility index (DI) may aid diagnosis and treatment of pediatric achalasia. Thirty-three (21 male, 12 female) patients underwent POEM and EndoFLIP. The authors were able to show that certain measurements, such as Eckardt scores, had a direct correlation to achalasia symptoms. The higher the Eckardt score, the lower DI. While manometry is the current gold standard test for the diagnosis of achalasia, it is not tolerated well in children which makes the EndoFLIP a potential alternative for the pediatric population. The authors concluded that EndoFLIP could be beneficial in the pediatric population in the diagnosis and management of achalasia. The DI could be used to guide the extent of myotomy and monitor the results of any lower esophageal sphincter LES focused interventions, however further studies are necessary to better support the use of EndoFLIP as a diagnostic tool. Limitations included small sample size, lack of manometry data for comparison and lack of long-term data.

A Hayes report (2022) for the use of EndoFLIP evaluated the device for its use in assessment of esophageal and gastric disorders. The evidence is insufficient to support any conclusions for any particular use of the device for any one disorder and found no improvement or benefit to health outcomes. Out of 13 studies, 11 reviewed and evaluated EndoFLIP for detecting or predicting disease course or treatment response. The other two studies evaluated the clinical utility for guidance and management of gastric disorders in addition to bariatric surgery. While the body of evidence for EndoFLIP was large for a variety of topics, the quality of evidence was low with a large range of uses. Limitations included lack of comparison groups, lack of systematic evaluation and lack of long-term outcomes.

An ECRI (2020) report on EndoFLIP for identification of GI motility disorders assessed two prospective and two retrospective single-center case series texts from January of 2015 thru May of 2020. The evidence found was considered inconclusive with mixed results. Limitations included a high risk of bias due to retrospective design, small sample size and lack of comparison groups.

In a systematic review, Desprez et al. (2020) summarized the available data in the literature on the use of the EndoFLIP® system in the gastrointestinal tract. A search was conducted using MEDLINE-PubMed, Cochrane Library, and Google Scholar databases. A total of 95 studies were found and 32 of these reported results in patients with achalasia. The authors found the EndoFLIP system demonstrated its relevance in the diagnosis of atypical achalasia and in the prediction of the treatment outcome in achalasia, but the precise esophago-gastric junction (EGJ) distensibility index (DI) threshold associated with success and prolonged response remains to be determined and therefore warrants additional studies. In addition, review of thirteen studies was completed which investigated EGJ in GERD patients. Patients were assessed with upper GI endoscopy and esophageal pH monitoring in addition to symptomatic evaluation. The authors found the patients with GERD seemed to have a higher EGJ DI than the healthy volunteers tested, but the relevance of the EndoFLIP® system in the prediction of the efficacy of fundoplication remained unconfirmed. Limitations included no standard protocol found when using the EndoFLIP® in the esophagus along with cost and availability in daily practice. In conclusion, while the authors found EndoFLIP® might have some promise, its use in the body of the esophagus, other esophageal diseases (GERD, eosinophilic esophagitis), and other sphincter regions (anal canal, pylorus) necessitates the need for additional confirmatory studies.

Yoo et al. (2019) assessed the utility of EndoFLIP in fifty-two patients with achalasia treated with POEM and hypothesized that improvement in the distensibility index (DI) correlated with the postoperative clinical outcome of POEM. EndoFLIP was performed before and after POEM by well-trained endoscopists in an outpatient setting. Esophagogastroduodenoscopy (EGD) was done just before EndoFLIP to measure esophageal length, four days following the procedure and again in outpatient clinic to evaluate the development of reflux esophagitis. The DI of the esophagogastric junction (EGJ) was measured at the point of narrowest perimeter of the functional lumen image during volumetric distension of the saline bag. After the probe was inserted, the DI and cross-sectional area (CSA) of the EGJ were measured at the site of the narrowest point. DI and CSA were measured twice during volumetric distension of the saline bag, each of them with 30 mL and 40 mL. The authors' conclusion was EndoFLIP was useful in predicting the treatment response and post procedure reflux of POEM in patients with achalasia. Limitations included lack of comparison groups, lack of long-term outcomes and small sample size.

Clinical Practice Guidelines

American College of Gastroenterology (ACG)

In a ACG clinical guideline (Camilleri et al., 2022) for gastroparesis, the ACG suggests pyloromyotomy over no treatment for patients with gastroparesis that have not responded to medical therapy and continue to experience symptoms (conditional recommendation, low quality of evidence). The preferred myotomy method (laparoscopic or endoscopic) is however not specified.

In a 2021 ACG published clinical guideline (Katz et al., 2021) for the diagnosis and management of GERD, the following recommendations are cited:

- Recommend antireflux surgery as an option for long-term treatment of patients with objective evidence of GERD, (strong recommendation; moderate level of evidence).
- Recommend consideration of magnetic sphincter augmentation (MSA) as an alternative to laparoscopic fundoplication for patients with regurgitation who fail medical management (strong recommendation, moderate level of evidence).
- Consideration of transoral incisionless fundoplication (TIF) for patients with troublesome regurgitation or heartburn who do not wish to undergo antireflux surgery and who do not have severe reflux esophagitis or hiatal hernias (conditional recommendation; low level of evidence).
- Do not recommend radiofrequency energy (Stretta) as an antireflux procedure due to inconsistent data on the efficacy of the device (conditional recommendation; low level of evidence).

American Gastroenterological Association (AGA)

The AGA recommends POEM be considered as a primary therapy for type III achalasia. Given the complexity of the POEM procedure, the AGA also recommends the procedure be performed by experienced physicians in high-volume centers to achieve procedure competence (Kahrilas et al., 2017).

An AGA clinical practice update (Khashab et al., 2023) on gastric peroral endoscopic myotomy for gastroparesis offered expert advice regarding cognitive, procedural, and post-procedural aspects on performance of gastric peroral endoscopic

myotomy for the treatment of refractory gastroparesis. After review of the available evidence, it was concluded G-POEM is safe with a high technical success rate when performed by an experienced endoscopic specialist.

An AGA clinical practice update offers guidance on best practices for personalized diagnosis and treatment of gastroesophageal reflux disease (GERD). The AGA recommends a stepwise approach for individuals presenting with GERD symptoms, aimed at identifying underlying mechanisms. When symptoms persist despite lifestyle changes and optimal medication use, management strategies should be tailored based on factors such as the integrity of the anti-reflux barrier, presence of visceral hypersensitivity and hypervigilance, confirmation of PPI-refractory GERD, symptom patterns, body mass index, and esophageal (and gastric) motor function. (Yadlapati et al., 2022).

American Society for Gastrointestinal Endoscopy (ASGE)

In a 2015 clinical guideline on the role of endoscopy in the management of GERD, ASGE suggests that endoscopic antireflux therapy be considered for selected patients with uncomplicated GERD after careful discussion with the patient regarding potential adverse effects, benefits, and other available therapeutic options.

The ASGE identifies laparoscopic Heller myotomy, pneumatic dilation, and POEM as effective therapeutic modalities for patients with achalasia. The decision made between these treatment options should depend on achalasia type, local expertise, and patient preference (Khashab et al., 2020).

American Society of General Surgeons (ASGS)

In 2014, the ASGS published a position statement regarding its support for the LINX procedure. ASGS states that total management of GERD will likely rely upon a combination of medical and surgical care in the current and near future. ASGS recommends that when considering a surgical procedure, the procedure will need to provide safe control of GERD with minimal side effects. The ASGS states, "Based on currently available information and the experience of our members with the procedure, we do support the LINX procedure as a mechanism for controlling GERD when it is placed by properly trained laparoscopic surgeons with experience in foregut surgery and the management of GERD patients."

In April 2011, the ASGS published a position statement regarding the use of TIF stating that it supports the use of TIF in patients with symptomatic chronic GERD who are not responsive to a standard dose of PPI therapy (ASGS, 2011). The ASGS also supports its use for patients who wish to avoid lifetime drug therapy for this condition. The ASGS also supports the adoption of the procedure by trained general surgeons as a less invasive alternative to more conventional surgical techniques, stating that the preferred surgical technique should be based on the discretion and judgment of the surgeon and the patient's clinical circumstances.

In a statement regarding coverage for TIF, ASGS states that there is a sufficient body of peer reviewed literature that establishes transoral fundoplication as reasonable and medically necessary for a subset of patients who are candidates for surgical fundoplication; specifically, patients who either cannot obtain satisfactory relief from standard PPI therapy or who wish to avoid a lifetime of dependence on such medications, and present with a two centimeter or smaller hiatal hernia (ASGS, 2011).

National Institute for Health and Care Excellence (NICE)

The National Institute for Health and Care Excellence (NICE, 2023) conducted a rapid review of the published literature on the efficacy and safety of laparoscopic insertion of a magnetic ring for gastro-esophageal reflux disease (GORD). They evaluated evidence from three systematic and meta-analyses, one RCT, three non-randomized comparative studies, and two case series. The committee noted that the magnetic ring use has evolved over time and is considered adequate. In addition, it was stated that the use of the device should be performed by clinicians that have specific training in the procedure.

The NICE guideline on endoscopic radiofrequency ablation for GERD considers the evidence on this procedure to be adequate in the short and medium term but there is uncertainty about longer-term outcomes. Regarding efficacy, there is evidence of symptomatic relief but objective evidence on reduction of reflux is inconclusive (NICE, 2013).

Society of American Gastrointestinal and Endoscopic Surgeons (SAGES)

SAGES guidelines (Kohn et al., 2021) on the use of POEM for the treatment of achalasia make the following recommendations:

- The panel suggests that adult and pediatric patients with type I and II achalasia may be treated with either POEM or laparoscopic Heller myotomy based on surgeon and patient's shared decision-making (conditional recommendation, very low certainty evidence).

- Based on the panel's collective experience, they suggest POEM over laparoscopic Heller myotomy for type III adult or pediatric achalasia (expert opinion).

The following recommendations are made by SAGES (Auyang et al., 2013) on endoluminal treatments for GERD:

- Results for EsophyX appear to be mixed and lacking long term data. Further studies are required to define optimal techniques and to further evaluate the device and its safety (Quality of Evidence: ++). GRADE Recommendation: Weak).
- The evidence for the Stretta device demonstrates it is considered appropriate therapy for patients > than 18 years of age with GERD and show symptoms of heartburn, regurgitation, or both for 6 months or more, have been partially or completely responsive to anti-secretory pharmacologic therapy, and have declined laparoscopic fundoplication (Quality of Evidence: ++++). GRADE Recommendation: Strong).

The SAGES Technology and Value Assessment Committee (TVAC) updated its safety and effectiveness analysis of the LINX Reflux Management System.

- Review of published studies suggests that magnetic sphincter augmentation is safe with no reported deaths and a 0.1% rate of intra/perioperative complications.
- Long-term efficacy of LINX appears good for typical GERD symptoms with reduced acid exposure, improved GERD symptoms, and freedom from PPI in 85-88% at three to five years.
- Dysphagia resolves in most patients and the incidence is roughly 10% at one year and 4% at three years. The need for endoscopic dilation ranges from 6-12% and the primary reason for explantation appears to be persistent dysphagia with a rate in larger series from 3-6%.
- Erosion appears to be rare, with one case reported in the first 1,000 patients, one additional published case report, a large series reporting two erosions, and several additional reports in the FDA MAUDE dataset (true number unknown, as multiple entries in this dataset may be made for each patient). Based on very limited literature, erosion can be successfully treated with explantation (Telem et al., 2017).

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Several endoscopic antireflux (endoluminal) devices have received approval by the FDA for treatment of gastroesophageal reflux disease (GERD).

The Stretta System (Mederi Therapeutics) was approved in April 2000 for radiofrequency thermal ablation treatment of GERD. Additional information is available at: http://www.accessdata.fda.gov/cdrh_docs/pdf10/k103017.pdf. (Accessed July 18, 2024)

The Bard EndoCinch Endoscopic Suturing System (Bard Endoscopic Technologies, Billerica, MA, a subsidiary of C.R. Bard Inc), was approved in January 2001 for endoscopic suturing in the treatment of GERD. Subsequent FDA approval was received in September 2007 for an updated version. Additional information is available at: http://www.accessdata.fda.gov/cdrh_docs/pdf7/k071651.pdf. (Accessed July 18, 2024)

The NDO Surgical Endoscopic Plication System was approved in September 2007 for endoscopic suturing in the treatment of GERD in patients who require and respond to pharmacological therapy. Additional information is available at: http://www.accessdata.fda.gov/cdrh_docs/pdf7/k071651.pdf. (Accessed July 18, 2024)

The current generation of EsophyX, EsophyX2, was cleared for marketing as substantially equivalent to the original EsophyX system with minor changes in November 2009 under the FDA510(k) process. The original system was cleared for marketing in September 2007 as substantially equivalent to the predicate devices NDO Surgical Endoscopic Plication System, Bard EndoCinch, and EGS StomaphyX Endoluminal Fasteners and Delivery System. According to the approval summary letter, EsophyX2 is indicated for:

- Use in transoral tissue approximation
- Full-thickness plication and ligation in the GI tract
- The treatment of symptomatic chronic gastroesophageal reflux disease in patients who require and respond to pharmacologic therapy
- Narrowing of the gastroesophageal junction
- Reduction of hiatal hernia < 2 cm in patients with symptomatic chronic gastroesophageal reflux disease

Refer to the following websites for more information:

- http://www.accessdata.fda.gov/cdrh_docs/pdf7/K071651.pdf
- <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm?db=PMN&id=k092400>

(Accessed July 18, 2024)

The Medigus Ultrasound Surgical Endostapler (MUSE™ System) received 510K approval on January 15, 2015 for the endoscopic placement of surgical staples in the soft tissue of the esophagus and stomach in order to create anterior partial fundoplication for treatment of symptomatic chronic GERD in patients who require and respond to pharmacological therapy. Refer to the following website for additional information:

https://www.accessdata.fda.gov/cdrh_docs/pdf14/k143634.pdf. (Accessed July 18, 2024)

The EndoFLIP® System is indicated for use in a clinical setting to measure pressure and dimensions in the esophagus, pylorus, and anal sphincters and received 510K approval on April 17, 2017. It is intended to be used as an adjunct procedure to other diagnostic methods as part of a comprehensive evaluation of patients with symptoms consistent with gastrointestinal motility disorders. Refer to the following website for additional information:

https://www.accessdata.fda.gov/cdrh_docs/pdf17/K170833.pdf. (Accessed July 18, 2024)

These products are Class II devices (moderate risk) deemed substantially equivalent to other endoscopic devices utilizing other procedures.

Torax Medical obtained FDA premarket approval (PMA) in March 2012 for the LINX Reflux Management System. Additional approvals for PMA supplements can be found on the FDA website. Refer to the following website for more information using PMA number P100049: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>. (Accessed July 18, 2024)

Durasphere is approved by the U.S. Food and Drug Administration (FDA) as an injectable bulking agent for gastro-urology use in the treatment of adult women with stress urinary incontinence due to intrinsic sphincter deficiency. Use of this product for esophageal reflux would be considered off-label use. Refer to the following website for more information, using PMA number P980053: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>. (Accessed July 18, 2024)

References

Abdelfatah MM, Noll A, Kapil N, et al. Long-term Outcome of Gastric Per-Oral Endoscopic Pyloromyotomy in Treatment of Gastroparesis. *Clin Gastroenterol Hepatol*. 2021 Apr;19(4):816-824.

Aiolfi A, Asti E, Bernardi D, et al. Early results of magnetic sphincter augmentation versus fundoplication for gastroesophageal reflux disease: systematic review and meta-analysis. *Int J Surg*. 2018 Apr ;52:82-88.

Al Ghamdi SS, Farha J, Moran RA, et al. Zenker's peroral endoscopic myotomy, or flexible or rigid septotomy for Zenker's diverticulum: a multicenter retrospective comparison. *Endoscopy*. 2022 Apr;54(4):345-351.

Albers DV, Kondo A, Bernardo WM. Endoscopic versus surgical approach in the treatment of Zenker's diverticulum: systematic review and meta-analysis. *Endosc Int Open*. 2016 Jun;4(6):E678-86.

Alicuben ET, Bell RCW, Jobe BA, et al. Worldwide experience with erosion of the magnetic sphincter augmentation device. *J Gastrointest Surg*. 2018 Aug;22(8):1442-1447.

American Society for Gastrointestinal Endoscopy (ASGE). The role of endoscopy in the management of GERD. *Gastroenterology* 2015 June; 81(6):1305-1310.

American Society of General Surgeons (ASGS). American Society of General Surgeons (ASGS) LINX statement of support from ASGS. 2014. <https://theasgs.org/position-statements/linx-statement-of-support-from-asgs/>. Accessed July 18, 2024.

Antoniou SA, Koch OO, Kaindlstorfer A, et al. Endoscopic full-thickness plication versus laparoscopic fundoplication: a prospective study on quality of life and symptom control. *Surg Endosc*. 2012 Apr;26(4):1063-8.

Arts J, Bisschops R, Blondeau K, et al. A double-blind sham-controlled study of the effect of radiofrequency energy on symptoms and distensibility of the gastro-esophageal junction in GERD. *Am J Gastroenterol*. 2012 Feb;107(2):222-30.

Asti E, Gianluca Bonitta G, Lovece A, et al. Longitudinal comparison of quality of life in patients undergoing laparoscopic Toupet fundoplication versus magnetic sphincter augmentation. *Observational cohort study with propensity score analysis. Medicine (Baltimore)*. 2016 Jul; 95(30): e4366.

Auyang ED, Carter P, Rauth T, et al.; SAGES Guidelines Committee. SAGES clinical spotlight review: endoluminal treatments for gastroesophageal reflux disease (GERD). *Surg Endosc*. 2013 Aug;27(8):2658-72.

Aziz AM, El-Khayat HR, Sadek A, et al. A prospective randomized trial of sham, single-dose Stretta, and double-dose Stretta for the treatment of gastroesophageal reflux disease. *Surg Endosc*. 2010 Apr;24(4):818-25.

Aziz M, Gangwani MK, Hagbini H, et al. Gastric peroral endoscopic myotomy versus surgical phloromyotomy/pyloroplasty for refractory gastroparesis: Systematic review and meta-analysis. *Endosc Int Open*. 2023;11(4):E322-E329.

Baret F, Jacques J, Pioche M, et al. Evaluation of the safety profile of endoscopic pyloromyotomy by G-POEM: a French multicenter study. *Therap Adv Gastroenterol*. 2022 Oct 3;15:17562848221122472.

Bell R, Lipham J, Louie B, et al. Laparoscopic magnetic sphincter augmentation versus double-dose proton pump inhibitors for management of moderate-to-severe regurgitation in GERD: a randomized controlled trial. *Gastrointest Endosc*. 2019 Jan;89(1):14-22.e1.

Bell R, Lipham J, Louie BE, et al. Magnetic sphincter augmentation superior to proton pump inhibitors for regurgitation in a 1-year randomized trial. *Clin Gastroenterol Hepatol*. 2020 Jul;18(8):1736-1743.e2.

Birk J, Pruitt R, Haber G, et al. The Plicator procedure for the treatment of gastroesophageal reflux disease: a registry study. *Surg Endosc*. 2009 Feb;23(2):423-31.

Bonavina L, DeMeester T, Fockens P, et al. Laparoscopic sphincter augmentation device eliminates reflux symptoms and normalizes esophageal acid exposure: one- and 2-year results of a feasibility trial. *Ann Surg*. 2010 Nov;252(5):857-62.

Bonavina L, Saino GI, Bona D, et al. Magnetic augmentation of the lower esophageal sphincter: results of a feasibility clinical trial. *J Gastrointest Surg*. 2008 Dec;12(12):2133-40.

Budnicka A, Januszewicz W, Białek AB. Peroral endoscopic myotomy in the management of zenker's diverticulum: A retrospective multicenter study. *J Clin Med*. 2021 Jan 7;10(2):187.

Camilleri M, Kuo B, Nguyen L, et al. ACG Clinical Guideline: Gastroparesis. *Am J Gastroenterol*. 2022 Aug 1;117(8):1197-1220.

Chandan S, Mohan BP, Khan SR, et al. Clinical efficacy and safety of magnetic sphincter augmentation (MSA) and transoral incisionless fundoplication (TIF2) in refractory gastroesophageal reflux disease (GERD): a systematic review and meta-analysis. *Endosc Int Open*. 2021 Apr;9(4):E583-E598.

Chandan S, Mohan BP, Chandan OC, et al. Clinical efficacy of per-oral endoscopic myotomy (POEM) for spastic esophageal disorders: a systematic review and meta-analysis. *Surg Endosc*. 2020 Feb;34(2):707-718.

Chen, D, Barber, C, McLoughlin, P, et al. Systematic review of endoscopic treatments for gastroesophageal reflux disease. *Br J Surg*. 2009;96(2):128-136.

Coron E, Sebillé V, Cadiot G, et al. Clinical trial: Radiofrequency energy delivery in proton pump inhibitor-dependent gastro-oesophageal reflux disease patients. *Aliment Pharmacol Ther*. 2008;28 (9):1147-1158.

Chung CS, Huang TY, Lin CL, et al. Efficacy and safety of gastric peroral endoscopic myotomy (G-POEM) for refractory gastroparesis: 3-year follow up results. *J Formos Med Assoc*. 2022 Jul;121(7):1334-1341.

de Moura EGH, Sallum RAA, Nasi A, et al. Endoscopic polymer injection and endoluminal plication in treatment of gastroesophageal reflux disease: evaluation of long-term results. *Endosc Int Open*. 2018 May;6(5):E630-E636.

de Moura ETH, Jukemura J, Ribeiro IB, et al. Peroral endoscopic myotomy vs laparoscopic myotomy and partial fundoplication for esophageal achalasia: A single-center randomized controlled trial. *World J Gastroenterol*. 2022 Sep 7;28(33):4875-4889.

Desart K, Rossidis G, Michel M, et al. Gastroesophageal reflux management with the LINX® System for gastroesophageal reflux disease following laparoscopic sleeve gastrectomy. *J Gastrointest Surg*. 2015 Oct;19(10):1782-6.

Desprez C, Roman S, Leroi AM, et al. The use of impedance planimetry (Endoscopic Functional Lumen Imaging Probe, EndoFLIP®) in the gastrointestinal tract: A systematic review. *Neurogastroenterol Motil*. 2020 Sep;32(9):e13980.

ECRI Institute. Clinical Evidence Assessment. EndoFLIP Impedance Planimetry System (Medtronic plc.) for Identifying Gastrointestinal Motility Disorders. Plymouth Meeting, PA. October 2018. Updated June 2020.

ECRI Institute. Clinical Evidence Assessment. LINX Reflux Management System (Ethicon US, LLC) for Treating Gastroesophageal Reflux Disease. Plymouth Meeting, PA. March 2020. Updated May 2023.

Fass R, Cahn F, Scotti DJ, et al. Systematic review and meta-analysis of controlled and prospective cohort efficacy studies of endoscopic radiofrequency for treatment of gastroesophageal reflux disease. *Surg Endosc*. 2017 Dec;31(12):4865-4882.

Ferrari D, Asti E, Lazzari V, et al. Six to 12-year outcomes of magnetic sphincter augmentation for gastroesophageal reflux disease. *Sci Rep*. 2020 Aug 13;10(1):13753.

Ganz RA, Edmundowicz SA, Taiganides PA, et al. Long-term outcomes of patients receiving a magnetic sphincter augmentation device for gastroesophageal reflux. *Clin Gastroenterol Hepatol*. 2016 May;14(5):671-7.

Ganz RA, Fallon E, Wittchow T, Klein D. A new injectable agent for the treatment of GERD: results of the Durasphere pilot trial. *Gastrointest Endosc*. 2009 Feb;69(2):318-23.

Ganz RA, Peters JH, Horgan S, et al. Esophageal sphincter device for gastroesophageal reflux disease. *NEJM*. 2013; 368(8): 719-727.

Hayes, Inc. Comparative Effectiveness Review. Comparative Effectiveness Review Laparoscopic Surgery for Gastroesophageal Reflux Disease to Medical Management. Hayes Inc; Lansdale, PA: December 2018; archived May 2023.

Hayes, Inc. Health Technology Assessment. Comparative Effectiveness Review of Endoscopic Therapy for Gastroesophageal Reflux Disease. Hayes, Inc; Lansdale, PA: December 2017; archived January 2023.

Hayes, Inc. Health Technology Assessment. Endolumenal Functional Lumen Image Probe (EndoFLIP; Medtronic Inc.) for Evaluation of Esophageal and Gastric Disorders. Hayes, Inc; Lansdale, PA: June 2019; updated June 2022.

Hayes, Inc. Health Technology Assessment. Magnetic Sphincter Augmentation (LINX Reflux Management System) for Treatment of Gastroesophageal Reflux Disease. Hayes Inc; Lansdale, PA. December 2018; updated August 2022.

Hayes, Inc. Health Technology Assessment. Peroral Endoscopic Myotomy for Treatment of Esophageal Achalasia. Hayes, Inc; Lansdale, PA: December 2019; updated March 2023.

Hayes, Inc. Health Technology Assessment. Stretta (Respiratory Technology Corp.) Radiofrequency Treatment for Gastroesophageal Reflux Disease. Hayes, Inc.; May 5, 2023.

He C, Li M, Lu B, et al. Long-term efficacy of peroral endoscopic myotomy for patients with achalasia: Outcomes with a median follow-up of 36 months. *Dig Dis Sci*. 2019;64(3):803-810.

Huang Z, Cui Y, Li Y, et al. Peroral endoscopic myotomy for patients with achalasia with previous Heller myotomy: a systematic review and meta-analysis. *Gastrointest Endosc*. 2021 Jan;93(1):47-56.e5.

Kahrilas PJ, Katzka D, Richter JE. Clinical Practice Update: The Use of Per-Oral Endoscopic Myotomy in Achalasia: Expert Review and Best Practice Advice from the AGA Institute. *Gastroenterology*. 2017;153(5):1205-1211.

Kalapala R, Karyampudi A, Nabi Z, et al. Endoscopic full-thickness plication for the treatment of PPI-dependent GERD: results from a randomised, sham controlled trial. *Gut*. 2022 Apr;71(4):686-694.

Kalapala R, Shah H, Nabi Z, et al. Treatment of gastroesophageal reflux disease using radiofrequency ablation (Stretta procedure): an interim analysis of a randomized trial. *Indian J Gastroenterol*. 2017 Sep;36(5):337-342.

Katz PO, Dunbar KB, Schnoll-Sussman FH, et al. ACG Clinical Guideline for the Diagnosis and Management of Gastroesophageal Reflux Disease. *Am J Gastroenterol*. 2022 Jan 1;117(1):27-56.

Khan MA, Kumbhari V, Ngamruengphong S, et al. Is POEM the answer for management of spastic esophageal disorders? A systematic review and meta-analysis. *Dig Dis Sci*. 2017 Jan;62(1):35-44.

Khashab MA, Familiari P, Draganov PV, et al. Peroral endoscopic myotomy is effective and safe in non-achalasia esophageal motility disorders: an international multicenter study. *Endosc Int Open*. 2018 Aug;6(8):E1031-E1036.

Khashab MA, Vela MF, Thosani N, et al. ASGE guideline on the management of achalasia. *Gastrointest Endosc*. 2020;91(2):213-227.e6.

Khashab MA, Wang AY, Cai Q. AGA Clinical Practice Update on Gastric Peroral Endoscopic Myotomy for Gastroparesis: Commentary. *Gastroenterology*. 2023 Jun;164(7):1329-1335.e1.

Kim HJ, Kwon CI, Kessler WR, et al. Long-term follow-up results of endoscopic treatment of gastroesophageal reflux disease with the MUSE™ endoscopic stapling device. *Surg Endosc*. 2016 Aug;30(8):3402-8.

Kohn GP. Peroral endoscopic myotomy for achalasia—a review. *Ann Laparosc Endosc Surg* 2019; 4:89.

Kohn GP, Dirks RC, Ansari MT et al. SAGES guidelines for the use of peroral endoscopic myotomy (POEM) for the treatment of achalasia. *Surg Endosc*. 2021 May;35(5):1931-1948.

Labonde A, Lades G, Debourdeau A, et al. Gastric peroral endoscopic myotomy in refractory gastroparesis: long-term outcomes and predictive score to improve patient selection. *Gastrointest Endosc*. 2022 Sep;96(3):500-508.e2.

Lipham JC, Demeester TR, Ganz RA, et al. The LINX® reflux management system: confirmed safety and efficacy now at 4 years. *Surg Endosc*. 2012 Apr 27.

Lipham JC, Taiganides PA, Louie BE, et al. Safety analysis of first 1000 patients treated with magnetic sphincter augmentation for gastroesophageal reflux disease. *Dis Esophagus*. 2015;28(4):305-311.

Li H, Peng W, Huang S, et al. The 2 years' long-term efficacy and safety of peroral endoscopic myotomy for the treatment of achalasia: a systematic review. *J Cardiothorac Surg*. 2019;14(1):1. Published 2019 Jan 3.

Louie BE, Smith CD, Smith CC, et al. Objective evidence of reflux control after magnetic sphincter augmentation: one year results from a post approval study. *Ann Surg*. 2019 Aug;270(2):302-308.

Marano L, Pallabazzer G, Solito B, et al. Surgery or peroral esophageal myotomy for achalasia: A systematic review and meta-analysis. *Medicine (Baltimore)*. 2016;95(10):e3001.

Martinek J, Hustak R, Mares J, et al. Endoscopic pyloromyotomy for the treatment of severe and refractory gastroparesis: a pilot, randomised, sham-controlled trial. *Gut*. 2022 Nov;71(11):2170-2178.

Medtronic. EndoFLIP™. Available at: <https://www.medtronic.com/covidien/en-us/products/motility-testing/endoflip-impedance-planimetry-system.html>. Accessed July 18, 2024.

Medigus website. MUSE system. Available at <https://www.attikourismedical.com/product/endoscopy-surgical/surgical/medigus-ultrasonic-surgical-endostapler/>. Accessed July 18, 2024.

Mohan BP, Chandan S, Jha LK, et al. Clinical efficacy of gastric per-oral endoscopic myotomy (G-POEM) in the treatment of refractory gastroparesis and predictors of outcomes: a systematic review and meta-analysis using surgical pyloroplasty as a comparator group. *Surg Endosc*. 2020 Aug;34(8):3352-3367.

National Institute for Health and Care Excellence (NICE). Endoscopic radiofrequency ablation for gastro-oesophageal reflux disease. Interventional procedures guidance [IPG461]. August 2013.

National Institute for Health and Care Excellence (NICE). Laparoscopic insertion of a magnetic ring for gastro-oesophageal reflux disease. Interventional procedures guidance [IPG749]. January 2023.

National Library of Medicine. Diffuse Esophageal Spasm. March 2023. Available at: <https://www.ncbi.nlm.nih.gov/>. Accessed July 18, 2024.

Niu C, Zhang J, Iyer C, et al. Endoscopic full-thickness plication for the treatment of gastroesophageal reflux disease: A systematic review and meta-analysis. *Endosc Int Open*. 2024 Feb 12;12(2):E199-E210.

Noar M, Squires P, Khan S. Radiofrequency energy delivery to the lower esophageal sphincter improves gastroesophageal reflux patient-reported outcomes in failed laparoscopic Nissen fundoplication cohort. *Surg Endosc*. 2017 Jul;31(7):2854-2862.

Noar MD, Squires P, Noar E, et al. Long-term maintenance effect of radiofrequency energy delivery for refractory GERD: a decade later. *Surg Endosc*. 2014 Aug;28(8):2323-33.

North A, Tewari N. Peroral endoscopic myotomy compared to laparoscopic Heller myotomy and pneumatic dilation in the treatment of achalasia: a systematic review. *Dis Esophagus*. 2024 Jan 1;37(1):doad055.

Parmar AD, Tessler RA, Chang HY, et al. Two-stage explantation of a magnetic lower esophageal sphincter augmentation device due to esophageal erosion. *J Laparoendosc Adv Surg Tech A*. 2017 Aug;27(8):829-833.

Rafeeqi T, Salimi-Jazi F, Cunningham A, et al. The utility of Endoscopic Functional Luminal Imaging (EndoFLIP) in the diagnosis and management of children with achalasia. *J Pediatr Surg*. 2023 Apr;58(4):639-642.

Reynolds JL, Zehetner J, Wu P, et al. Laparoscopic magnetic sphincter augmentation vs laparoscopic Nissen fundoplication: A matched-pair analysis of 100 Patients. *J Am Coll Surg*. 2015 Jul;221(1):123-8.

Riegler M, Schoppman SF, Bonavina L, et al. Magnetic sphincter augmentation and fundoplication for GERD in clinical practice: one-year results of a multicenter, prospective observational study. *Surg Endosc*. 2015 May;29(5):1123-9.

Rinsma NF, Farré R, Bouvy ND, et al. The effect of endoscopic fundoplication and proton pump inhibitors on baseline impedance and heartburn severity in GERD patients. *Neurogastroenterol Motil*. 2015 Feb;27(2):220-8.

Rothstein R, Filipi C, Caca K, et al. Endoscopic full-thickness plication for the treatment of gastroesophageal reflux disease: A randomized, sham-controlled trial. *Gastroenterology*. 2006;131(3):704-712.

Saino G, Bonavina L, Lipham JC, et al. Magnetic sphincter augmentation for gastroesophageal reflux at 5 years: final results of a pilot study show long-term acid reduction and symptom improvement. *J Laparoendosc Adv Surg Tech A*. 2015 Oct 1; 25(10): 787-792.

Saleh CMG, Familiari P, Bastiaansen BAJ, et al. The efficacy of peroral endoscopic myotomy vs pneumatic dilation as treatment for patients with achalasia suffering from persistent or recurrent symptoms after laparoscopic heller myotomy: a randomized clinical trial. *Gastroenterology*. 2023 Jun;164(7):1108-1118.e3.

Schizas D, Mastoraki A, Papoutsi E, et al. LINX® reflux management system to bridge the "treatment gap" in gastroesophageal reflux disease: A systematic review of 35 studies. *World J Clin Cases*. 2020;8(2):294-305.

Schwartz MP, Schreinemakers JR, Smout AJ. Four-year follow-up of endoscopic gastroplication for the treatment of gastroesophageal reflux disease. *World J Gastrointest Pharmacol Ther*. 2013 Nov 6;4(4):120-6.

Skubleny D, Switzer NJ, Dang J, et al. LINX® magnetic esophageal sphincter augmentation versus Nissen fundoplication for gastroesophageal reflux disease: a systematic review and meta-analysis. *Surg Endosc*. 2017 Aug;31(8):3078-3084.

Smith CD, Ganz RA, Lipham JC, et al. Lower esophageal sphincter augmentation for gastroesophageal reflux disease: the safety of a modern implant. *J Laparoendosc Adv Surg Tech A*. 2017 Jun;27(6):586-591.

Society of American Gastrointestinal and Endoscopic Surgeons (SAGES). Position Statement on Endolumenal Therapies for Gastrointestinal Diseases. 2009. Available at: <https://www.sages.org/publications/guidelines/position-statement-on-endolumenal-therapies-for-gastrointestinal-diseases/>. Accessed August 18, 2024.

Stojilkovic T, Staudinger K, Dennis J. A systematic review of the long-term clinical success of gastric peroral endoscopic myotomy for refractory gastroparesis. *Cureus*. 2023 May 30;15(5):e39709.

Telem D, Wright A, Shah P, et al. SAGES Technology and Value Assessment Committee (TAVAC) safety and effectiveness analysis. LINX® Reflux Management System (Torax Medical, Inc.). March 2017. Available at: <https://www.sages.org/publications/tavac/tavac-safety-and-effectiveness-analysis-linx-reflux-management-system/>. Accessed July 18, 2024.

Valinoti AC, Angeramo CA, Dreifuss N, et al. Magnetic sphincter augmentation device for gastroesophageal reflux disease: effective, but postoperative dysphagia and risk of erosion should not be underestimated. a systematic review and meta-analysis. *Arq Bras Cir Dig*. 2024 Mar 4;36:e1781.

Viswanath Y, Maguire N, Obuobi RB, et al. Endoscopic day case antireflux radiofrequency (Stretta) therapy improves quality of life and reduce proton pump inhibitor (PPI) dependency in patients with gastro-oesophageal reflux disease: a prospective study from a UK tertiary centre. *Frontline Gastroenterol*. 2019 Apr;10(2):113-119.

von Renteln D, Schiefke I, Fuchs KH, et al. Endoscopic full-thickness plication for the treatment of gastroesophageal reflux disease using multiple Plicator implants: 12-month multicenter study results. *Surg Endosc*. 2009 Aug;23(8):1866-75.

Warren HF, Brown LM, Mihura M, et al. Factors influencing the outcome of magnetic sphincter augmentation for chronic gastroesophageal reflux disease. *Surg Endosc*. 2018 Jan;32(1):405-412.

Warren HF, Reynolds JL, Lipham JC, et al. Multi-institutional outcomes using magnetic sphincter augmentation versus Nissen fundoplication for chronic gastroesophageal reflux disease. *Surg Endosc*. 2016 Aug;30(8):3289-96. Epub 2015 Nov 5.

Weitzendorfer M, Spaun GO, Antoniou SA, et al. Clinical feasibility of a new full-thickness endoscopic plication device (GERDx™) for patients with GERD: results of a prospective trial. *Surg Endosc*. 2018 May;32(5):2541-2549.

Werner YB, Hakanson B, Martinek J, et al. Endoscopic or surgical myotomy in patients with idiopathic achalasia. *N Engl J Med*. 2019;381(23):2219-2229.

Yadlapati R, Gyawali CP, Pandolfino JE; et al. AGA clinical practice update on the personalized approach to the evaluation and management of GERD: expert review. *Clin Gastroenterol Hepatol*. 2022 May;20(5):984-994.e1.

Yang J, Novak S, Ujiki M. An international study on the use of peroral endoscopic myotomy in the management of Zenker's diverticulum. *Gastrointest Endosc*. 2020 Jan;91(1):163-168.

Yoo IK, Choi SA, Kim WH, et al. Assessment of clinical outcomes after peroral endoscopic myotomy via esophageal distensibility measurements with the endoluminal functional lumen imaging probe. *Gut Liver*. 2019 Jan 15;13(1):32-39.

Zacherl J, Roy-Shapira A, Bonavina L, et al. Endoscopic anterior fundoplication with the Medigus Ultrasonic Surgical Endostapler (MUSE™) for gastroesophageal reflux disease: 6-month results from a multi-center prospective trial. *Surg Endosc*. 2015 Jan;29(1):220-9.

Zerbib F, Sacher-Huvelin S, Coron E, et al. Randomised clinical trial: oesophageal radiofrequency energy delivery versus sham for PPI-refractory heartburn. *Aliment Pharmacol Ther*. 2020 Aug;52(4):637-645.

Zhang H, Huang S, Xia H, et al. The role of peroral endoscopic myotomy for Zenker's diverticulum: a systematic review and meta-analysis. *Surg Endosc*. 2022 May;36(5):2749-2759.

Zhang H, Zhang J, Jiang A, et al. Gastric peroral endoscopic myotomy for gastroparesis: A systematic review of efficacy and safety. *Gastroenterol Hepatol*. 2019 Aug-Sep;42(7):413-422. English, Spanish.

Zhuang QJ, Tan ND, Chen SF, et al. Magnetic sphincter augmentation in treating refractory gastroesophageal reflux disease: A systematic review and meta-analysis. *J Dig Dis*. 2021 Dec;22(12):695-70.

Policy History/Revision Information

Date	Summary of Changes
07/01/2025	<p>Coverage Rationale</p> <ul style="list-style-type: none">Added language to clarify:<ul style="list-style-type: none">The per oral endoscopic myotomy (POEM) procedure is proven and medically necessary for <i>treating individuals with</i> Achalasia or Diffuse Esophageal SpasmPer oral endoscopic myotomy (POEM) is unproven and not medically necessary for all other indications (e.g., Zenker's diverticula) due to insufficient evidence <i>of efficacy</i> <p>Supporting Information</p> <ul style="list-style-type: none">Updated <i>Clinical Evidence</i> and <i>References</i> sections to reflect the most current informationArchived previous policy version CS079KY.10

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state, or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state, or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state, or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state, or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare uses InterQual® for the primary medical/surgical criteria, and the American Society of Addiction Medicine (ASAM) for substance use, in administering health benefits. If InterQual® does not have applicable criteria, UnitedHealthcare may also use UnitedHealthcare Medical Policies, Coverage Determination Guidelines, and/or Utilization Review Guidelines that have been approved by the Kentucky Department for Medicaid Services. The UnitedHealthcare Medical Policies, Coverage Determination Guidelines, and Utilization Review Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.