

Lower Extremity Invasive Diagnostic and Endovascular Procedures

Policy Number: CS166.G
Effective Date: August 1, 2021

[➔ Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	2
Applicable Codes	3
Description of Services	4
Clinical Evidence	5
U.S. Food and Drug Administration	7
References	7
Policy History/Revision Information	8
Instructions for Use	9

<p>Related Community Plan Policies</p> <ul style="list-style-type: none"> Pneumatic Compression Devices Surgical and Ablative Procedures for Venous Insufficiency and Varicose Veins
<p>Commercial Policy</p> <ul style="list-style-type: none"> Lower Extremity Invasive Diagnostic and Endovascular Procedures
<p>Medicare Advantage Coverage Summary</p> <ul style="list-style-type: none"> Cardiovascular Diagnostic Procedures

Application

This Medical Policy does not apply to the states listed below; refer to the state-specific policy/guideline, if noted:

State	Policy/Guideline
Indiana	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for Indiana Only)
Kentucky	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for Kentucky Only)
Louisiana	None
Mississippi	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for Mississippi Only)
Nebraska	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for Nebraska Only)
New Jersey	Lower Extremity Vascular Angiography (for New Jersey Only)
North Carolina	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for North Carolina Only)
Pennsylvania	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for Pennsylvania Only)
Tennessee	Lower Extremity Invasive Diagnostic and Endovascular Procedures (for Tennessee Only)

Coverage Rationale

Note: This policy does not apply to upper extremities.

Diagnosis

Lower extremity vascular angiography is proven and medically necessary for evaluating arterial disease, aneurysms, perivascular masses, and trauma related injuries of the lower extremity under certain circumstances.

Angiography for known chronic lower extremity arterial disease (peripheral vascular disease, occlusion or stenosis of arteries of the leg, claudication and limb ischemia) requires all of the following:

- Abnormal ankle-brachial index (ABI) or an abnormal computed tomography angiography (CTA), magnetic resonance angiography (MRA) or duplex ultrasound; and
- One of the following:
 - Claudication – as evidenced with an ABI \leq 0.90; or
 - Limb Ischemia – as evidenced with an ABI $<$ 0.40

For additional medical necessity clinical coverage criteria, refer to the InterQual® 2021, Apr. 2021 Release, CP: Imaging, Imaging, Peripheral Vascular for:

- Angiogram, X-ray Extremity, Unilateral
- Angiogram, X-ray Extremity, Bilateral

Click [here](#) to view the InterQual® criteria.

Treatment

Endovascular revascularization procedures (e.g., stents, angioplasty and/or atherectomy) for treating lower extremity ischemia are proven and medically necessary in individuals who meet the following indication-specific criteria:

- Claudication due to atherosclerotic disease of the aortoiliac and/or femoropopliteal arteries when all of the following criteria are met:
 - Impaired ability to work and/or perform activities of daily living (ADL)
 - All of the following conservative therapies have been tried and failed:
 - At least twelve (12) weeks of a [Supervised](#) or [Structured](#) Exercise Program
 - Pharmacologic therapy
 - Smoking cessation, if applicable
 - Moderate to severe ischemic peripheral artery disease with Ankle-Brachial Index (ABI) \leq 0.69
 - Imaging results show anatomic location and severity of occlusion (stenosis \geq 50%) (e.g., duplex ultrasound, CTA, MRA or invasive angiography). If duplex ultrasound does not demonstrate a stenosis \geq 50%, another imaging modality will be necessary to demonstrate the extent of stenosis.
- Chronic Limb-Threatening Ischemia (CLTI) when all of the following criteria are met:
 - One or more of the following:
 - Pain at rest
 - Nonhealing wound or ulcer due to ischemia
 - Gangrene
 - Moderate to severe ischemic peripheral artery disease and any of the following:
 - Ankle-Brachial Index (ABI) \leq 0.69; or
 - Ankle pressure $<$ 50 mmHg; or
 - Toe-Brachial Index \leq 0.70; or
 - Toe pressure $<$ 30 mmHg; or
 - Transcutaneous Oxygen Pressure (TcPO₂) $<$ 25 mmHg
 - Imaging results show anatomic location and severity of occlusion (stenosis \geq 50%) (e.g., duplex ultrasound, CTA, MRA or invasive angiography). If duplex ultrasound does not demonstrate a stenosis \geq 50%, another imaging modality will be necessary to demonstrate the extent of stenosis.

Due to insufficient evidence of efficacy, endovascular revascularization procedures (e.g., stents, angioplasty and/or atherectomy) for treating lower extremity ischemia are unproven and not medically necessary in the following circumstances:

- Claudication due to isolated infrapopliteal (e.g., anterior tibial, posterior tibial or peroneal) artery disease
- To prevent the progression of Claudication to CLTI
- Individual is asymptomatic
- Treatment of a nonviable limb

Definitions

Ankle-Brachial Index (ABI): The ABI compares the systolic blood pressure in the ankle to the systolic blood pressure in the arm and indicates how well blood is flowing in the limbs (Gerhard-Herman et al., 2017). Studies have shown that ABI can

differentiate between normal and diseased limbs with a sensitivity of 97% and a specificity of 100% and that the resting ABI is a significant predictor of disease severity (Mohler, 2003). An ABI less than 0.90 indicates peripheral artery disease (PAD).

- Mild: 0.70 to 0.89
- Moderate: 0.40 – 0.69
- Severe: Less than 0.40

Chronic Limb-Threatening Ischemia (CLTI): A condition characterized by chronic (≥ 2 weeks) ischemic rest pain, nonhealing wound/ulcers or gangrene in one or both legs attributable to arterial occlusive disease (Gerhard-Herman et al., 2017). Also referred to as critical limb ischemia (CLI).

Claudication: Fatigue, discomfort, cramping or pain of vascular origin in the muscles of the lower extremities that is consistently induced by exercise and consistently relieved by rest (within 10 min) (Gerhard-Herman et al., 2017).

Structured Exercise Program: Components of a Structured Exercise Program include all of the following (Gerhard-Herman, et al., 2017):

- Program takes place in the personal setting of the patient rather than in a clinical setting
- Program is self-directed with guidance of healthcare provider(s)
- Healthcare providers prescribe an exercise regimen similar to that of a [Supervised](#) Program
- Patient counseling ensures understanding of how to begin and maintain the program and how to progress the difficulty of the walking (by increasing distance or speed)
- Program may incorporate behavioral change techniques, such as health coaching or use of activity monitors

Supervised Exercise Program: Components of a Supervised Exercise Program include all of the following (Gerhard-Herman, et al., 2017):

- Program takes place in a hospital or outpatient facility
- Program uses intermittent walking exercise as the treatment modality
- Program can be standalone or within a cardiac rehabilitation program
- Program is directly supervised by [qualified] healthcare provider(s)
- Training is performed for a minimum of 30–45 minutes per session and sessions are performed at least 3 times per week for a minimum of 12 weeks
- Training involves intermittent bouts of walking to moderate-to-maximum claudication, alternating with periods of rest
- Warm-up and cool-down periods precede and follow each session of walking

Toe-Brachial Index (TBI): The TBI compares blood pressure in the toe to blood pressure in the arm and indicates how well blood is flowing in the limbs. The test may be performed when ABI results are abnormally high due to calcified arteries in the legs.

Transcutaneous Oxygen Pressure (TcPO₂): A TcPO₂ test measures the oxygen level of tissue under the skin and may be used as an indirect measure of blood flow.

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
37220	Revascularization, endovascular, open or percutaneous, iliac artery, unilateral, initial vessel; with transluminal angioplasty
37221	Revascularization, endovascular, open or percutaneous, iliac artery, unilateral, initial vessel; with transluminal stent placement(s), includes angioplasty within the same vessel, when performed

CPT Code	Description
37222	Revascularization, endovascular, open or percutaneous, iliac artery, each additional ipsilateral iliac vessel; with transluminal angioplasty (List separately in addition to code for primary procedure)
37223	Revascularization, endovascular, open or percutaneous, iliac artery, each additional ipsilateral iliac vessel; with transluminal stent placement(s), includes angioplasty within the same vessel, when performed (List separately in addition to code for primary procedure)
37224	Revascularization, endovascular, open or percutaneous, femoral, popliteal artery(s), unilateral; with transluminal angioplasty
37225	Revascularization, endovascular, open or percutaneous, femoral, popliteal artery(s), unilateral; with atherectomy, includes angioplasty within the same vessel, when performed
37226	Revascularization, endovascular, open or percutaneous, femoral, popliteal artery(s), unilateral; with transluminal stent placement(s), includes angioplasty within the same vessel, when performed
37227	Revascularization, endovascular, open or percutaneous, femoral, popliteal artery(s), unilateral; with transluminal stent placement(s) and atherectomy, includes angioplasty within the same vessel, when performed
37228	Revascularization, endovascular, open or percutaneous, tibial, peroneal artery, unilateral, initial vessel; with transluminal angioplasty
37229	Revascularization, endovascular, open or percutaneous, tibial, peroneal artery, unilateral, initial vessel; with atherectomy, includes angioplasty within the same vessel, when performed
37230	Revascularization, endovascular, open or percutaneous, tibial, peroneal artery, unilateral, initial vessel; with transluminal stent placement(s), includes angioplasty within the same vessel, when performed
37231	Revascularization, endovascular, open or percutaneous, tibial, peroneal artery, unilateral, initial vessel; with transluminal stent placement(s) and atherectomy, includes angioplasty within the same vessel, when performed
37232	Revascularization, endovascular, open or percutaneous, tibial/peroneal artery, unilateral, each additional vessel; with transluminal angioplasty (List separately in addition to code for primary procedure)
37233	Revascularization, endovascular, open or percutaneous, tibial/peroneal artery, unilateral, each additional vessel; with atherectomy, includes angioplasty within the same vessel, when performed (List separately in addition to code for primary procedure)
37234	Revascularization, endovascular, open or percutaneous, tibial/peroneal artery, unilateral, each additional vessel; with transluminal stent placement(s), includes angioplasty within the same vessel, when performed (List separately in addition to code for primary procedure)
37235	Revascularization, endovascular, open or percutaneous, tibial/peroneal artery, unilateral, each additional vessel; with transluminal stent placement(s) and atherectomy, includes angioplasty within the same vessel, when performed (List separately in addition to code for primary procedure)
75710	Angiography, extremity, unilateral, radiological supervision and interpretation
75716	Angiography, extremity, bilateral, radiological supervision and interpretation

CPT® is a registered trademark of the American Medical Association

Description of Services

Peripheral artery disease (PAD) is a narrowing of vessels due to atherosclerosis that limits blood flow to the limbs. PAD most commonly affects arteries in the legs. While many people with PAD do not have any symptoms, some will have leg pain, numbness or cramping during exercise that is relieved by rest (claudication). Risk factors include age, smoking, diabetes, obesity, high blood pressure and high cholesterol. Some of the tests used to diagnose PAD are physical exam, Ankle-Brachial Index (ABI), ankle pressure, Toe-Brachial Index, toe pressure, Transcutaneous Oxygen Pressure, noninvasive imaging, and invasive angiography.

PAD is associated with an increased risk of heart attack, stroke and when left untreated, can lead to chronic limb-threatening ischemia (CLTI). Treatment options include lifestyle changes, medications, endovascular techniques and surgery. Endovascular

techniques to treat claudication and CLTI include balloon dilation (angioplasty), stents and atherectomy. The technique chosen for endovascular treatment depends on many factors including lesion characteristics such as anatomic location, lesion length and degree of calcification (Gerhard-Herman et al., 2017; National Heart, Lung and Blood Institute website).

Clinical Evidence

A Cochrane systemic review by Fakhry et al. (2018) assessed the effectiveness of endovascular revascularization compared with no specific therapy for intermittent claudication or compared with a conservative therapy option such as supervised exercise or drug therapy. The review included ten studies with a total of 1087 participants. The results showed that endovascular revascularization and supervised exercise are comparable treatment options in improving walking distances and quality of life in individuals with intermittent claudication. Combination therapy (endovascular revascularization with either supervised exercise or drug therapy) seemed to result in greater improvements than those seen with supervised exercise or drug therapy alone. (The ERASE trial by Fakhry et al., 2015 and the CLEVER trial by Murphy et al., 2015, which were previously cited in this policy, are included in this systematic review.)

Malgor et al. (2015) conducted a systematic review to evaluate the efficacy of three treatment strategies for individuals with claudication. Primary outcome measures included mortality, amputation, walking distance, quality of life, patency and measures of blood flow (ABI). The review included eight systematic reviews and 12 trials enrolling 1548 patients. Compared with medical management, each of the three treatments (surgery, endovascular therapy and exercise therapy) was associated with improved walking distance, claudication symptoms and quality of life. Evidence supporting superiority of one of the three approaches was limited. However, blood flow parameters improved faster and better with both forms of revascularization compared with exercise or medical management. Compared with endovascular therapy, open surgery may be associated with longer length of hospital stay and higher complication rates but resulted in more durable patency (moderate-quality evidence). (The CLEVER trial by Murphy et al., 2012, which was previously cited in this policy, is included in this systematic review.)

Vemulapalli et al. (2015) conducted a systematic review and a network meta-analysis to evaluate the comparative effectiveness of medical therapy, supervised exercise training, endovascular intervention and surgical revascularization in patients with claudication. Outcomes assessed included walking distance, claudication distance, all-cause mortality and quality of life. Thirty-five studies (n=7475) were included in the analysis. A meta-analysis of 16 studies suggested that, compared with usual care, maximal walking measures were improved to a greater extent with supervised exercise than with medical therapy or endovascular intervention. A meta-analysis of 12 studies demonstrated that exercise training and endovascular intervention, but not cilostazol, improved initial claudication measures compared with usual care. A meta-analysis of 13 studies suggested that although all treatment modalities were superior to usual care, there was no significant difference between modalities in respect to quality of life. The authors noted that heterogeneity in functional endpoints, single-arm observational study design and poor subgroup reporting significantly limit comparative effectiveness analysis in PAD. Further studies with attention to study design, standardized efficacy and safety endpoints, and appropriate subgroup reporting are needed. (The multicenter CLEVER trial by Murphy et al., 2012, which was previously cited in this policy, is included in this systematic review.)

A National Institute for Health and Care Excellence (NICE) clinical guideline offers recommendations on the diagnosis and management of PAD (NICE, 2012; updated 2020).

Clinical Practice Guidelines

American College of Cardiology (ACC)/American Heart Association (AHA)/Society for Cardiovascular Angiography and Interventions (SCAI)/Society of Interventional Radiology (SIR)/Society for Vascular Medicine (SVM)

In a multi-society report, Bailey et al. (2019) published appropriate use criteria for peripheral artery interventions. The panel recommends that patients with PAD and intermittent claudication should first be treated with guideline-directed medical therapy and structured exercise. Revascularization should be considered only in patients who continue to have lifestyle-limiting claudication despite these noninvasive approaches. In situations where medical therapy is insufficient, the selection of surgical or endovascular revascularization depends on several factors including patient risk level and lesion characteristics, such as anatomic location, length and presence of stenosis or occlusion. For patients with CLTI, both endovascular or surgical revascularization procedures are considered appropriate and critical for the reduction of high morbidity and mortality rates associated with limb loss and cardiovascular events.

American Diabetes Association (ADA)

ADA guidelines address the evaluation of PAD in diabetic individuals. Recommended testing is based on the presence and severity of symptoms. The guidelines state that urgent vascular imaging and revascularization should be considered in individuals with a diabetic foot ulcer and an ankle pressure <50 mmHg, toe pressure <30 mmHg, or a TcPO₂ <25 mmHg (ADA, 2021).

American Heart Association (AHA)/American College of Cardiology (ACC)

AHA/ACC guidelines for the diagnosis and management of patients with lower extremity PAD make the following recommendations for diagnostic imaging procedures (Gerhard-Herman et al., 2017):

- Duplex ultrasound, CTA, or MRA of the lower extremities is useful to diagnose anatomic location and severity of stenosis for patients with symptomatic PAD in whom revascularization is considered
- Invasive angiography is useful for patients with CLTI in whom revascularization is considered
- Invasive angiography is reasonable for patients with lifestyle-limiting claudication with an inadequate response to guideline-directed management and therapy for whom revascularization is considered
- Invasive and noninvasive angiography (i.e., CTA, MRA) should not be performed for the anatomic assessment of patients with asymptomatic PAD

AHA/ACC guidelines for the diagnosis and management of patients with lower extremity PAD make the following recommendations for surgical and endovascular revascularization procedures (Gerhard-Herman et al., 2017):

Claudication

- Revascularization is a reasonable treatment option for the patient with lifestyle-limiting claudication with an inadequate response to guideline-directed management and therapy
- Endovascular procedures are effective as a revascularization option for patients with lifestyle-limiting claudication and hemodynamically significant aortoiliac occlusive disease
- Endovascular procedures are reasonable as a revascularization option for patients with lifestyle-limiting claudication and hemodynamically significant femoropopliteal disease
- The usefulness of endovascular procedures as a revascularization option for patients with claudication due to isolated infrapopliteal artery disease is unknown
- Endovascular procedures should not be performed in patients with PAD solely to prevent progression to CLTI

CLTI

- In patients with CLTI, revascularization should be performed when possible to minimize tissue loss
- An evaluation for revascularization options should be performed by an interdisciplinary care team before amputation in the patient with CLTI
- Endovascular procedures are recommended to establish in-line blood flow to the foot in patients with nonhealing wounds or gangrene
- A staged approach to endovascular procedures is reasonable in patients with ischemic rest pain
- Evaluation of lesion characteristics can be useful in selecting the endovascular approach for CLTI

International Working Group on the Diabetic Foot (IWGDF)

IWGDF guidelines on the prevention and management of diabetic foot disease state that in patients with either an ankle pressure <50mm Hg or an ABI <0.5, consider urgent vascular imaging and, when findings suggest it is appropriate, revascularization. Also consider revascularization if the toe pressure is <30 mmHg or TcPO₂ is <25 mmHg. Clinicians might consider revascularization at higher pressure levels in patients with extensive tissue loss or infection (Schaper et al., 2020).

Society for Vascular Surgery (SVS)

SVS guidelines provide a comprehensive set of recommendations for the evaluation and management of CLTI. Vein bypass may be preferred for average-risk patients with advanced limb threat and high complexity disease, while those with less complex anatomy, intermediate severity limb threat or high patient risk may be favored for endovascular intervention. All patients with CLTI should be afforded best medical therapy including the use of antithrombotic, lipid-lowering, antihypertensive

and glycemic control agents, as well as counseling on smoking cessation, diet, exercise and preventive foot care (Conte et al., 2019).

Separate SVS guidelines provide a comprehensive set of recommendations for the evaluation and management of asymptomatic disease and intermittent claudication. Emphasis is placed on risk factor modification, medical therapies and broader use of exercise programs to improve cardiovascular health and functional performance. Revascularization for intermittent claudication is an appropriate therapy for selected patients with disabling symptoms, after a careful risk-benefit analysis. Treatment should be individualized based on comorbid conditions, degree of functional impairment and anatomic factors. Invasive treatments for intermittent claudication should provide predictable functional improvements with reasonable durability. A minimum threshold of a >50% likelihood of sustained efficacy for at least 2 years is suggested as a benchmark. Endovascular approaches are favored for most candidates with aortoiliac disease and for selected patients with femoropopliteal disease in whom anatomic durability is expected to meet this minimum threshold. Conversely, caution is warranted in the use of interventions for intermittent claudication in anatomic settings where durability is limited (extensive calcification, small-caliber arteries, diffuse infrainguinal disease, poor runoff). Surgical bypass may be a preferred strategy in good-risk patients with these disease patterns or in those with prior endovascular failures. Common femoral artery disease should be treated surgically, and the saphenous vein is the preferred conduit for infrainguinal bypass grafting. Patients who undergo invasive treatments for intermittent claudication should be monitored regularly in a surveillance program to record subjective improvements, assess risk factors, optimize compliance with cardioprotective medications and monitor hemodynamic and patency status (Conte et al., 2015).

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

The FDA has approved several stents and stent systems for the treatment of PAD of the lower extremities. See the following website (use product codes NIO and NIP) for more information:

<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm>. (Accessed April 21, 2021)

The FDA has approved several catheter systems used for the treatment of PAD of the lower extremities. See the following website (use product code DQY) for more information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed April 21, 2021)

References

Ahmed O, Hanley M, Bennett SJ, et al.; Expert Panel on Vascular Imaging. ACR Appropriateness Criteria[®] Vascular claudication—assessment for revascularization. *J Am Coll Radiol*. 2017 May;14(5S):S372-S379.

American Diabetes Association. Standards of medical care in diabetes - 2021. Available at:

https://care.diabetesjournals.org/content/diacare/suppl/2020/12/09/44.Supplement_1.DC1/DC_44_S1_final_copyright_stamped.pdf. Accessed April 21, 2021.

Bailey SR, Beckman JA, Dao TD, et al. ACC/AHA/SCAI/SIR/SVM 2018 Appropriate use criteria for peripheral artery intervention: a report of the American College of Cardiology Appropriate Use Criteria Task Force, American Heart Association, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, and Society for Vascular Medicine. *J Am Coll Cardiol*. 2019 Jan 22;73(2):214-237.

Conte MS, Bradbury AW, Kolh P, et al.; GVG Writing Group. Global vascular guidelines on the management of chronic limb-threatening ischemia. *J Vasc Surg*. 2019 Jun;69(6S):3S-125S.e40. Erratum in: *J Vasc Surg*. 2019 Aug;70(2):662.

Conte MS, Pomposelli FB, Clair DG, et al.; Society for Vascular Surgery Lower Extremity Guidelines Writing Group. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: management of asymptomatic disease and claudication. *J Vasc Surg*. 2015 Mar;61(3 Suppl):2S-41S. Erratum in: *J Vasc Surg*. 2015 May;61(5):1382.

Fakhry F, Fokkenrood HJP, Spronk S, et al. Endovascular revascularisation versus conservative management for intermittent claudication. *Cochrane Database of Systematic Reviews* 2018, Issue 3. Art. No.: CD010512.

Fakhry F, Spronk S, van der Laan L, et al. Endovascular revascularization and supervised exercise for peripheral artery disease and intermittent claudication: a randomized clinical trial. *JAMA*. 2015 Nov 10;314(18):1936-44.

Francois CJ, Skulborstad EP, Kalva SP, et al.; Expert Panel on Vascular Imaging. ACR Appropriateness Criteria® Nonatherosclerotic peripheral arterial disease. *J Am Coll Radiol*. 2019 May;16(5S):S174-S183.

Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the management of patients with lower extremity peripheral artery disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2017 Mar 21;69(11):e71-e126. Erratum in: *J Am Coll Cardiol*. 2017 Mar 21;69(11):1521.

Malgor RD, Alahdab F, Elraiyah TA, et al. A systematic review of treatment of intermittent claudication in the lower extremities. *J Vasc Surg*. 2015 Mar;61(3 Suppl):54S-73S. Erratum in: *J Vasc Surg*. 2015 May;61(5):1382.

Misra S, Shishehbor MH, Takahashi EA, et al.; American Heart Association Council on Peripheral Vascular Disease; Council on Clinical Cardiology; and Council on Cardiovascular and Stroke Nursing. Perfusion assessment in critical limb ischemia: principles for understanding and the development of evidence and evaluation of devices: A Scientific Statement from the American Heart Association. *Circulation*. 2019 Sep 17;140(12):e657-e672.

Mohler ER 3rd. Peripheral arterial disease: identification and implications. *Arch Intern Med*. 2003 Oct 27;163(19):2306-14.

Murphy TP, Cutlip DE, Regensteiner JG, et al.; CLEVER Study Investigators. Supervised exercise versus primary stenting for claudication resulting from aortoiliac peripheral artery disease: six-month outcomes from the claudication: exercise versus endoluminal revascularization (CLEVER) study. *Circulation*. 2012 Jan 3;125(1):130-9.

Murphy TP, Cutlip DE, Regensteiner JG, et al. Supervised exercise, stent revascularization, or medical therapy for claudication due to aortoiliac peripheral artery disease: the CLEVER study. *J Am Coll Cardiol*. 2015 Mar 17;65(10):999-1009. Erratum in: *J Am Coll Cardiol*. 2015 May 12;65(18):2055.

National Heart, Lung and Blood Institute (NHLBI) website. Peripheral artery disease. Updated September 2020. <https://www.nhlbi.nih.gov/health-topics/peripheral-artery-disease>. Accessed April 21, 2021.

National Institute for Health and Care Excellence (NICE). CG147. Peripheral arterial disease: diagnosis and management. August 2012. Updated December 2020.

Schaper NC, van Netten JJ, Apelqvist J, et al.; IWGDF Editorial Board. Practical guidelines on the prevention and management of diabetic foot disease (IWGDF 2019 update). *Diabetes Metab Res Rev*. 2020 Mar;36 Suppl 1:e3266.

Vemulapalli S, Dolor RJ, Hasselblad V, et al. Comparative effectiveness of medical therapy, supervised exercise, and revascularization for patients with intermittent claudication: a network meta-analysis. *Clin Cardiol*. 2015 Jun;38(6):378-86.

Policy History/Revision Information

Date	Summary of Changes
08/01/2021	<p>Application</p> <ul style="list-style-type: none"> Added language to indicate this policy does not apply to the states of Mississippi and Pennsylvania; refer to the state-specific policy version
08/01/2021	<p>Coverage Rationale</p> <p><i>Diagnosis</i></p> <ul style="list-style-type: none"> Replaced language indicating: <ul style="list-style-type: none"> “Lower extremity vascular angiography is proven and medically necessary for evaluating arterial disease, aneurysms, and trauma related injuries of the lower extremity” with “lower extremity vascular angiography is proven and medically necessary for evaluating arterial disease, aneurysms, <i>perivascular masses</i>, and trauma related injuries of the lower extremity <i>under certain circumstances</i>” “Angiography for known lower extremity arterial disease (peripheral vascular disease, occlusion or stenosis of arteries of the leg, claudication and limb ischemia) requires all of the [listed criteria]” with “angiography for known <i>chronic</i> lower extremity arterial disease (peripheral vascular disease, occlusion or stenosis of arteries of the leg, claudication and limb ischemia) requires all of the [listed criteria]” Removed language indicating angiography [is proven and medically necessary] for trauma related injuries including, but not limited to, one of the following findings:

Date	Summary of Changes
	<ul style="list-style-type: none"> ○ Clinically significant hematoma ○ Knee dislocation (confirmed by x-ray) ○ Need for localization of the source of bleeding and vascular injury found on duplex ultrasound ● Updated language pertaining to <i>additional</i> medical necessity coverage criteria; replaced reference to “InterQual® 2020” with “InterQual® 2021” <p>Treatment</p> <ul style="list-style-type: none"> ● Revised coverage criteria; added criterion requiring “if duplex ultrasound does not demonstrate a stenosis ≥ 50%, another imaging modality will be necessary to demonstrate the extent of stenosis” <p>Definitions</p> <ul style="list-style-type: none"> ● Updated definition of “Ankle-Brachial Index (ABI)” <p>Supporting Information</p> <ul style="list-style-type: none"> ● Updated <i>Description of Services</i>, <i>Clinical Evidence</i>, and <i>References</i> sections to reflect the most current information ● Archived previous policy version CS166.F

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.