

# Embolization of the Ovarian and Iliac Veins for Pelvic Congestion Syndrome (for North Carolina Only)

Policy Number: CSNCT0574.01  
Effective Date: July 1, 2021

[Instructions for Use](#)

Table of Contents	Page
<a href="#">Application</a> .....	1
<a href="#">Coverage Rationale</a> .....	1
<a href="#">Definitions</a> .....	1
<a href="#">Applicable Codes</a> .....	2
<a href="#">Description of Services</a> .....	2
<a href="#">Clinical Evidence</a> .....	2
<a href="#">U.S. Food and Drug Administration</a> .....	4
<a href="#">References</a> .....	5
<a href="#">Policy History/Revision Information</a> .....	6
<a href="#">Instructions for Use</a> .....	6

Related Policies
None

## Application

This Medical Policy only applies to the state of North Carolina.

## Coverage Rationale

Embolization of the Ovarian Vein or Internal Iliac Vein is unproven and not medically necessary for treating Pelvic Congestion Syndrome due to insufficient evidence of efficacy.

## Definitions

**Embolization:** A procedure that uses particles, such as tiny gelatin sponges or beads, to block a blood vessel. Embolization may be used to stop bleeding or to block the flow of blood to a tumor or abnormal area of tissue. Types of embolization are arterial embolization, chemoembolization, and radioembolization. (National Cancer Institute)

**Fluoroscopy:** A radiological imaging technique that converts real-time X-rays from an X-ray machine into video images, usual for guiding diagnostic and interventional procedures. (FDA)

**Internal Iliac Vein (Hypogastric Vein):** Veins that originate deep in the pelvic region and extend to the lower portion of the abdomen, where they are joined with the right and left iliac veins, that together form the common iliac veins. (Merriam-Webster)

**Ovarian Vein:** One of a pair of veins that emerge from the broad ligament near the ovaries and the uterine tubes. (Mosby's Pocket Dictionary)

**Pelvic Congestion Syndrome (PCS):** A syndrome involving chronic pelvic pain usually associated with the Varices or Varicosities in the pelvic area. (Merck Manual)

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

**Coding Clarification:** According to the American Medical Association (AMA), CPT code 37241 is specific to venous embolization or occlusion and excludes lower extremity venous incompetency. Coding instructions state that 37241 should not be used to report treatment of incompetent extremity veins. For sclerosis of veins or endovenous ablation of incompetent extremity veins, see 36468-36479 (CPT Assistant, 2014).

CPT Code	Description
37241	Vascular embolization or occlusion, inclusive of all radiological supervision and interpretation, intraprocedural roadmapping, and imaging guidance necessary to complete the intervention; venous, other than hemorrhage (e.g., congenital or acquired venous malformations, venous and capillary hemangiomas, varices, varicoceles)

*CPT® is a registered trademark of the American Medical Association*

Diagnosis Code	Description
I86.2	Pelvic varices
N94.89	Other specified conditions associated with female genital organs and menstrual cycle
R10.2	Pelvic and perineal pain

## Description of Services

Pelvic Congestion Syndrome (PCS), also known as pelvic venous incompetence (PVI), causes noncyclic pelvic pain and discomfort, lasting for at least six months, and typically affects women of reproductive age. Varicosities of the Ovarian Veins and/or Internal Iliac Veins are believed to lead to PCS. For those patients who fail to adequately respond to conventional treatments (i.e., pharmacological therapy or surgical intervention), embolization therapy of the Ovarian Vein and/or Internal Iliac Vein has been proposed as an alternative (Nasser et al., 2014).

Patients with PCS may be treated with Ovarian Vein embolization following venography to visualize the affected veins (Bittles et al., 2008; Nasser et al., 2014). Under Fluoroscopy, an interventional radiologist guides a catheter to the affected vein, and inserts inert embolic agents to completely seal the vein. As a result, blood flow is rerouted, thereby reducing pressure within the targeted veins. Several types of embolic agents may be used, and include, but are not limited to, metal coils, sclerosing agents, and gelatin sponges. These agents may either be temporary or permanent. Since the Ovarian Veins and Internal Iliac Veins are in close proximity, embolization of the Internal Iliac Veins may also be necessary (Nasser et al., 2014).

## Clinical Evidence

The body of evidence in the peer-reviewed medical literature regarding embolization of the Ovarian Vein or Internal Iliac Veins for the treatment of PCS is insufficient and poor quality. While some studies suggest favorable results of embolization for the treatment of PCS, additional well-designed randomized controlled trials are necessary to establish the relative safety and efficacy of the embolization procedure.

A Hayes Health Technology Assessment (March 2020) states there is a lack of robust evidence regarding the efficacy on the use of ovarian or internal iliac vein embolization for the treatment of pelvic congestion syndrome (PCS). The overall low quality

of evidence primarily reflects individual study limitations, lack of comparison treatments, limited follow up, loss to follow-up and concerns regarding precision.

Guirola et al. (2018) provided one-year outcomes from a randomized, prospective, single-center study which compared fibered platinum coils (FPC) versus vascular plugs (VP) in 100 women with pelvic congestion syndrome (PCS). Patients were randomized to either FPC (n=50) or VP (n=50). Primary outcome (clinical success at one year using a VAS), number of devices, procedure and fluoroscopy times, radiation doses, costs, and complications were compared, and participants were followed at one, three, six, and 12 months. Clinical success and subjective improvement were not significantly different at one-year follow-up (89.7% for FPCs vs 90.6% for VPs; P = .760). The authors concluded that embolization for PCS resulted in pain relief in 90% of patients; clinical success was not affected by embolic device. Longer-term outcomes are needed to evaluate embolization procedures for the treatment of PCS.

Champaneria et al. (2016) conducted a systematic review of treatment to determine the effectiveness of treatment for PCS, and concluded that the data supporting its diagnosis and treatment are limited and of variable methodological quality. Their assessment revealed that embolization appears to provide symptomatic relief in the majority of women and is safe; however, the majority of included studies of embolization were relatively small case series and only a single randomized controlled trial was considered at risk of potential biases. There is scope and demand for considerable further research in which adequately powered randomized trials are essential to provide evidence on the effectiveness of embolization.

Daniels et al. (2016) conducted a systematic review to evaluate the effectiveness of embolization of incompetent pelvic veins performed to reduce CPP. Twenty-one prospective case series and one poor-quality randomized trial of embolization (involving a total of 1,308 women) were identified. The authors found that early substantial relief from pain was observed in approximately 75% of women undergoing embolization, and generally increased over time and was sustained. In addition, significant pain reductions following treatment were observed in all studies that measured pain on a visual analog scale. Repeat intervention rates were generally low. There were few data on the impact on menstruation, ovarian reserve, or fertility, but no concerns were noted. Transient pain was common following foam embolization, and there was a < 2% risk of coil migration. In the authors' opinion, embolization appears to provide symptomatic relief of CPP in the majority of women and is safe, although the quality of the evidence is low.

O'Brien and Gillespie (2015) conducted a systematic review of the diagnosis and treatment of PCS. Thirty-seven references were small series including fewer than 50 patients or individual case reports documenting medical, surgical, or endovascular treatment of PCS. The majority of these papers demonstrated successful treatment of symptoms from PCS with embolization of one or both ovarian veins in addition to treatment of refluxing internal iliac vein branches. In addition, open surgery and, more recently, endovascular stenting of LRV obstruction have shown some promise in alleviating symptoms attributed to nutcracker syndrome. Whereas a fairly large body of data regarding transcatheter ovarian vein embolization exists, the authors summarized that these studies are limited to relatively small clinical series and retrospective reviews. The success rates for the reduction of chronic pelvic pain (CPP) in these studies range from 47% to 94% with average follow-ups of 12 to 36 months. The authors concluded that ultimately, there remains an uncertainty as to the optimal technique for ovarian vein embolization, although a combination of coils and sclerosants has demonstrated clinical efficacy in a number of studies described before and is the most common published technique for ovarian vein embolization. In addition, there is currently no evidence to suggest a difference in symptomatic relief with regard to unilateral vs bilateral ovarian vein embolization.

In an evaluation of pelvic vein embolization indications, techniques and outcomes, Lopez (2015) summarized that evidence remains poor for its efficacy, and although initially anecdotal by way of case reports and small series, data is accumulating in larger series. There remains, however, a lack of robust evidence of its effectiveness, and this partly reflects the challenges of actually making the diagnosis clinically and radiologically, as well as the difficulty in assessing outcome. For pelvic congestion syndrome, symptomatic response is usually subjective but visual analogue scales (or variations thereof) have most often been used to attempt to identify a more objective outcome.

Hansrani et al. (2015) conducted a well-designed systematic review of the literature to evaluate the safety and effectiveness of transvenous occlusion of incompetent pelvic varicosities. Study authors selected 13 studies (n=866) that evaluated patients had CPP, PCS, or pelvic pain. The interventions generally consisted of transvenous occlusion of the ovarian and internal iliac veins (via the femoral or jugular veins) using metallic coils, sclerosants, or glue. A total of 10 studies were prospective uncontrolled, two were retrospective, and one was a randomized controlled trial (RCT) that included untreated controls. In nine of 13 studies, patients experienced significant improvement in pelvic pain and other PCS symptoms following embolization of the pelvic

varicosities when compared with baseline symptoms. One study reported 13% of recurrence at five years of follow-up. Embolization was generally considered technically successful, with 98 to 100% of veins occluded at first attempt. Adverse events included coil migration in 1.6% of patients, abdominal pain in 1.2%, and vein perforation in 0.6%. One serious complication was reported as coil migration to the lungs.

Nasser et al. (2014) conducted a retrospective review (n=113) in women with PCS who underwent embolization of the ovarian and pelvic varicose veins. The primary outcome was pain assessment using VAS. Patients were followed for a period of one year. Of the 113 included patients, 13 (10%) were lost to follow-up. At the end of follow-up, 37% had complete resolution of symptoms, 53% of patients had no pelvic pain and 47% had partial pain relief. There was also a significant reduction in the mean score of total associated symptoms at 12 months (2.69 at baseline to 0.92 at post-procedure). Complications were considered relatively minimal, with four cases of coil migrations. No other serious complications were reported.

In a single-center case series, Laborda et al. (2013) reported long-term results in 202 women with chronic pelvic pain (CPP). Inclusion criteria were: lower limb varices and CPP for more than six months, pelvic veins >6 mm on ultrasonography, and either venous reflux or presence of communicating veins. The primary outcomes were pain assessment using a visual analog scale (VAS), and patient satisfaction. Technical and clinical successes were also evaluated as secondary outcomes. Follow-up evaluations were conducted at one, three, and six months, and each year thereafter for five years. At five-years of follow-up, 11% of women were lost to follow-up, while 89% were available for evaluation. Study results demonstrated a significant improvement in pain symptoms (7.34±0.7 at baseline versus 0.78±1.2 at follow-up; P<0.0001). Technical success was considered 100%. Clinical success was observed in nearly 94% of all patients and approximately 33% experienced complete resolution of symptoms. Mean individual satisfaction scores were 7.39 [standard deviation (SD), 1.5; scale 0-9]. Major complications included four cases of coil migration and six cases of groin hematoma. Methodological limitations of this study include the case series design, the lack of appropriate controls for comparison, and the lack of diagnostic criteria during the patient selection process.

In a smaller case series, Hocquet et al. (2013) assessed the safety and efficacy of embolization for PCS (n=33). Average duration of follow-up was 26 months (range, 3-59). Patients experienced a significant reduction in pain following the procedure. The average VAS at baseline was 7.37 (SD, 0.99) compared with 1.36 (SD, 1.73) after embolization (P<0.0001). A total of 20 patients (~61%) had complete symptom resolution, 11 patients (~33%) had partial resolution, and two patients (6%) had no improvement.

Additional prospective and retrospective case series evaluated a small number of patients with PCS (Dorobisz et al., 2017; Marcelin et al., 2017; Siqueira et al., 2016; Nasser et al., 2014; Castenmiller et al., 2013; Laborda et al., 2013; Meneses et al., 2013; Smith et al., 2012; Tinelli et al., 2012; Mallios et al., 2011). Well-designed randomized controlled trials with a larger patient population and longer term outcomes are needed to further evaluate embolization procedures for pelvic congestion syndrome.

## Clinical Practice Guidelines

### *Society for Vascular Surgery (SVS)/ American Venous Forum (AVF)*

In a guideline published by the SVS and the AVF in 2011, guideline authors suggest “treatment of pelvic congestion syndrome and pelvic varices with coil embolization, plugs, or transcatheter sclerotherapy, used alone or together (2B).” The 2B recommendation indicates a “weak” recommendation based on moderate quality evidence, where the benefits of the technology are considered closely balanced with risks and burdens (Gloviczki et al., 2011).

Other professional society clinical guidelines that address embolization procedures for PCS were not identified.

## U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Numerous products used for vascular embolization, including sclerosing agents, and other substances, have been approved by the FDA. These products are generally classified under the product code: KRD (device, vascular, for promoting embolization), indexed in the Center for Devices and Radiological Health (CDRH) 510(k) database or Premarket Search Strategy. Available at: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm>. (Accessed February 18, 2020)

## References

- American Medical Association. Current Procedural Terminology: CPT Professional Edition. AMA Press.
- Bittles M, Hoffer E. Gonadal vein embolization: treatment of varicocele and pelvic congestion syndrome. *Semin Intervent Radiol*. 2008 Sep;25(3):261-70.
- Castenmiller PH, de Leur K, de Jong TE, et al. Clinical results after coil embolization of the ovarian vein in patients with primary and recurrent lower-limb varices with respect to vulval varices. *Phlebology*. 2013 Aug;28(5):234-8.
- Champaneria R, Shah L, Moss J, et al. The relationship between pelvic vein incompetence and chronic pelvic pain in women: systematic reviews of diagnosis and treatment effectiveness. *Health Technol Assess*. 2016 Jan;20(5):1-108.
- Daniels JP, Champaneria R, Shah L, et al. Effectiveness of embolization or sclerotherapy of pelvic veins for reducing chronic pelvic pain: a systematic review. *J Vasc Interv Radiol*. 2016 Oct;27(10):1478-1486.
- Dorobisz TA, Garcarek JS, Kurcz J, et al. Diagnosis and treatment of pelvic congestion syndrome: Single-centre experiences. *Adv Clin Exp Med*. 2017 Mar-Apr;26(2):269-276.
- Gloviczki P, Comerota AJ, Dalsing MC, et al. The care of patients with varicose veins and associated chronic venous diseases: clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum. *J Vasc Surg*. 2011; 53:2S-48S.
- Guirola JA, Sánchez-Ballestin M, Sierre S, et al. A randomized trial of endovascular embolization treatment in pelvic congestion syndrome: fibered platinum coils versus vascular plugs with 1-year clinical outcomes. *J Vasc Interv Radiol*. 2018 Jan;29(1):45-53.
- Hansrani V, Abbas A, Bhandari S, et al. Trans-venous occlusion of incompetent pelvic veins for chronic pelvic pain in women: a systematic review. *Eur J Obstet Gynecol Reprod Biol*. 2015 Feb;185:156-63.
- Hayes, Inc. Health Technology Assessment. Ovarian or Internal Iliac Vein Embolization for Treatment of Pelvic Congestion Syndrome. Lansdale PA: Hayes, Inc.; March 2020.
- Hocquelet A, Bras YL, Balian E, et al. Evaluation of the efficacy of endovascular treatment of pelvic congestion syndrome. *Diagnostic and Interventional Imaging*. 2014; 95:301-306.
- Laborda A, Medrano J, de Blas I, et al. Endovascular treatment of pelvic congestion syndrome: visual analog scale (VAS) long-term follow-up clinical evaluation in 202 patients. *Cardiovasc Intervent Radiol*. 2013 Aug;36(4):1006-14.
- Lopez, AJ. Female pelvic vein embolization: indications, techniques, and outcomes. *Cardiovasc Intervent Radiol* (2015) 38:806–820.
- Mallios A, Laurian C, Houbballah R, et al. Curative treatment of pelvic arteriovenous malformation—an alternative strategy: transvenous intra-operative embolisation. *Eur J Vasc Endovasc Surg*. 2011 Apr;41(4):548-53.
- Marcelin C, Izaaryene J, Castelli M, et al. Embolization of ovarian vein for pelvic congestion syndrome with ethylene vinyl alcohol copolymer (Onyx®). *Diagn Interv Imaging*. 2017 Jun 21. pii: S2211-5684(17)30152-3.
- Meneses L, Fava M, Diaz P. Embolization of incompetent pelvic veins for the treatment of recurrent varicose veins in lower limbs and pelvic congestion syndrome. *Cardiovasc Intervent Radiol*. 2013; 36(1):128-132.
- Merck Manual. Available at: <https://www.merckmanuals.com/professional/gynecology-and-obstetrics>. Accessed March 2, 2021.
- Merriam-Webster Dictionary. Available at: <https://www.merriam-webster.com/dictionary>. Accessed March 2, 2021.
- Mosby's Pocket Dictionary of Medicine, Nursing & Health Professions - E-Book; page 970. Available at: <https://books.google.com/>. Accessed March 2, 2021.
- Nasser F, Cavalcante RN, Affonso BB, et al. Safety, efficacy, and prognostic factors in endovascular treatment of pelvic congestion syndrome. *Int J Gynaecol Obstet*. 2014; 125(1):65-68.
- National Cancer Institute Dictionary of Cancer Terms. Available at: <https://www.cancer.gov/publications/dictionaries>. Accessed March 2, 2021.
- O'Brien MT, Gillespie DL. Diagnosis and treatment of the pelvic congestion syndrome. *J Vasc Surg Venous Lymphat Disord*. 2015 Jan;3(1):96-106.

Siqueira FM, Monsignore LM, Rosa-E-Silva JC, et al. Evaluation of embolization for periuterine varices involving chronic pelvic pain secondary to pelvic congestion syndrome. Clinics (Sao Paulo). 2016 Dec 1;71(12):703-708.

Smith PC. The outcome of treatment for pelvic congestion syndrome. Phlebology. 2012; 27 Suppl 1:74-77.

Tinelli A, Prudenzeno R, Torsello M, et al. Suprapubic percutaneous sclero-embolization of symptomatic female pelvic varicocele under local anesthesia. Eur Rev Med Pharmacol Sci. 2012 Jan;16(1):111-7.

U.S. Food and Drug Administration. Available at: <https://www.fda.gov/radiation-emitting-products/medical-x-ray-imaging/fluoroscopy>. Accessed March 2, 2021.

## Policy History/Revision Information

Date	Summary of Changes
07/01/2021	<ul style="list-style-type: none"><li>New Medical Policy</li></ul>

## Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual<sup>®</sup> criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.