

Outpatient Surgical Procedures – Site of Service

Guideline Number: CS143.N
Effective Date: June 1, 2022

[➔ Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	2
References	3
Guideline History/Revision Information	4
Instructions for Use	4

Related Community Plan Policy
• Obstructive and Central Sleep Apnea Treatment
Commercial Policy
• Outpatient Surgical Procedures – Site of Service
Medicare Advantage Coverage Summary
• Hospital Services (Inpatient and Outpatient)

Application

This Utilization Review Guideline only applies to the states of Arizona, Florida, Maryland, Michigan, Mississippi, Missouri, New York, Ohio, Rhode Island, Texas, and Washington.

Refer to the guidelines listed below for the following states:

State	Policy/Guideline
New Jersey	Outpatient Surgical Procedures – Site of Service (for New Jersey Only)
North Carolina	Outpatient Surgical Procedures – Site of Service (for North Carolina Only)
Pennsylvania	Outpatient Surgical Procedures – Site of Service (for Pennsylvania Only)
Tennessee	Outpatient Surgical Procedures – Site of Service (for Tennessee Only)

Coverage Rationale

UnitedHealthcare members may choose to receive surgical procedures in an ambulatory surgical center (ASC) or other locations. We are conducting site of service medical necessity reviews, however, to determine whether the outpatient hospital department is medically necessary, in accordance with the terms of the member’s benefit plan. If the outpatient hospital department is not considered medically necessary, this location will not be covered under the member’s plan.

Certain planned surgical procedures performed in a hospital outpatient department are considered medically necessary for an individual who meets any of the following criteria:

- Advanced liver disease (MELD Score > 8)
- Advance surgical planning determines an individual requires overnight recovery and care following a surgical procedure
- Anticipated need for transfusion
- Bleeding disorder requiring replacement factor or blood products or special infusion products to correct a coagulation defect
- Cardiac arrhythmia (symptomatic arrhythmia despite medication)
- Chronic obstructive pulmonary disease (COPD) (FEV1 < 50%)
- Coronary artery disease ([CAD]/peripheral vascular disease [PVD]) (ongoing cardiac ischemia requiring medical management or recently placed [within 1 year] drug eluting stent)
- Developmental stage or cognitive status warranting use of a hospital outpatient department

- End stage renal disease ([hyperkalemia above reference range] receiving peritoneal or hemodialysis)
- History of cerebrovascular accident (CVA) or transient ischemic attack (TIA) (recent event [< 3 months])
- History of myocardial infarction (MI) (recent event [< 3 months])
- Individuals with drug eluting stents (DES) placed within one year or bare metal stents (BMS) or plain angioplasty within 90 days unless acetylsalicylic acid and antiplatelet drugs will be continued by agreement of surgeon, cardiologist and anesthesia
- Less than 19 years of age unless otherwise required by federal, state or contractual requirements
- Ongoing evidence of myocardial ischemia
- Poorly Controlled asthma (FEV1 $< 80\%$ despite medical management)
- Pregnancy
- Prolonged surgery (> 3 hours)
- Resistant hypertension (Poorly Controlled)
- Severe valvular heart disease
- Sleep apnea (moderate to severe Obstructive Sleep Apnea [OSA])
- Uncompensated chronic heart failure (CHF) (NYHA class III or IV)
- Uncontrolled diabetes with recurrent diabetic ketoacidosis (DKA) or severe hypoglycemia

A planned surgical procedure performed in a hospital outpatient department is considered medically necessary if there is an inability to access an ambulatory surgical center for the procedure due to any one of the following:

- There is no geographically accessible ambulatory surgical center that has the necessary equipment for the procedure; or
- There is no geographically accessible ambulatory surgical center available at which the individual's physician has privileges; or
- An ASC's specific guideline regarding the individual's weight or health conditions that prevents the use of an ASC

Documentation Requirements

Documentation requirements include the following, when applicable:

- History
- Physical examination including patient weight and co-morbidities
- Surgical plan
- Physician privileging information related to the need for the use of the hospital outpatient department
- American Society of Anesthesiologists (ASA) score, as applicable
- Specific criteria (see coverage rationale) that qualifies the individual for the site of service requested

Planned Surgical Procedures List

Site of service medical necessity reviews will be conducted for certain surgical procedures only when performed in an outpatient hospital setting. For the complete list of surgical procedure codes requiring prior authorization for each state, refer to the [UnitedHealthcare Community Plan Prior Authorization List](#). (Accessed May 12, 2021)

Definitions

Check the member specific benefit plan document or any applicable federal or state contractual or regulatory requirements. In the event of a conflict, the federal, state or contractual definitions for benefit plan coverage supersede this Utilization Review Guideline.

ASA Physical Status Classification System Risk Scoring Tool: The American Society of Anesthesiologists (ASA) physical status classification system was developed to offer clinicians a simple categorization of a patient's physiological status that can be helpful in predicting operative risk. The ASA score is a subjective assessment of a patient's overall health that is based on five classes (ASA, 2020).

Obstructive Sleep Apnea (OSA): The American Academy of Sleep Medicine (AASM) defines OSA as a sleep related breathing disorder that involves a decrease or complete halt in airflow despite an ongoing effort to breathe. OSA severity is defined as:

- Mild for AHI or RDI ≥ 5 and < 15
- Moderate for AHI or RDI ≥ 15 and ≤ 30

- Severe for AHI or RDI > 30/hr (AASM, 2021)

Poorly Controlled: Requiring three or more drugs to control blood pressure. (Sheppard, 2017).

References

- Adamson P, Peters W, Janney C, Panchbhavi V. The safety of foot and ankle procedures at an ambulatory surgery center. *J Orthop*. 2020 Mar 28;21:203-206.
- American Academy of Sleep Medicine (AASM). Obstructive Sleep Apnea. Available at: <https://aasm.org/resources/factsheets/sleepapnea.pdf>. Accessed September 16, 2021.
- American Heart Association. Classes of Heart Failure. Available at: http://www.heart.org/HEARTORG/Conditions/HeartFailure/AboutHeartFailure/Classes-of-Heart-Failure_UCM_306328_Article.jsp. Accessed April 10, 2021.
- American Society of Anesthesiologists (ASA) [Physical Status Classification System](#). December 13, 2020. Accessed September 16, 2021.
- American Society of Anesthesiologists. Guidelines for ambulatory anesthesia and surgery. October 17, 2018.
- American Society of Anesthesiologists. Guidelines for patient care in anesthesiology. October 26, 2016.
- Bhattacharyya N. Ambulatory pediatric otolaryngologic procedures in the United States: characteristics and perioperative safety. *Laryngoscope*. 2010 Apr;120(4):821-5.
- Bilimoria K, Liu Y, Paruch J, et al. Development and evaluation of the Universal ACS NSQIP Surgical Risk Calculator: a decision aide and informed consent tool for patients and surgeons. *J Am Coll Surg*. 2013 November; 217(5): 833–842.e3.
- Brolin TJ, Mulligan RP, Azar FM, et al. Neer Award 2016: Outpatient total shoulder arthroplasty in an ambulatory surgery center is a safe alternative to inpatient total shoulder arthroplasty in a hospital: a matched cohort study. *J Shoulder Elbow Surg*. 2017;26(2):204-8.
- Brophy RH, Bansal A, Rogalski BL, et al. Risk factors for surgical site infections after orthopaedic surgery in the ambulatory surgical center setting. *J Am Acad Orthop Surg*. 2019 Oct 15;27(20):e928-e934.
- Cancienne JM, Brockmeier SF, Gulotta LV, et al. Ambulatory total shoulder arthroplasty: A comprehensive analysis of current trends, complications, readmissions, and costs. *J Bone Joint Surg Am*. 2017;99(8):629- 37.
- Epstein LJ, Kristo D, Strollo PJ Jr, et al. Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med*. 2009 Jun 15; 5(3):263-76.
- Fleisher LA, Fleischmann KE, Auerbach AD, et al. American College of Cardiology; American Heart Association. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. 2014 Dec 9;64(22):e77-137.
- Ford MC, Walters JD, Mulligan RP, et al. Safety and cost-effectiveness of outpatient unicompartmental knee arthroplasty in the ambulatory surgery center: A Matched Cohort Study. *Orthop Clin North Am*. 2020 Jan;51(1):1-5.
- Friedlander DF, Krimphove MJ, Cole AP, et al. Where Is the value in ambulatory versus inpatient surgery? *Ann Surg*. 2021 May 1;273(5):909-916.
- Friedman L S. Surgery in the patient with liver disease. *Trans Am Clin Climatol Assoc*. 2010; 121: 192–205.
- Goyal N, Chen AF, Padgett SE, et al. Otto Aufranc Award: A multicenter, randomized study of outpatient versus inpatient total hip arthroplasty. *Clin Orthop*. 2017;475(2):364-7.
- Joshi G, Ahmad S; Riad W. et al. Selection of obese patients undergoing ambulatory surgery: a systematic review of the literature. *Anesthesia & Analgesia*. November 2013; 117(5): 1082–1091.
- Joshi G, Ankichetty P, Gan T, and Chung F. Society for Ambulatory Anesthesia Consensus Statement on preoperative selection of adult patients with obstructive sleep apnea scheduled for ambulatory surgery. *Anesthesia & Analgesia*: November 2012; 115(5): 1060–1068.

Joshi G; Chung F; Vann Mary Ann, et al. Society for Ambulatory Anesthesia Consensus Statement on perioperative blood glucose management in diabetic patients undergoing ambulatory surgery. *Anesthesia & Analgesia*. December 2010; 111(6): 1378–1387.

Lovald S, Ong K, Lau E, et al. Patient selection in outpatient and short-stay total knee arthroplasty. *J Surg Orthop Adv*. 2014 Spring;23(1):2-8.

MacKoul P, Danilyants N, Baxi R, et al. Laparoscopic hysterectomy outcomes: hospital vs ambulatory surgery center. *JSLs*. 2019 Jan-Mar;23(1):e2018.00076.

Maganti K, Rigolin VH, Sarano ME, Bonow RO. Valvular heart disease: diagnosis and management. *Mayo Clin Proc*. 2010 May;85(5):483-500.

Mathis MR, Naughton NN, Shanks AM, et al. Patient selection for day case-eligible surgery: identifying those at high risk for major complications. *Anesthesiology*. 2013 Dec;119(6):1310-21.

Medicare Claims Processing Manual. Chapter 14 - Ambulatory Surgical Centers.

National Kidney Foundation. Clinical update on hyperkalemia. A chronic risk for CKD patients and a potential barrier to recommended CKD treatment. https://www.kidney.org/sites/default/files/02-10-6785_HBE_Hyperkalemia_Bulletin.pdf. Accessed April 10, 2021.

Orthopedic Certification. Pathways to excellence in patient care. Joint Commission.

Sankar A, Johnson SR, Beattie WS, et al. Reliability of the American Society of Anesthesiologists physical status scale in clinical practice. *Br J Anaesth*. 2014 Sep;113(3):424-32.

Sheffer BW, Kelly DM, Spence DD, et al. Can pediatric orthopaedic surgery be done safely in a freestanding ambulatory surgery center? Review of 3780 cases. *J Pediatr Orthop*. 2021 Jan;41(1):e85-e89.

Sheppard JP, Martin U, McManus RJ. Diagnosis and management of resistant hypertension. *Heart*. 2017 Aug;103(16):1295-1302.

Toy PC, Fournier MN, Throckmorton TW, et al. Low rates of adverse events following ambulatory outpatient total hip arthroplasty at a free-standing surgery center. *J Arthroplasty*. 2018 Jan;33(1):46-50.

Whippey A, Kostandoff G, Ma HK, et al. Predictors of unanticipated admission following ambulatory surgery in the pediatric population: a retrospective case-control study. *Paediatr Anaesth*. 2016 Aug;26(8):831-7.

UnitedHealthcare Community Plan Prior Authorization List. Available at: <https://www.uhcprovider.com/en/health-plans-by-state.html>. Accessed May 5, 2021.

Guideline History/Revision Information

Date	Summary of Changes
06/01/2022	<p>Application <i>Florida and Mississippi</i></p> <ul style="list-style-type: none"> Added language to indicate this Utilization Review Guideline applies to the states of Florida and Mississippi (retired state-specific policy versions) <p>Supporting Information</p> <ul style="list-style-type: none"> Archived previous policy version CS143.M

Instructions for Use

This Utilization Review Guideline provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this guideline, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Utilization Review Guideline is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. The UnitedHealthcare Utilization Review Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.