

# Papzimeos™ (Zopapogene Imadenovec-Drba)

**Policy Number:** CS2026D0141B  
**Effective Date:** April 1, 2026

[➔ Instructions for Use](#)

<b>Table of Contents</b>	<b>Page</b>
<a href="#">Application</a> .....	1
<a href="#">Coverage Rationale</a> .....	1
<a href="#">Applicable Codes</a> .....	1
<a href="#">Background</a> .....	2
<a href="#">Clinical Evidence</a> .....	2
<a href="#">U.S. Food and Drug Administration</a> .....	2
<a href="#">References</a> .....	2
<a href="#">Policy History/Revision Information</a> .....	3
<a href="#">Instructions for Use</a> .....	3

<b>Commercial Policy</b>
<ul style="list-style-type: none"> <li><a href="#">Papzimeos™ (Zopapogene Imadenovec-Drba)</a></li> </ul>

## Application

This Medical Benefit Drug Policy does not apply to the states listed below; refer to the state-specific policy/guideline, if noted:

State	Policy/Guideline
Kansas	None
North Carolina	None
Ohio	<a href="#">Papzimeos™ (Zopapogene Imadenovec-Drba) (for Ohio Only)</a>

## Coverage Rationale

**Papzimeos is proven and medically necessary for the treatment of recurrent respiratory papillomatosis in patients who meet all of the following criteria:**

- Diagnosis of recurrent respiratory papillomatosis (RRP); **and**
- Patient is 18 years of age or older; **and**
- Patient has required surgery (i.e., surgical resection of papillomas, laser ablation of papillomas) to remove laryngotracheal papillomas prior to treatment with Papzimeos; **and**
- Surgical debulking of present visible papilloma will be performed prior to the initial, third, and fourth dose of Papzimeos; **and**
- Prescribed by or in consultation with a specialist knowledgeable in the treatment of recurrent respiratory papillomatosis (e.g., otolaryngologist, pulmonologist, oncologist); **and**
- Patient has not received a previous complete treatment course (i.e., four doses over a 12-week interval) with Papzimeos; **and**
- Dosing is in accordance with the United States Food and Drug Administration approved labeling; **and**
- Authorization will be issued for no more than one treatment course (i.e., four doses) per lifetime

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
J3404	Injection, zopapogene imadenovec-drba suspension, per therapeutic dose

  

Diagnosis Code	Description
D14.1	Benign neoplasm of larynx
J38.7	Other diseases of larynx

## Background

Papzimeos is a non-replicating adenoviral vector-based immunotherapy designed to express a fusion antigen of selected regions of human papillomavirus (HPV) proteins expressed in HPV 6- and HPV 11-infected cells. Papzimeos is designed to generate an immune response directed against HPV 6 and HPV 11 proteins in patients with recurrent respiratory papillomatosis.

## Clinical Evidence

Papzimeos was evaluated in an open-label, single-arm study in 38 adults with recurrent respiratory papillomatosis (RRP). Patients received Papzimeos on days 1, 15, 43 and 85. The primary efficacy endpoint was the percentage of patients with a complete response to Papzimeos treatment, defined as no requirement of surgical intervention in the 12 months after treatment. In the 35 patients that received a dose of  $5 \times 10^{11}$  particle units (PU) per injection, the median age at RRP diagnosis was 35 years (range 1-68) and median age at the start of treatment was 49 years (range 20-88). 23 (66%) patients were diagnosed with adult-onset RRP and 12 (34%) with juvenile-onset disease. Most patients were male [20 (57%) of 35] and required up to hundreds of lifetime clinically indicated interventions to control RRP [median 40 (range 3 to > 400)]. Four (11%) of 35 patients had pulmonary RRP evaluable by RECIST criteria. The median number of previous medical treatments for RRP was 2 (range 1-8). The median number of clinically indicated interventions in the 12 months before the study treatment was 4 (3-10). The median pretreatment anatomic Derkay score was 8 (3-31). The median pretreatment VHI-10 score was 24 (6-40), indicating severe dysphonia. At the dose of  $5 \times 10^{11}$  PU per injection, 18 of 35 patients received a complete response at 12 months [complete response rate 51% (95% CI: 34, 69)]. At the dose of  $1 \times 10^{11}$  PU per injection, no patient (0 of 3) received a complete response. Warnings and precautions for Papzimeos include injection site reactions and thrombotic events. The most common adverse reactions ( $\geq 5\%$ ) with Papzimeos use were injection site reactions, fatigue, chills, pyrexia, myalgia, and nausea.

## U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Papzimeos is a non-replicating adenoviral vector-based immunotherapy indicated for the treatment of adults with recurrent respiratory papillomatosis.

## References

1. Papzimeos [prescribing information]. Germantown, MD: Precigen, Inc; August 2025.
2. Norberg SM, Valdez J, Napier S, et al. PRGN-2012 gene therapy in adults with recurrent respiratory papillomatosis: a pivotal phase 1/2 clinical trial. *Lancet Respir Med*. 2025;13(4):318-326. doi:10.1016/S2213-2600(24)00368-0.
3. Stachler RJ, Francis DO, Schwartz SR, et al. Clinical Practice Guideline: Hoarseness (Dysphonia) (Update). *Otolaryngol Head Neck Surg*. 2018;158(1\_suppl):S1-S42. doi:10.1177/0194599817751030.
4. Lawlor C, Balakrishnan K, Bottero S, et al. International Pediatric Otolaryngology Group (IPOG): Juvenile-onset recurrent respiratory papillomatosis consensus recommendations. *Int J Pediatr Otorhinolaryngol*. 2020;128:109697. doi:10.1016/j.ijporl.2019.109697.
5. Sidell DR, Balakrishnan K, Best SR, et al. Systemic Bevacizumab for Treatment of Respiratory Papillomatosis: International Consensus Statement. *Laryngoscope*. 2021;131(6):E1941-E1949. doi:10.1002/lary.29343.
6. American Academy of Otolaryngology – Head and Neck Surgery (AAO-HNS). Position statement: Recurrent respiratory papillomatosis and Gardasil vaccination. AAO-HNS. April 5, 2021. Accessed December 1, 2025. <https://www.entnet.org/resource/position-statement-recurrent-respiratory-papillomatosis-and-gardasil-vaccination/>.

## Policy History/Revision Information

Date	Summary of Changes
04/01/2026	<p data-bbox="337 207 500 239"><b>Application</b></p> <p data-bbox="337 239 444 270"><b>Indiana</b></p> <ul data-bbox="337 270 1135 302" style="list-style-type: none"><li data-bbox="337 270 1135 302">• Removed content/language pertaining to the state of <b>Louisiana</b></li></ul> <p data-bbox="337 302 477 333"><b>Louisiana</b></p> <ul data-bbox="337 333 1135 365" style="list-style-type: none"><li data-bbox="337 333 1135 365">• Removed content/language pertaining to the state of <b>Louisiana</b></li></ul> <p data-bbox="337 365 407 396"><b>Ohio</b></p> <ul data-bbox="337 396 915 428" style="list-style-type: none"><li data-bbox="337 396 915 428">• Added reference link to state-specific version</li></ul> <p data-bbox="337 428 586 459"><b>Applicable Codes</b></p> <ul data-bbox="337 459 1492 533" style="list-style-type: none"><li data-bbox="337 459 1492 533">• Updated list of applicable HCPCS codes to reflect quarterly edits; replaced C9399, J3490, and J3590 with J3404</li></ul> <p data-bbox="337 533 662 564"><b>Supporting Information</b></p> <ul data-bbox="337 564 964 596" style="list-style-type: none"><li data-bbox="337 564 964 596">• Archived previous policy version CS2026D0141A</li></ul>

## Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.