

Somatostatin Analogs

Policy Number: CS2020D0036P
Effective Date: December 1, 2020

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Applicable Codes	3
Background	5
Clinical Evidence	5
U.S. Food and Drug Administration	12
Centers for Medicare and Medicaid Services	13
References	13
Policy History/Revision Information	16
Instructions for Use	16

Related Community Plan Policy
• Oncology Medication Clinical Coverage
Commercial Policy
• Somatostatin Analogs

Application

This Medical Benefit Drug Policy does not apply to the states listed below; refer to the state-specific policy/guideline, if noted:

State	Policy/Guideline
Indiana	None
Kansas	None
Kentucky	Somatostatin Analogs (for Kentucky Only)
Louisiana	Somatostatin Analogs (for Louisiana Only)
North Carolina	None

Coverage Rationale

Refer to the Medical Benefit Drug Policy titled [Oncology Medication Clinical Coverage](#) for updated information based on the National Comprehensive Cancer Network (NCCN) Drugs & Biologics Compendium for oncology indications.

Sandostatin (octreotide acetate) and Sandostatin LAR (octreotide acetate LAR) are proven for the treatment of one of the following:

- Bleeding gastroesophageal varices associated with liver disease^{4,7,17,38,43}
Octreotide acetate is medically necessary for the treatment of bleeding esophageal varices when both of the following criteria are met:
 - Diagnosis of bleeding esophageal varices associated with liver disease; and
 - Octreotide acetate will be used as an adjunct to endoscopic therapy
- Diarrhea, chemotherapy and/or radiation-induced^{5,41}
- Diarrhea, refractory HIV/AIDS-related^{6,13,15,24,28,31,35,39}
Octreotide acetate is medically necessary for the treatment of refractory HIV/AIDS-related diarrhea when both of the following criteria are met:
 - Diagnosis of HIV/AIDS-related diarrhea; and

- History of failure, contraindication, or intolerance to standard therapy (e.g., loperamide, diphenoxylate/atropine)
- Malignant bowel disease^{27,29}

Sandostatin immediate release (IR) is proven and medically necessary for the treatment of acromegaly when both of the following criteria are met:

- Diagnosis of acromegaly by one of the following:³²
 - Serum GH level > 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis
 - Elevated serum IGF-1 levels (above the age and gender adjusted normal range as provided by the physician's lab) at time of diagnosis;
- and
- One of the following:^{22,26}
 - Inadequate response to one of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy;
 - or
 - Not a candidate for any of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy

Sandostatin LAR is proven and medically necessary for the treatment of acromegaly when all of the following criteria are met:

- Diagnosis of acromegaly by one of the following:³²
 - Serum GH level > 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis
 - Elevated serum IGF-1 levels (above the age and gender adjusted normal range as provided by the physician's lab) at time of diagnosis;
- and
- One of the following:^{22,26}
 - Inadequate response to one of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy;
 - or
 - Not a candidate for any of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy;
- and
- Initial treatment with octreotide immediate release (IR) has been shown to be effective and tolerated³⁷

Signifor and Signifor LAR (pasireotide diaspate) are proven and medically necessary for the treatment of Cushing's disease when both of the following criteria are met:

- Diagnosis of Cushing's disease;⁴⁹ and
- One of the following:
 - Inadequate response to pituitary surgery; or
 - Not a candidate for pituitary surgery

Signifor LAR (pasireotide) is proven and medically necessary for the treatment of acromegaly when both of the following criteria are met:

- Diagnosis of acromegaly by one of the following:⁵³
 - Serum GH level > 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis
 - Elevated serum IGF-1 levels (above the age and gender adjusted normal range as provided by the physician's lab) at time of diagnosis;
- and

- One of the following:^{22,26}
 - Inadequate response to one of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy;
 - or
 - Not a candidate for any of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy

Somatuline Depot (lanreotide) is proven and medically necessary for the treatment of acromegaly when both of the following criteria are met:

- Diagnosis of acromegaly by one of the following:³²
 - Serum GH level > 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis
 - Elevated serum IGF- 1 levels (above the age and gender adjusted normal range as provided by the physician's lab) at time of diagnosis;
- and
- One of the following:^{22,26}
 - Inadequate response to one of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy;
 - or
 - Not a candidate for any of the following:
 - Surgery
 - Radiotherapy
 - Dopamine agonist (e.g., bromocriptine, cabergoline) therapy

Somatostatin analogs are unproven and not medically necessary for treating the following conditions:

- Chylothorax
- Dumping syndrome
- Pancreatitis
- Persistent hyperinsulinemic hypoglycemia of infancy
- Prevention of postoperative complications following pancreatic surgery
- Short bowel syndrome

Somatostatin analogs are unproven for treating other conditions not listed above as proven due to the lack of published clinical evidence of safety and/or efficacy in published peer-reviewed medical literature.

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
J1930	Injection, lanreotide, 1 mg
J2353	Injection, octreotide, depot form for intramuscular injection, 1 mg
J2354	Injection, octreotide, non-depot form for subcutaneous or intravenous injection, 25 mcg
J2502	Injection, pasireotide long acting, 1 mg

Diagnosis Code	Description
B20	Human immunodeficiencyvirus [HIV] disease
C7A.010	Malignant carcinoid tumor of the duodenum
C7A.011	Malignant carcinoid tumor of the jejunum
C7A.012	Malignant carcinoid tumor of the ileum
C7A.019	Malignant carcinoid tumor of the small intestine, unspecified portion
C7A.020	Malignant carcinoid tumor of the appendix
C7A.021	Malignant carcinoid tumor of the cecum
C7A.022	Malignant carcinoid tumor of the ascending colon
C7A.024	Malignant carcinoid tumor of the descending colon
C7A.025	Malignant carcinoid tumor of the sigmoid colon
C7A.026	Malignant carcinoid tumor of the rectum
C7A.029	Malignant carcinoid tumor of the large intestine, unspecified portion
C7A.092	Malignant carcinoid tumor of the stomach
C7A.094	Malignant carcinoid tumor of the foregut, unspecified
C7A.095	Malignant carcinoid tumor of the midgut, unspecified
C7A.096	Malignant carcinoid tumor of the hindgut, unspecified
C25.4	Malignant neoplasm of endocrine pancreas
E22.0	Acromegaly and pituitary gigantism
E24.0	Pituitary-dependent Cushing's disease
E34.0	Carcinoid syndrome
E34.4	Constitutional tall stature
I85.01	Esophageal varices with bleeding
I85.11	Secondary esophageal varices with bleeding
K52.0	Gastroenteritis and colitis due to radiation
K52.89	Other specified noninfective gastroenteritis and colitis
K52.9	Noninfective gastroenteritis and colitis, unspecified
K56.1	Intussusception
K56.2	Volvulus
K56.50	Intestinal adhesions [bands], unspecified as to partial versus complete obstruction
K56.51	Intestinal adhesions [bands], with partial obstruction
K56.52	Intestinal adhesions [bands] with complete obstruction
K56.600	Partial intestinal obstruction, unspecified as to cause
K56.601	Complete intestinal obstruction, unspecified as to cause
K56.609	Unspecified intestinal obstruction, unspecified as to partial versus complete obstruction
K56.690	Other partial intestinal obstruction
K56.691	Other complete intestinal obstruction
K56.699	Other intestinal obstruction unspecified as to partial versus complete obstruction
K59.9	Functional intestinal disorder, unspecified
R19.7	Diarrhea, unspecified

Background

Sandostatin is a cyclic octapeptide prepared as a clear sterile solution of octreotide acetate salt, in a buffered lactic acid solution for administration by deep subcutaneous (SC) or intravenous (IV) injection. It is a long-acting octapeptide with pharmacologic actions mimicking those of the natural hormone somatostatin. The principal effects of octreotide include inhibition of growth hormone (GH), glucagon, and insulin. Other effects include diminution of luteinizing hormone response to gonadotropin-releasing hormone, reduction of splanchnic blood flow, and inhibition of release of several gastrointestinal hormones, including serotonin, gastrin, vasoactive intestinal peptide, secretin, motilin, and pancreatic polypeptide.³⁶

Sandostatin LAR is a long-acting dosage form that maintains all of the clinical and pharmacological characteristics of the immediate-release dosage form with the added feature of slow release of octreotide from the site of injection, reducing the need for frequent administration. It is indicated in patients in whom initial treatment with Sandostatin injection has been shown to be effective and tolerated. Sandostatin LAR is designed to be injected intramuscularly (intragluteally) once every 4 weeks and must be administered under the supervision of a physician.³⁷

Signifor is an injectable cyclohexapeptide somatostatin analogue. Pasireotide exerts its pharmacological activity via binding to somatostatin receptors (SSTRs). Five human somatostatin receptor subtypes are known: SSTR 1, 2, 3, 4, and 5. These receptor subtypes are expressed in different tissues under normal physiological conditions. Corticotroph tumor cells from Cushing's disease patients frequently over-express SSTR5 whereas the other receptor subtypes are often not expressed or are expressed at lower levels. Pasireotide binds and activates the SSTRs resulting in inhibition of ACTH secretion, which leads to decreased cortisol secretion.⁴⁹

Signifor LAR is a long-acting release form of pasireotide, a somatostatin analogue. Pasireotide exerts its pharmacological activity via binding to somatostatin receptors (SSTRs). Five human somatostatin receptor subtypes are known: SSTR 1, 2, 3, 4, and 5. These receptor subtypes are expressed in different tissues under normal physiological conditions. Corticotroph tumor cells from Cushing's disease patients frequently over-express SSTR5 whereas the other receptor subtypes are often not expressed or are expressed at lower levels. Pasireotide binds and activates the SSTRs resulting in inhibition of ACTH secretion, which leads to decreased cortisol secretion.^{49,53}

Somatuline Depot is a prolonged-release formulation for deep subcutaneous injection. It is a synthetic octapeptide with a biological activity similar to naturally occurring somatostatin. Like somatostatin, lanreotide is an inhibitor of various endocrine, neuroendocrine, exocrine, and paracrine functions.⁴⁷

Clinical Evidence

Octreotide

Proven

Bleeding Gastroesophageal Varices

Octreotide alone may not be useful for acute variceal bleeding due to the risk for tachyphylaxis, and results of meta-analyses of trials of octreotide are controversial. A Cochrane review of trials comparing somatostatin or its analogues with placebo in general showed that the drugs did not significantly reduce mortality. Overall, units of blood transfused were reduced, and the number of patients failing initial hemostasis was reduced in the studies.¹⁷

A randomized, controlled clinical trial (n=105) compared the efficacy of octreotide (50 g/hr for 48 hours) combined with sclerotherapy versus sclerotherapy alone in patients with acute bleeding from gastroesophageal varices. Initial control of bleeding was achieved in 46/51 (90.2%) patients who received combined treatment compared to 41/54 (75.9%) patients (p=0.05) in the sclerotherapy alone group. Rebleeding after the first 48 hours was less in the octreotide treated patients 2/46 vs. 8/41 patients (p=0.003). The octreotide treated patients had a better short term (5 days) survival without rebleeding 44/51 vs. 33/54 (p=0.003) and shorter hospital stay, 5.31 ± 3.87 days vs. 6.63 ± 3.86 (p=0.008) as compared to sclerotherapy alone group. The blood transfusion requirement was also less in the combined treatment group 3.88 ± 2.80 vs. 5.37 ± 3.15 units (p=0.002).³⁸

The efficacy of subcutaneous octreotide, administered after emergency sclerotherapy, was investigated to prevent rebleeding of esophageal varices. After a bolus injection of octreotide 50 mcg, the standard therapy (ST) group (n=34) received octreotide infusion at a rate of 50 mcg/hr until endoscopic sclerotherapy was performed within 36 hours. The same procedure was applied to another 27 patients in the maintenance therapy (MT) group in which octreotide was given at 100 mcg/8hr via the subcutaneous route after sclerotherapy for five days. In both groups, sclerotherapy was repeated on the 5th-7th day. Patients were followed for three weeks for rebleeding. Nine patients rebled in the ST group but only one patient bled in the MT group (3.7% vs. 26.5% vs. 3.7%; p<0.05). Transfusion requirement and duration of hospitalization period were similar in both groups.⁴²

Another meta-analysis showed that use of agents such as octreotide in combination with endoscopic therapy improved initial control of bleeding and 5-day hemostasis, without differences in mortality or severe adverse events, compared to endoscopic therapy alone.⁴

Corley et al. present a meta-analysis on the safety and efficacy of octreotide for esophageal variceal hemorrhage. Octreotide improved control of esophageal variceal hemorrhage compared with all alternative therapies combined (relative risk [RR], 0.63; 95% confidence interval [CI], 0.51-0.77); vasopressin/terlipressin (RR, 0.58; 95% CI, 0.42-0.81); or no additional intervention/placebo (among patients that received initial sclerotherapy/banding before randomization) (RR, 0.46; 95% CI, 0.32-0.67). Octreotide had comparable efficacy to immediate sclerotherapy for control of bleeding (RR, 0.94; 95% CI, 0.55-1.62), fewer major complications than vasopressin/terlipressin (RR, 0.31; 95% CI, 0.11-0.87), and a complication profile comparable to no intervention/placebo (RR, 1.06; 95% CI, 0.72-1.55). The results favor octreotide over vasopressin/terlipressin in the control of esophageal variceal bleeding and suggest it is a safe and effective adjunctive therapy after variceal obliteration techniques. Trials are needed to determine the optimal dose, route, and duration of octreotide treatment.⁷

Chemotherapy and/or Radiation-Induced Diarrhea

A panel of oncology experts recommends that if mild to moderate chemotherapy-induced diarrhea persists for more than 48 hours despite treatment with loperamide, it should be discontinued and the patient started on a second-line antidiarrheal agent such as octreotide. However, in the majority of mild to moderate cases of radiation-induced diarrhea, octreotide may not be sufficiently effective. Aggressive management of complicated cases of chemotherapy-induced diarrhea should involve intravenous fluids, octreotide, and antibiotics. For patients presenting with a complicated case of radiation-induced diarrhea, hospitalization may be required and octreotide therapy may or may not be appropriate.⁵

Although the somatostatin analog octreotide is currently used in the treatment of chemotherapy-induced diarrhea and secretory diarrhea associated with various disorders, its role in the management of radiation enteritis is not well defined. Yavuz, et al. performed a randomized study (n=61) that compared octreotide acetate with diphenoxylate hydrochloride plus atropine sulfate, the drug commonly used as therapy for acute radiation-induced diarrhea (ARID). Within 3 days, ARID completely resolved in 20 patients in the octreotide arm vs. only 4 in the diphenoxylate/atropine arm (p=0.002). On the diphenoxylate/atropine arm, 15/28 patients were required to discontinue pelvic radiotherapy; on the octreotide arm, 6/33 patients were required to discontinue pelvic radiotherapy for an average of 1.89 ± 0.5 and 0.45 ± 0.2 days, respectively (p=0.003). Octreotide seems to be more effective than conventional therapy with diphenoxylate and atropine in controlling ARID and eliminating the need for radiotherapy interruptions.⁴¹

Malignant Bowel Obstruction

Octreotide administration is recommended early in the diagnosis of malignant bowel obstruction due to high efficacy and tolerability.²⁹

Researchers investigated improvements in symptoms caused by gastrointestinal obstruction following administration of octreotide acetate (Sandostatin[®]) injection through steroid administration. Patients (n=37) hospitalized with malignant gastrointestinal obstructions were enrolled in the present study and 27 of them were investigated for gastrointestinal symptoms. Octreotide acetate was administered intravenously (IV) to all 27 patients. Out of them, 17 showed a marked response, 4 a good response, and 6 no response. The overall response rate was 77.8%. Octreotide acetate with a steroid was administered to 19 patients; 13 showed a marked response, 4 a good response, and 2 no response at all. Multiple logistic regression analysis showed that that steroid administration improved the efficacy of octreotide acetate after adjusting for infusion dose (p=0.03). Researchers concluded that IV administration of octreotide acetate with steroid can effectively improve gastrointestinal symptoms due to malignant gastrointestinal obstruction without adverse events.⁴⁴

A systematic review was conducted that included fifteen randomized controlled trials or observational reports with a significant number of patients (total n=281) treated with octreotide for malignant bowel obstruction. The authors reported a therapeutic success ranging between 60% and 90%. Despite the limited number of controlled studies, the large experience acquired through 20 years suggests that octreotide is the first-choice antisecretory agent for malignant bowel obstruction.²⁷

Refractory HIV/AIDS-Related Diarrhea

Agents utilized for symptomatic treatment include loperamide, diphenoxylate/atropine, paregoric, deodorized tincture of opium.³¹

Fifty-one patients with refractory uncontrolled AIDS related diarrhea were treated with octreotide in a prospective, open-label study. All fifty-one patients completed the 28 day protocol. Stool frequency and volume decreased significantly ($p < 0.001$). 41.2% (21) were considered to be partial or complete responders (decrease in daily stool volume by > 50% of initial collections or reduction to 250 mL/d). Of the responders, 67% (14 of 21) were negative for pathogens at initial screening compared to 30% (9 of 30) of nonresponders ($p < 0.01$). The study concluded that patients with refractory AIDS related diarrhea, especially those without pathogens, may respond favorably to octreotide. Limitations of this study include small sample size and lack of randomization.⁶

Although a 3 week study of 129 patients with refractory AIDS-associated diarrhea and a baseline stool weight of > 500 g/day did not show octreotide to be more effective than placebo (48% vs. 39% response, respectively), those with a baseline stool weight of 1,000-2,000 g/day did show improvement with octreotide ($p = 0.06$).³⁹

Several small reports also support octreotide's use in refractory AIDS-related diarrhea.^{13,15,24,28,35}

Unproven

Chylothorax

A Cochrane review of octreotide in the treatment of congenital or acquired chylothorax in neonates concluded that no practice recommendation can be made based on the evidence identified. Search included randomized or quasi-randomized controlled trials of octreotide in the treatment of congenital or acquired chylothorax in term or preterm neonates, with any dose, duration or route of administration. The authors reported that no randomized controlled trials were identified. Nineteen case reports of 20 neonates with chylothorax in whom octreotide was used either subcutaneously or intravenously were identified. Fourteen case reports described successful use (resolution of chylothorax), four reported failure (no resolution), and one reported equivocal results following use of octreotide. The timing of initiation, dose, duration and frequency of doses varied markedly. A prospective registry of chylothorax patients and a subsequent multicenter randomized controlled trial are needed to assess the safety and efficacy of octreotide in the treatment of chylothorax in neonates.⁹

In a retrospective review, Landvoigt examined the efficacy of octreotide in resolving chylothoraces in infants and children following cardiac surgery. Eight courses of octreotide treatment were identified in seven patients who met the inclusion criteria. The median duration of therapy was 5 days, and dosing ranged from 1 to 4 mcg/kg/hr. Treatment did not result in an overall decrease in average chest tube output after 3 days of therapy. However, in two patients (29%) the chylothoraces ultimately resolved during the octreotide infusion. Treatment was well tolerated, and no serious side effects were noted. In contrast to previously published reports, the author found that octreotide therapy for postoperative chylothoraces was successful in only a minority of cases.²³

Roehr et al. systematically reviewed the evidence on the efficacy and safety of somatostatin and octreotide in treating young children with chylothorax. Thirty-five children treated for primary or secondary chylothorax were identified. Ten of the 35 children had been given somatostatin, as an IV infusion at a median dose of 204 mcg/kg/day, for a median duration of 9.5 days. The remaining 25 children had received octreotide, either as an IV infusion at a median dose of 68 mcg/kg/day over a median 7 days, or SC at a median dose of 40 mcg/kg/day and a median duration of 17 days. A positive treatment effect was evident for both somatostatin and octreotide in the majority of reports. Minor side effects have been reported, however caution should be exercised in patients with an increased risk of vascular compromise as to avoid serious side effects. Systematic clinical research is needed to establish treatment efficacy and to develop a safe treatment protocol.³⁴

Kalomenidis performed a literature review to examine the role of somatostatin and its synthetic analog, octreotide, in the treatment of chylothorax. Several case reports and series have shown that octreotide is safe and probably effective in both

children and adults with chylothorax of different origins. The property of somatostatin and octreotide to induce leak closure is attributed to a decelerating effect on lymph flow, although their exact mechanism of action is not well defined. In successful cases, a substantial reduction of lymph drainage through the chest tube is evident within the first few days of commencing the drug, and treatment lasts for 1-2 weeks. Treatment failure has been also reported, however. Although accumulating evidence suggests that octreotide is a putative novel therapeutic intervention for chylothorax, it is imperative that randomized controlled studies are conducted in order to fully elucidate the efficacy and safety of this treatment.²¹

Dumping Syndrome

Octreotide therapy is effective in controlling severe dumping symptoms during short-term follow-up but little is known about long-term results. Didden et al. report on the long-term results of 34 patients with severe dumping syndrome treated with subcutaneous or depot intramuscular (long-acting release) octreotide. All patients had excellent initial relief of symptoms during octreotide subcutaneous therapy. However, during follow-up, 16 patients stopped therapy because of side effects (n=9) or loss of efficacy (n=7). Four patients died. Fourteen patients (41%) remained using octreotide (follow-up 93 ± 15 months), seven on octreotide subcutaneous and seven on octreotide long-acting release. The authors concluded that long-term efficacy of octreotide is much less favorable compared with short-term treatment.¹⁰

In a systematic review of seven randomized, controlled trials, Li-Ling found that although sample sizes were small in all the studies, compared with the control cases, octreotide pre-treatment resulted in significant improvement in symptoms in nearly all patients. However, long term use of octreotide for dumping syndrome was limited by severe side effects.²⁵

Vecht et al. reported the results of an open-label study including 20 patients with severe dumping symptoms after gastric surgery treated with octreotide. Mean follow-up was 37 ± 9 months (range 1-107 months). Doses of octreotide ranged from 25 to 200 mcg/day. Initial relief of symptoms was achieved in all subjects, but after three months of therapy symptom relief persisted in 80% of patients. Mean body weight increased by 2.4 ± 1.2 kg despite a significant increase in faecal fat excretion from 10 ± 2 g/24 h to 24 ± 3 g/24 h. Reasons for discontinuation of therapy were diminished efficacy in the longer term in 4 patients and side-effects in 7 patients. Biliary complications were encountered in 3 patients. Self-administration of octreotide provides an effective symptomatic treatment of severe dumping, even on the long-term. However, its use is frequently limited by the occurrence of side-effects.⁴⁰

Pancreatitis

Omata et al. performed a recent meta-analysis of double-blinded randomized controlled trials that analyzed the efficacy of somatostatin or octreotide for the prevention of post-ERCP pancreatitis and had a primary outcome measure of acute pancreatitis following ERCP. A comprehensive literature review revealed seventeen studies (n=3,818) employing a variety of methods of administration in various populations with different risks of developing post-ERCP pancreatitis. The investigators concluded that somatostatin may have significant preventive efficacy against post-ERCP pancreatitis, especially when used in appropriate diagnostic or therapeutic procedures or with high-dose administration as a 12-h infusion or a bolus. High-dose octreotide may also prevent post-ERCP pancreatitis. The efficacy of both somatostatin and octreotide in these contexts is expected to be confirmed by large high-quality randomized controlled trials in the future.³³

Zhang et al. conducted a comprehensive literature search to examine the effects of octreotide on post-endoscopic retrograde cholangiopancreatography pancreatitis (PEP). Seventeen randomized controlled trials (n=2,784) were analyzed and divided into two groups according to the total dosage of octreotide: <0.5 mg (OCT1) and ≥ 0.5 mg (OCT2). The investigators concluded that octreotide is effective in preventing post-ERCP pancreatitis and hyperamylasemia, but must be given at dosages ≥ 0.5 mg. However, there are insufficient data to determine the optimal route of administration for octreotide or its optimal timing.⁴³

Heinrich et al. performed an evidence-based analysis to assess the best available treatment for acute pancreatitis (AP), looking at the value of aprotinin, lexipafant, gabexate mesylate and octreotide treatment. Recommendations were based on the available level of evidence (A=large randomized; B=small randomized; C=prospective trial). None of the evaluated medical treatments is recommended (level A).²⁰

Uncertainties still exist about the clinical benefit of pharmacological prevention of post-endoscopic retrograde cholangiopancreatography (ERCP) pancreatitis by either antisecretory drugs such as somatostatin and its long-acting analogue octreotide, or protease inhibitors such as gabexate mesilate. Recent, large-scale prospective studies have reported a fourfold reduction in acute pancreatitis as compared to a placebo with the prophylactic administration of either gabexate mesilate or

somatostatin, whereas octreotide was found to be ineffective. An initial meta-analysis of all available controlled trials on this topic has confirmed these findings. Current literature does not support the prophylactic use of either somatostatin or gabexate mesilate for the prevention of ERCP-related pancreatic damage, even in patients deemed to be at high risk for complications.²

Persistent Hyperinsulinemic Hypoglycemia of Infancy

Long-term experience with octreotide in patients with persistent hyperinsulinemic hypoglycemia of infancy is limited, including information about possible side effects such as growth suppression. Appropriate dose and place in therapy in combination with other agents also need to be established.³

Postoperative Complications Following Pancreatic Surgery

Graham et al. conducted a prospective study of prophylactic long-acting octreotide to prevent postoperative pancreatic fistula (POPF) in high-risk patients undergoing pancreaticoduodenectomy. Sixty-eight patients evaluated for the study were divided into two groups: pancreatic ducts ≤ 3 mm (high risk, $n=36$) and those with ducts > 3 mm (low risk, $n=32$). High-risk patients were treated preoperatively with depot octreotide and begun on an intravenous drip for 24 hours. Low-risk patients underwent pancreaticoduodenectomy without pharmacologic intervention. In contrast, the control cohort represented 106 retrospectively analyzed patients who underwent a pancreaticoduodenectomy without depot octreotide injection without regard to low- or high-risk status. Overall, POPF was 11 of 68 (16%). Nine of 36 high risk patients treated with depot octreotide developed POPF (25%), and 2 of 32 low risk patients developed POPF (6%). In the control cohort of high-risk patients, 9 of 44 (20%) and 3 of 62 (5%) low-risk patients developed POPF ($p=0.628$ when comparing the development of POPF in high-risk patients with or without pharmacologic intervention). The authors concluded that prophylactic use of depot octreotide in high-risk patients does not result in reduced incidence of POPF. However, duct size has a significant impact on the occurrence of POPF.¹⁸

A recent Cochrane review of somatostatin analogues (SSAs) for pancreatic surgery concluded that SSAs reduce perioperative complications but do not reduce perioperative mortality. In those undergoing pancreatic surgery for malignancy, they shorten hospital stay. Further adequately powered trials with low risk of bias are necessary. Based on the current available evidence, somatostatin and its analogues are recommended for routine use in patients undergoing pancreatic resection for malignancy. There is currently no evidence to support their routine use in pancreatic surgeries performed for other indications.¹⁹

In a meta-analysis by Zeng et al., eight studies were reviewed to evaluate the efficacy of somatostatin and its analogues in the prevention of postoperative complications after pancreaticoduodenectomy. The use of somatostatin or its analogues did not significantly benefit for reducing the incidence of pancreatic fistula (odds ratio [OR] 95% confidence interval [CI], 0.64-1.37; $p=0.73$), total pancreas-specific postoperative complications (OR 95% CI, 0.63-1.42; $p=0.79$), delayed gastric emptying (OR 95% CI, 0.50-1.78; $p=0.86$), total complication (OR 95% CI, 0.73-1.70; $p=0.61$), mortality (OR 95% CI, 0.59-7.72; $p=0.97$) and length of postoperative hospital stay (weighted mean difference 95% CI, -7.74 to 4.47; $p=0.60$). The use of somatostatin and its analogues does not significantly reduce postoperative complications after pancreaticoduodenectomy.⁴²

Several clinical trials have evaluated the use of octreotide to prevent the development of pancreatic fistula after pancreatic surgery with conflicting recommendations. Alghamdi et al. conducted a meta-analysis of seven randomized controlled trials ($n=1,359$), reporting comparisons between octreotide and a control. The primary outcome was the incidence of postoperative pancreatic fistula, and the secondary outcome was the postoperative mortality. In these studies, sample sizes ranged from 75 to 252 patients. In total, 679 patients were given octreotide and 680 patients formed the control group. Perioperative octreotide is associated with a significant reduction in the incidence of pancreatic fistula after elective pancreatic surgery, with a relative risk of 0.59 (95% confidence interval 0.41-0.85, $p=0.004$). However, this risk reduction was not associated with a significant difference in postoperative mortality ($p>0.05$). Further studies are warranted to confirm the results of this meta-analysis and to define which patient subgroups might benefit the most from prophylactic octreotide administration.¹

Short Bowel Syndrome

Nehra et al. assessed the effects of octreotide acetate depot in patients with short bowel syndrome by conducting a 15-week, prospective, open-label study of eight patients (five women and three men; mean age 52 yr., range 37-72 yr.). Treatment with octreotide acetate depot significantly increased small bowel transit time ($p=0.03$). Changes in body weight, urine volume, stool weight, fecal fat excretion, stool sodium and potassium excretion, or gastric emptying rate were highly variable, and no overall significance was observed. Octreotide acetate depot for 15 weeks significantly prolonged small bowel transit time. However, octreotide acetate treatment needs to be assessed further in multicenter studies assessing dose, frequency of administration and a larger sample size.³⁰

Signifor

Petersenn et al. conducted a randomised, double-blind study, to investigate the safety and efficacy of pasireotide in adult patients with persistent/recurrent or de novo Cushing's disease.⁵⁰ Patients with mean urinary free cortisol at or below the upper limit of normal or clinical benefit at month 12 could continue receiving pasireotide during this open-ended, open-label phase. For the 16 patients that received 5 years of pasireotide treatment, the median (95% confidence interval) percentage change from baseline in mean urinary free cortisol was -82.6% (-89.0, -41.9) and -81.8% (-89.8, -67.4) at months 12 and 60. Eleven patients had mean urinary free cortisol \leq upper limit of normal at month 60. Improvements in clinical signs were sustained during long-term treatment. The safety profile of pasireotide at 5 years was similar to that reported after 12 months. Fifteen of 16 patients experienced a hyperglycaemia-related adverse event; glycated haemoglobin levels were stable between months 6 and 60. Adverse events related to hyperglycaemia, bradycardia, gallbladder/biliary tract, and liver safety were most likely to first occur by month 6, and severity did not tend to worsen over time. The authors conclude that the use of pasireotide is an effective long-term therapy for some patients with Cushing's disease.

In a double-blind, phase 3 study, Colao et al evaluated the efficacy of pasireotide on urinary free cortisol.⁵¹ Adults with Cushing's disease and a urinary free cortisol level of at least 1.5 times the upper limit of the normal were randomly assigned to receive subcutaneous pasireotide at a twice daily dose of 600 μ g or 900 μ g. At month 3, patients with urinary free cortisol 2 times the upper limit of the normal range or less, and not exceeding their baseline level remained on their randomly assigned dose. All other patients received an increase in dose of 300 μ g twice daily. The primary end point was a urinary free cortisol level at or below the upper limit of the normal at 6 months without an increased dose. Open-label treatment continued for a total of 12 months. The primary endpoint was met by 12 of 82 patients in the 600- μ g group and 21 of 80 patients in the 900- μ g group. The median urinary free cortisol level decreased by approximately 50% by month 2 and remained stable in both groups. Patients with baseline levels not exceeding 5 times the upper limit of the normal more frequently achieved a normal urinary free cortisol level than patients with higher baseline levels. Serum and salivary cortisol and plasma corticotropin levels decreased, as well as clinical signs and symptoms of Cushing's disease. Hyperglycemia-related adverse events occurred in 118 of 162 patients. Additionally, other adverse events were similar to those associated with other somatostatin analogues. Even with declines in cortisol levels, blood glucose and glycated hemoglobin levels increased shortly after the initiation of treatment and then stabilized; glucose-lowering medication was initiated in 74 of 162 patients. The authors concluded that there was a significant decrease in cortisol levels in patients receiving pasireotide with Cushing's disease. This supports its potential use as a targeted treatment for corticotropin secreting pituitary adenomas.

Signifor LAR

In this double-blind extension to a multicenter, 12-month, Phase III core study, Sheppard et al evaluated the efficacy and safety of pasireotide LAR and octreotide LAR after up to 26 months' treatment.⁵⁶ Patients with GH <2.5 μ g/L and IGF-1 $\leq 1 \times$ ULN at month 12, or patients considered to be experiencing clinical benefit, were eligible to continue receiving their randomized therapy in this extension. Efficacy and safety were evaluated for up to 26 months. Overall, 120 patients who completed the core study continued receiving pasireotide LAR or octreotide LAR in this extension study. At month 25, biochemical control, defined as GH <2.5 μ g/L and normal IGF-1, was achieved by 48.6% and 45.7% of patients in the pasireotide LAR and octreotide LAR arms respectively. In total, 74.7 % of pasireotide LAR and 71.6 % of octreotide LAR patients had tumor volume decrease ≥ 20 % from baseline to month 26. Most adverse events were mild or moderate. Hyperglycemia-related adverse events were seen in 62.9 and 25.0 % of pasireotide LAR and octreotide LAR patients, respectively. The authors conclude that GH and IGF-1 suppression is maintained for up to 25 months during pasireotide LAR treatment. Additionally, they conclude that the safety profile of pasireotide LAR is typical of a somatostatin analogue, except for the frequency and degree of hyperglycemia.

In the PAOLA trial, Gadelha et al evaluated the efficacy and safety of pasireotide long-acting release compared with octreotide or lanreotide in patients with inadequately controlled acromegaly.⁵⁵ In this randomized, phase 3 trial, patients 18 years and older with acromegaly who were inadequately controlled, and had received 30 mg octreotide long-acting or 120 mg lanreotide as monotherapy for 6 months or longer were enrolled. Patients were randomly assigned in a 1:1:1 ratio to receive 40 mg pasireotide long-acting release once every 28 days, 60 mg pasireotide long-acting release once every 28 days, or continued treatment with octreotide or lanreotide (active control) for 24 weeks. Patients were stratified according to previous treatment and growth hormone concentrations at screening. The primary endpoint was number of patients achieving biochemical control, defined as mean growth hormone concentration less than 2.5 μ g/L and normalized IGF-1 concentration. Enrolled patients were randomly assigned to pasireotide 40 mg, pasireotide 60 mg, or active control groups. At 24 weeks, ten (15%) patients in the pasireotide 40 mg group and 13 (20%) patients in the pasireotide 60 mg group achieved biochemical control, compared with

no patients in the active control group. The most common adverse events were hyperglycemia, diabetes, and diarrhea. The authors concluded that pasireotide provides superior efficacy compared with continued treatment with octreotide or lanreotide.

Coloa et al evaluated the superiority of pasireotide LAR over octreotide LAR in medically naive patients with acromegaly in a multicenter prospective, randomized, double-blind study.⁵⁴ Enrollment included 358 patients with medically naive acromegaly. Patients either had previous pituitary surgery but no medical treatment or were de novo with a visible pituitary adenoma on magnetic resonance imaging. In the study, patients receiving pasireotide LAR 40 mg/28 days were compared to patients receiving octreotide LAR 20 mg/28 days for 12 months. At months 3 and 7, patients who had IGF-1 levels above the upper limit of normal had the option of having their doses titrated to pasireotide LAR 60mg or octreotide LAR 30mg. The primary outcome was the proportion of patients in each treatment group achieving biochemical control, defined at GH 2.5 µg/L and normal IGF-1 at month 12. Biochemical control was achieved by significantly more pasireotide LAR patients than octreotide LAR patients. In pasireotide LAR and octreotide LAR patients, respectively, 38.6% and 23.6% (P.002) achieved normal IGF-1, and 48.3% and 51.6% achieved GH 2.5 µg/L. 31.0% of pasireotide LAR and 22.2% of octreotide LAR patients who did not achieve biochemical control did not receive the recommended dose increase. Hyperglycemia-related adverse events were more common with pasireotide LAR (57.3% vs 21.7%). The authors conclude that pasireotide LAR demonstrated superior efficacy over octreotide LAR and is a viable new treatment option for acromegaly.

Somatuline Depot

Salvatori et al. evaluated the 2-year effectiveness and safety of lanreotide depot/autogel (LAN), as well as treatment convenience and acromegaly symptom relief, from the Somatuline[®] Depot for Acromegaly (SODA) registry, in a post-marketing, open-label, observational, multicenter, registry study. Patients with acromegaly treated with LAN were eligible for enrollment. The following items were collected as part of the registry: demographics, LAN dose, extended dosing interval (EDI), insulin-like growth factor 1 (IGF-1), growth hormone (GH), glycated hemoglobin, adverse events (AEs), injection convenience, as well as symptom data. IGF-1 levels below age- and gender-adjusted upper normal limit (ULN) were achieved in 71.2% at month (M) 12 and 74.4% at M24; GH ≤2.5 µg/L in 83.3% at M12 and 80.0% at M24; GH <1.0 µg/L in 61.7% at M12 and 61.4% at M24. Both IGF-1 < ULN and GH ≤2.5 µg/L were achieved in 65.0% at M12 and 54.8% at M24; both IGF-1 < ULN and GH < 1.0 µg/L were achieved in 51.7 and 42.9% at M12 and M24, respectively. EDI regimen was 5.0% at baseline and 12.0% at M24. Acromegaly symptoms appeared stable or improved at 24 months. Arthralgia was the most common adverse event at 25.7%. At M24, 73.1% of patients rated LAN as convenient. The authors conclude that SODA indicates 2-year biochemical control with majority of patients achieving both IGF-1 < ULN and GH ≤2.5 µg/L. In addition, LAN was generally well tolerated during the 24 month observation period with no new or unexpected safety signals reported.

Chanson et al. conducted an open-label, multicenter, phase III, 48-week trial to assess the efficacy and safety of 48 weeks titrated dosing of lanreotide.⁴⁸ Patients with active acromegaly (IGF-I levels > 1-3 times upper limit of age-adjusted normal range) were recruited and received 12 injections of lanreotide Autogel at 28-day intervals. Dosing during the 16-week fixed-dose phase was 90 mg; in the 32-week dose-titration phase, patients received 60, 90 or 120 mg based on GH and IGF-I levels. At the end of the study, an intention-to-treat analysis was performed to determine the proportion of patients with normalized age-adjusted IGF-I levels. GH levels, clinical acromegaly signs, and safety were secondary measures. Fifty-seven of 63 patients completed the study. Lanreotide resulted in normalized age-adjusted IGF-I levels in 27 patients (43%, 95% CI 31–55). Mean GH levels decreased from 6.2 to 1.5 µg/l at study end, with 53 of 62 patients (85%) having GH levels ≤ 2.5 µg/l (95% CI 76.7–94.3) and 28 of 62 patients (45%) with levels < 1 µg/l (95% CI 32.8–57.6). Twenty-four (38%) had both normal IGF-I levels and GH levels ≤ 2.5 µg/l. Symptoms of acromegaly reduced significantly in most patients during the study. The most common adverse events were gastrointestinal in nature. The authors concluded that at 48 weeks lanreotide treatment, titrated for optimal hormonal control, controlled IGF-I and GH levels effectively, reduced acromegaly symptoms and was well tolerated.

Professional Societies

Acromegaly

Endocrine Society & European Society of Endocrinology (2014)⁴⁶

The Task Force of the Endocrine Society Clinical Guidelines Subcommittee published an evidence based guideline regarding the evaluation and management of acromegaly. The guidelines state (Strong recommendations= the number 1, weak recommendations= the number 2; quality of evidence):

- Preoperative use of somatostatin analogues to reduce surgical risk from severe comorbidities (2; very low quality)

- The use of somatostatin analogues (e.g., octreotide) or pegvisomant in a patient with significant disease, as the initial adjuvant medical therapy (2; low quality).
- The addition of pegvisomant or cabergoline in a patient with inadequate response to a somatostatin analogue (2; low quality).
- The use of somatostatin analogue as primary therapy in a patient who cannot be cured by surgery, has extensive cavernous sinus invasion, does not have chiasmal compression, or is a poor surgical candidate (2; moderate quality).
- Discontinue long acting somatostatin analogue formulations and pegvisomant approximately 2 months before conceiving, with use of short acting octreotide as necessary until conception (2; low quality).

American Association of Clinical Endocrinologists

The recently updated guidelines of the American Association of Clinical Endocrinologists for the diagnosis and treatment of acromegaly list the somatostatin analogues (SSAs) octreotide and lanreotide with a Grade A recommendation. (Grade A = one or more conclusive level 1 publications exist demonstrating benefit >> risk; recommendation is based upon strong evidence; recommendation is considered first-line therapy.)²²

Bleeding Gastroesophageal Varices

American College of Gastroenterology

The American Association for the Study of Liver Diseases and the Practice Parameters Committee of the American College of Gastroenterology's Practice Guidelines for the Prevention and Management of Gastroesophageal Varices and Variceal Hemorrhage in Cirrhosis recommend octreotide as a useful adjunct to endoscopic therapy. Pharmacological therapy (somatostatin or its analogues) should be initiated as soon as variceal hemorrhage is suspected and continued for 3-5 days after diagnosis is confirmed (Class I, Level A). (Class I - conditions for which there is evidence and/or general agreement that a given diagnostic evaluation, procedure or treatment is beneficial, useful, and effective. Level A - data derived from multiple randomized clinical trials or meta-analyses.)¹⁶

Refractory HIV/AIDS-Related Diarrhea

American College of Gastroenterology

The American College of Gastroenterology's Practice Guidelines on Acute Infectious Diarrhea in Adults consider octreotide an effective alternative for otherwise refractory cases of AIDS-induced diarrhea. Whereas the drug is best used in pathogen-negative diarrhea, it may be useful in some patients with microsporidiosis and possibly other otherwise non treatable conditions. Because octreotide must be administered by injection and it is quite expensive, it should be considered a last resort to symptomatic management.¹²

American Gastroenterological Association

An American Gastroenterological Association Technical Review on Acute Pancreatitis lists somatostatin and octreotide as pharmacological options to limit pancreatic secretion. However, the review states that the data supporting the use of these agents is not very convincing. Of note, the largest single randomized trial (by far) of octreotide in 302 patients with moderate to severe acute pancreatitis found absolutely no effect on mortality, organ failure, or secondary infections. Somatostatin is not easily available in the United States, and the data on octreotide are controversial, so neither can currently be recommended as routine management for acute pancreatitis.¹⁴

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Sandostatin is indicated for the following:³⁶

- To reduce blood levels of growth hormone and IGF-I (somatomedin C) in acromegaly patients who have had inadequate response to or cannot be treated with surgical resection, pituitary irradiation and bromocriptine mesylate at maximally tolerated doses
- For symptomatic treatment of patients with metastatic carcinoid tumors where it suppresses or inhibits the severe diarrhea and flushing episodes associated with the disease
- In the treatment of the profuse watery diarrhea associated with VIP-secreting tumors

Sandostatin LAR Depot is indicated in patients in whom initial treatment with Sandostatin subcutaneous injection has been shown to be effective and tolerated for:³⁷

- Acromegaly:
 - Long-term maintenance therapy in acromegalic patients who have had an inadequate response to surgery and/or radiotherapy, or for whom surgery and/or radiotherapy is not an option. The goal of treatment in acromegaly is to reduce GH and IGF-1 levels to normal
- Carcinoid Tumors
- Long-term treatment of the severe diarrhea and flushing episodes associated with metastatic carcinoid tumors. Vasoactive Intestinal Peptide Tumors (VIPomas):
 - Long-term treatment of the profuse watery diarrhea associated with VIP-secreting tumors

Signifor is indicated for the treatment of adult patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative.⁴⁹

Signifor LAR is indicated for the treatment of:⁵³

- Patients with acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option
- Patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative

Somatuline Depot is indicated for:⁴⁷

- The long-term treatment of acromegalic patients who have had an inadequate response to or cannot be treated with surgery and/or radiotherapy
- The treatment of patients with unresectable, well- or moderately differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs) to improve progression-free survival
- The treatment of adults with carcinoid syndrome; when used, it reduces the frequency of short-acting somatostatin analogue rescue therapy

Centers for Medicare and Medicaid Services (CMS)

Medicare does not have a National Coverage Determination (NCD) for Sandostatin® (octreotide acetate) and Sandostatin® LAR (octreotide acetate LAR). Local Coverage Determinations (LCDs)/Local Coverage Articles (LCAs) exist. See the LCDs/LCAs for [Octreotide Acetate for Injectable Suspension \(Sandostatin LAR depot\)](#) and [Chemotherapy Administration](#).

Medicare does not have NCDs for Somatuline® Depot (lanreotide), Signifor® (pasireotide) and Signifor® LAR (pasireotide diaspertate). LCDs/LCAs do not exist at this time.

In general, Medicare may cover outpatient (Part B) drugs that are furnished "incident to" a physician's service provided that the drugs are not usually self-administered by the patients who take them. Refer to the [Medicare Benefit Policy Manual, Chapter 15, §50 - Drugs and Biologicals](#). (Accessed October 2, 2020)

References

1. Alghamdi AA, Jawas AM, Hart RS. Use of octreotide for the prevention of pancreatic fistula after elective pancreatic surgery: a systematic review and meta-analysis. *Can J Surg*. 2007 Dec;50(6):459-66.
2. Andriulli A, Caruso N, Quitadamo M, et al. Antisecretory vs. antiprotease drugs in the prevention of post-ERCP pancreatitis: the evidence-based medicine derived from a meta-analysis study. *JOP*. 2003 Jan;4(1):41-8.
3. Aynsley-Green, Hussain K, Hall J, et al. Practical management of hyperinsulinism in infancy. *Arch Dis Child Neonatal Ed*. 2000;82(2):F98-F107.
4. Banares R, Albillos A, Rincon D, et al. Endoscopic treatment versus endoscopic plus pharmacologic treatment for acute variceal bleeding: A meta-analysis. *Hepatology*. 2002;35(3):609-15.
5. Benson AB 3rd, Ajani JA, Catalano RB, et al. Recommended guidelines for the treatment of cancer treatment-induced diarrhea. *J Clin Oncol*. 2004 Jul 15;22(14):2918-26.

6. Cello JP, Grendell JH, Basuk, P, et al. Effect of octreotide on refractory AIDS-associated diarrhea. A prospective, multicenter clinical trial. *Annals Internal Medicine*. 1991;115(9):705-10.
7. Corley DA, Cello JP, Adkisson W, et al. Octreotide for acute esophageal variceal bleeding: a meta-analysis. *Gastroenterology*. 2001 Mar;120(4):946-54.
8. D'Amico G, Pietrosi G, Tarantino I, Pagliaro L. Emergency sclerotherapy versus vasoactive drugs for variceal bleeding in cirrhosis: a Cochrane meta-analysis. *Gastroenterology*. 2003 May;124(5):1277-91.
9. Das A, Shah PS. Octreotide for the treatment of chylothorax in neonates. *Cochrane Database of Systematic Reviews* 2010, Issue 9. Art. No.: CD006388. DOI: 10.1002/14651858.CD006388.pub2.
10. Didden P, Penning C, Masclee AA. Octreotide therapy in dumping syndrome: Analysis of long-term results. *Aliment Pharmacol Ther*. 2006 Nov 1;24(9):1367-75.
11. Dolan JT, Miltenburg DM, Granchi TS, et al. Treatment of metastatic breast cancer with somatostatin analogues—a meta-analysis. *Ann Surg Oncol*. 2001 Apr;8(3):227-33.
12. DuPont HL. Guidelines on Acute Infectious Diarrhea in Adults. *Am J Gastroenterol*. 1997 Nov;92(11):1962-75.
13. Fanning M, Monte M, Sutherland LR, et al. Pilot study of sandostatin (octreotide) therapy of refractory HIV-associated diarrhea. 1991;36(4):476-80.
14. Forsmark CE and Baillie J. AGA Institute Technical Review on Acute Pancreatitis. *Gastroenterology* 2007;132:2022-2044.
15. Garcia Compean D, Ramos Jimenez J, Guzman de la Garza F, et al. Octreotide therapy of large-volume refractory AIDS-associated diarrhea: a randomized controlled trial. *AIDS*. 1994;8(11):1563-7.
16. Garcia-Tsao G, Sanyal AJ, Grace ND, et al. Prevention and management of gastroesophageal varices and variceal hemorrhage in cirrhosis. *Am J Gastroenterol*. 2007 Sep;102(9):2086-102.
17. Gotzsche PC, Hrobjartsson A. Somatostatin analogues for acute bleeding oesophageal varices. *Cochrane Database Syst Rev*. 2008, Issue 3. Art. No.: CD000193. DOI: 10.1002/14651858.CD000193.pub3.
18. Graham JA, Johnson LB, Haddad N, et al. A prospective study of prophylactic long-acting octreotide in high-risk patients undergoing pancreatoduodenectomy. *The American Journal of Surgery*. 2011;201(4):481-485.
19. Gurusamy KS, Koti R, Fusai G, Davidson BR. Somatostatin analogues for pancreatic surgery. *Cochrane Database of Systematic Reviews* 2010, Issue 2. Art. No.: CD008370. DOI: 10.1002/14651858.CD008370.
20. Heinrich S, SchM, Rousson V, Clavien PA. Evidence-based treatment of acute pancreatitis: a look at established paradigms. *Ann Surg*. 2006 Feb;243(2):154-68.
21. Kalomenidis I. Octreotide and chylothorax. *Curr Opin Pulm Med*. 2006 Jul;12(4):264-7.
22. Katznelson L, Atkinson JLD, Cook DM, et al. American Association of Clinical Endocrinologists Medical Guidelines for Clinical Practice for the Diagnosis and Treatment of Acromegaly – 2011 Update. *Endocr Pract*. 2011;17(Suppl 4).
23. Landvoigt MT, Mullett CJ. Octreotide efficacy in the treatment of chylothoraces following cardiac surgery in infants and children. *Pediatr Crit Care Med*. 2006 May;7(3):245-8.
24. Liberti A, Bisogno A, Izzo E. Octreotide treatment in secretory and cryptosporidial diarrhea in patients with acquired immunodeficiency syndrome (AIDS): clinical evaluation. *J Chemother*. 1992;4(5):303-5.
25. Li-Ling J, Irving M. Therapeutic value of octreotide for patients with severe dumping syndrome—a review of randomised controlled trials. *Postgrad Med J*. 2001 Jul;77(909):441-2.
26. Melmed S, Barkan A, Molitch M, et al. Guidelines for Acromegaly Management: An Update. *J Clin Endocrinol Metab*. May 2009, 94 (5):1509-1517.
27. Mercadante S, Porzio G. Octreotide for malignant bowel obstruction: Twenty years after. *Crit Rev Oncol Hematol*. 2012 Sep;83(3):388-92. Epub 2012 Jan 25.
28. Moroni M, Esposito R, Cernuschi M, et al. Treatment of AIDS-related refractory diarrhoea with octreotide. *Digestion*. 1993;54 Suppl 1:30-2.
29. The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). Palliative Care. Version 1.2020. Accessed October 6, 2020.

30. Nehra V, Camilleri M, Burton D, et al. An open trial of octreotide long-acting release in the management of short bowel syndrome. *Am J Gastroenterol*. 2001 May;96(5):1494-8.
31. New York State Department of Health AIDS Institute. Gastrointestinal Complications of HIV. October 2006. Accessed September 7, 2012.
32. American Association of Clinical Endocrinologist (AACE) medical guidelines for clinical practice for the diagnosis and treatment of acromegaly. *Endocrine Practice*. 2004; 10(3): 213-225.
33. Omata F, Deshpande G, Tokuda Y, et al. Meta-analysis: somatostatin or its long-acting analogue, octreotide, for prophylaxis against post-ERCP pancreatitis. *J Gastroenterol* 2010; 45:885-895.
34. Roehr CC, Jung A, Proquitté H, et al. Somatostatin or octreotide as treatment options for chylothorax in young children: a systematic review. *Intensive Care Med*. 2006 May;32(5):650-7. Epub 2006 Mar 11.
35. Romeu J, Miro JM, Sirera G, et al. Efficacy of octreotide in the management of chronic diarrhoea in AIDS. *AIDS*. 1991;5(12):1495.
36. Sandostatin [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corporation, June 2020.
37. Sandostatin LAR Depot [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corporation, April 2019.
38. Shah HA, Mumtaz K, Jafri W, et al. Sclerotherapy plus octreotide versus sclerotherapy alone in the management of gastro-oesophageal variceal hemorrhage. *J Ayub Med Coll Abbottabad*. 2005 Jan-Mar;17(1):10-4.
39. Simon DM, Cello JP, Valenzuela J, et al. Multicenter trial of octreotide in patients with refractory acquired immunodeficiency syndrome-associated diarrhea. *Gastroenterology*. 1995 Jun;108(6):1753-60.
40. Vecht J, Lamers CBHW, and Masclee AAM. Long-term results of octreotide-therapy in severe dumping syndrome. *Clinical Endocrinology*. 1999;51:619-624.
41. Yavuz MN, Yavuz AA, Aydin F, et al. The efficacy of octreotide in the therapy of acute radiation-induced diarrhea: a randomized controlled study. *Int J Radiat Oncol Biol Phys*. 2002 Sep 1;54(1):195-202.
42. Yücesoy M, Başkol M, Keklik M, et al. Efficacy of five days of subcutaneous octreotide treatment after sclerotherapy in preventing rebleeding from esophageal varices. *Turk J Gastroenterol*. 2004 Sep;15(3):137-43.
43. Zeng Q, Zhang Q, Han S, et al. Efficacy of somatostatin and its analogues in prevention of postoperative complications after pancreaticoduodenectomy: a meta-analysis of randomized controlled trials. *Pancreas*. 2008 Jan;36(1):18-25.
44. Zhang Y, Chen QB, Gao ZY, et al. Meta-analysis: octreotide prevents post-ERCP pancreatitis, but only at sufficient doses. *Aliment Pharmacol Ther* 2009;29:1155-64.
45. Murakami H, Matsumoto H, Nakamura M, et al. Octreotide acetate-steroid combination therapy for malignant gastrointestinal obstruction. *Anticancer Res*. 2013 Dec;33(12):5557-60.
46. Katznelson L, Laws Jr ER, Melmed S, et al. Acromegaly: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*, November 2014, 99(11):3933–3951.
47. Somatuline Depot [prescribing information]. Cambridge, MA: Ipsen Biopharmaceuticals, Inc, June 2019.
48. Chanson P, Barson-Chazott, Kuhn JM et al. Control of IGF-I levels with titrated dosing of lanreotide autogel over 48 weeks in patients with acromegaly. *Clin Endocrinol (OXF)*. 2008; 69:299-305.
49. Signifor [prescribing information]. Lebanon, NJ: Recordati Rare Diseases Inc, June 2020.
50. Petersenn S, Salgado LR, Schopohl J, Portocarrero-Ortiz L, Arnaldi G, Lacroix A, Scaroni C, Ravichandran S, Kandra A, Biller BMK. Long-term treatment of Cushing's disease with pasireotide: 5-year results from an open-label extension study of a Phase III trial. *Endocrine*. 2017 Jul;57(1):156-165.
51. Colao A, Petersenn S, et al. A 12-Month Phase 3 Study of Pasireotide in Cushing's Disease. *N Engl J Med* 2012;366:914-24.
52. Salvatori R, Gordon MB, Woodmansee WW, Ioachimescu AG, Carver DW, Mirakhor B, Cox D, Molitch ME. A multicenter, observational study of lanreotide depot/autogel (LAN) in patients with acromegaly in the United States: 2-year experience from the SODA registry. *Pituitary*. 2017 Jul 24.
53. Signifor LAR [prescribing information]. East Lebanon, NJ: Recordati Rare Diseases Inc., June 2002.

54. Colao A, Bronstein MD, Freda P, et al. Pasireotide versus octreotide in acromegaly: a head-to-head superiority study. *J Clin Endocrinol Metab.* 2014;99(3):791–799.
55. Gadelha MR, Bronstein MD, Brue T, et al. Pasireotide versus continued treatment with octreotide or lanreotide in patients with inadequately controlled acromegaly (PAOLA): a randomised, phase 3 trial. *Lancet Diabetes Endocrinol.* 2014;2:875–884.
56. Sheppard M, Bronstein MD, Freda P, et al. Pasireotide LAR maintains inhibition of GH and IGF-1 in patients with acromegaly for up to 25 months: results from the blinded extension phase of a randomized, double-blind, multicenter, phase III study. *Pituitary.* 2014;18:385–394.

Policy History/Revision Information

Date	Summary of Changes
08/01/2021	<p>Application</p> <ul style="list-style-type: none"> Added language to indicate this policy does not apply to the states of Indiana and North Carolina
02/01/2021	<p>Application</p> <ul style="list-style-type: none"> Reformatted content Added language to indicate this policy does not apply to the state of Kentucky; refer to the state-specific policy version
12/01/2020	<p>Application</p> <ul style="list-style-type: none"> Added language to indicate this policy does not apply to the state of Louisiana; refer to the state-specific policy version <p>Supporting Information</p> <ul style="list-style-type: none"> Updated <i>CMS</i> and <i>References</i> sections to reflect the most current information Archived previous policy version CS2019D0036O

Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. The UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.