

Meniscus Implant and Allograft

Policy Number: SURGERY 096.19
Effective Date: January 1, 2023

[Instructions for Use](#)

Table of Contents	Page
Coverage Rationale	1
Documentation Requirements	1
Definitions	2
Applicable Codes	2
Description of Services	2
Clinical Evidence	3
U.S. Food and Drug Administration	5
References	5
Policy History/Revision Information	6
Instructions for Use	6

Related Policies

- [Articular Cartilage Defect Repairs](#)
- [Unicondylar Spacer Devices for Treatment of Pain or Disability](#)

Coverage Rationale

[Meniscus Allograft Transplantation](#) (MAT) with human cadaver tissue is proven and medically necessary for replacement of major meniscus loss due to trauma or previous meniscectomy when all of the following criteria are met:

- Individuals who are skeletally mature with documented closure of growth plates;
- Disabling knee pain causing [Functional Impairment](#) that is refractory to conservative treatment;
- Absence of more than half of the meniscus due to surgery or injury or the presence of a tear that cannot be repaired;
- Radiographic criteria established by a standing anteroposterior (AP) view demonstrates all of the following:
 - Normal alignment or correctable varus or valgus deformities;
 - No osteophytes or marginal osteophytes;
 - No irreparable articular cartilage defects;
 - No significant joint space narrowing;
- Normal knee biomechanics, or alignment and stability achieved concurrently with meniscal Transplantation;
- Minimal to absent degenerative changes in surrounding articular cartilage (Outerbridge Grade II or less);
- No evidence of active inflammatory arthritis or systemic arthritis

[Collagen Meniscus Implants](#) (CMI) are unproven and not medically necessary for treating meniscus injuries or tears due to insufficient evidence of efficacy.

Documentation Requirements

Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The documentation requirements outlined below are used to assess whether the member meets the clinical criteria for coverage but do not guarantee coverage of the service requested.

CPT Code*	Required Clinical Information
Meniscus Implant and Allograft	
29868	<p>Medical notes documenting the following, when applicable:</p> <ul style="list-style-type: none"> • Complete report(s) of diagnostic imaging (MRI, CT scan, x-rays, and bone scan) Note: For pediatric age, indicate status of growth plates • Condition requiring procedure • Severity of pain and details of functional disability(ies) interfering with activities of daily living (ADL) • Pertinent physical examination of the relevant joint • Co-morbid medical condition(s) • Prior therapies/treatments tried, failed, or contraindicated; include the dates and reason for discontinuation • Degree of degenerative changes in surrounding cartilage as measured by the Outerbridge Grade • If the location being requested is an inpatient stay, provide documentation to support site of care • Physician's treatment plan including pre-op discussion

*For code description, refer to the [Applicable Codes](#) section.

Definitions

Collagen Meniscal Implant (CMI): Resorbable and biocompatible Type I collagen matrix that was developed to restore the segmental loss of meniscal tissue in the knee. It consists of a porous cross-linked matrix scaffold that allows for the ingrowth of the body's own cells. (Warth and Rodkey, 2015)

Functional or Physical Impairment: A Physical or Functional or physiological Impairment causes deviation from the normal function of a tissue or organ. This results in a significantly limited, impaired, or delayed capacity to move, coordinate actions, or perform physical activities and is exhibited by difficulties in one or more of the following areas: physical and motor tasks; independent movement; performing basic life functions. (World Health Organization and World Bank (WHO), 2011)

Meniscal Allograft Transplantation (MAT): Transplant of the meniscus of the knee, which separates the thigh bone (femur) from the lower leg bone (tibia). The worn or damaged meniscus is removed and is replaced with a new one from a donor. The meniscus to be transplanted is taken from a cadaver, and, as such, is known as an allograft. (AAOS, 2021)

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies may apply.

CPT Code	Description
29868	Arthroscopy, knee, surgical; meniscal transplantation (includes arthrotomy for meniscal insertion), medial or lateral

CPT® is a registered trademark of the American Medical Association

HCPCS Code	Description
G0428	Collagen meniscus implant procedure for filling meniscal defects (e.g., CMI, collagen scaffold, Menaflex)

Description of Services

Meniscal cartilage is an integral structural component of the human knee, functioning to absorb shocks and providing load sharing, joint stability, congruity, proprioception, and lubrication and nutrition of the cartilage surfaces. Allografts are grafts of tissues made available from a live person or a human cadaver. Allografts from cadavers avoid morbidity from harvesting tissue

from a different site on the person requiring meniscus repair. The goal of meniscal allograft transplantation is to restore knee function and prevent further joint degeneration by replacing the damaged or destroyed meniscus with allograft tissue having similar properties as the damaged tissue.

The Collagen Meniscal Implant (CMI) is an implant derived from bovine collagen used to treat acute or chronic advanced meniscal loss or damage with the intent of relieving symptoms and preventing joint degeneration. The CMI is a flexible, sickle-shaped disc that mimics the shape of the native meniscus and is attached arthroscopically to native tissue with suture. The CMI is meant to serve as a temporary template to support migration of the host's cells to the meniscal deficiency, restoring meniscal volume and function.

Clinical Evidence

Collagen Meniscus Implants (CMIs)

There is insufficient evidence to demonstrate the efficacy of collagen meniscus implants for treating meniscus injuries or tears. Robust randomized controlled trial studies are needed along with long-term outcomes to establish the safety and efficacy of this procedure.

Veronesi et al. (2021) conducted a systematic review to collect and evaluate the available evidence on biosynthetic scaffolds for meniscus regeneration both in vivo and in clinical studies. Three databases were searched: A total of 46 in vivo preclinical studies and 30 clinical studies were identified. Sixteen natural, 15 synthetic, and 15 hybrid scaffolds were studied in vivo. Among them, only 2 scaffolds were evaluated in clinical studies: The Collagen Meniscus Implant was evaluated in 11 studies, and the polyurethane-based scaffold Actifit® was evaluated in 19 studies. Although positive outcomes were described in the short- to mid-term, the number of concurrent procedures and the lack of randomized trials are the major limitations of the available clinical literature. According to the authors, current solutions offer a significant but incomplete clinical improvement, and the regeneration potential is still unsatisfactory. Authors Rodkey et al., 2008; Bulgheroni et al., 2015; and Zaffagnini et al., 2011 which were previously cited in this policy are included in the Veronesi et al., 2021 systematic review.

In a follow up study to evaluate the long-term clinical results, reoperations, surgical failure and complications at a minimum of 20 years of follow-up, Lucidi et al. (2021) assessed the outcomes of the first 8 patients who received medial CMIs implanted in a pilot prospective study. The pilot study included 8 men with a final age of 55.2 ±8.9 years; however, one patient was not available for the long-term follow-up so data for the remaining 7 men were included in this study. The average follow-up was 21.5 ±0.5 years with evaluations utilizing the Cincinnati Knee Rating System, the visual analogical scale (VAS) and the Lysholm score. The authors did not perform any statistical analyses due to the small number of patients included in the study. The authors reported one failure with one patient who underwent a total knee arthroplasty after 13 years. Of the six patients who were not considered procedure failures, 3 were rated as "excellent" based on the Lysholm score, 1 as "good" and 2 as "fair" while the Cincinnati score and the VAS for pain were stable with respect to the previous follow-ups. The authors found that the CMI for partial meniscal resection could provide pain relief and good knee function at a minimum of 20-year follow-up. The authors concluded that CMI is a safe procedure with satisfactory results and a low failure rate at long-term follow-up.

Grassi et al. (2021) assessed the clinical outcomes and failures of lateral CMI implantation at a minimum 10-year follow-up. This study included 24 consecutive patients who underwent lateral CMI implantation for partial lateral meniscal defects and who were part of a previous study with a 2-year follow-up (cited as Zaffagnini et al., 2015). Outcome measures at the latest follow up included the Lysholm score, Knee injury and Osteoarthritis Outcome Score, visual analog scale (VAS) for pain, Tegner activity level, and EuroQol 5-Dimensions score. Data regarding complications and failures were collected, and patients were asked about their satisfaction with the procedure. Included in the final analysis were 19 patients (16 male, 3 female) with a mean age at surgery of 37.1 ±12.6 years and a mean follow-up of 12.4 ±1.5 years (range, 10-14 years). Five failures (26%) were reported: 1 CMI removal because of implant breakage and 4 joint replacements (2 unicompartmental knee arthroplasties and 2 total knee arthroplasties). The implant survival rate was 96% at 2 years, 85% at 5 years, 85% at 10 years, 77% at 12 years, and 64% at 14 years. Lysholm scores at the final follow-up were rated as "excellent" in 36% (5 of 14 nonfailures), "good" in 43% (6 of 14), and "fair" in 21% (3 of 14). The VAS score was 3.1 ±3.1, with only 16% (3 of 19 patients) reporting that they were pain-free; the median Tegner score was 3 (interquartile range, 2-5). All clinical scores decreased from the 2-year follow-up; however, with the exception of the Tegner score, they remained significantly higher compared with the preoperative status. Overall, 79% of patients were willing to undergo the same procedure. The authors concluded that lateral CMI implantation for partial lateral meniscal defects provided good long-term results, with a 10-year survival rate of 85% and a 14-year survival rate of 64%. At the

final follow-up, 58% of the patients had "good" or "excellent" Lysholm scores. However, there was a general decrease in outcome scores between the short- and the long-term follow-up. According to the authors, although this represents the first study to assess the long-term outcome of lateral meniscal replacement using a scaffold, several limitations are present. Four patients were lost to follow-up, thus creating a possible selection bias. Another limitation is the limited number of patients, which did not allow the performance of sophisticated statistical subanalyses to identify outcomes and failure predictors. According to the authors, additional factors such as a surgical learning curve, the time from meniscectomy to scaffold implantation and the cartilage status, and the time of the index surgery could be relevant and should be investigated in studies with a larger sample size.

Grassi et al. (2014) performed a systematic review to summarize and evaluate the clinical outcomes of the collagen meniscus implant (CMI) and its complication and failure rates. These data were then used to evaluate the results of the CMI at different follow-up time periods and investigate possible differences in the behavior of lateral and medial CMI. All studies evaluating medial or lateral CMI using the Lysholm score, visual analogue scale (VAS) for pain, Tegner activity scale and subjective or objective International Knee Documentation Committee (IKDC) scores were included in the systematic review. Eleven studies were included in the systematic review. The pooled number of patients involved in CMI surgery were 396. The Lysholm score and VAS for pain showed an improvement at six months up to ten years. No noticeable differences were present comparing short-term values of Lysholm score between medial and lateral CMI. The Tegner activity level reached its peak at 12 months after surgery and showed a progressive decrease through five and ten years post CMI implantation, however always remaining above the pre-operative level. Only a few knees were rated as "nearly abnormal" or "abnormal" at IKDC grading at all follow-up evaluations. The reviewers concluded the CMI could produce good and stable clinical results, particularly regarding knee function and pain, with low rates of complications and reoperations.

Harston et al. (2012) conducted a systematic review to examine collagen meniscus implant (CMI) effectiveness for improving patient function, symptoms, and activity level. Study methodologies, rehabilitation, and return to sports guidelines were also reviewed. A total of 11 studies with 520 subjects met inclusion criteria. The authors concluded that knee function, symptoms, and activity level generally improved following CMI use, but poor research report quality was common. They stated that additional well-designed long-term prospective studies are needed to better determine knee osteoarthritis prevention efficacy and appropriate patient selection.

An assessment by the California Technology Assessment Forum (CTAF), (Tice, 2010) concluded that the collagen meniscus implant does not meet CTAF criteria. The CTAF assessment found that the pivotal randomized clinical trial (citing Rodkey et al, 2008) failed to demonstrate any improvement in pain or symptoms in either arm of the trial and the trial has substantial risk for selection bias, confounding, and reporting bias because of the large number of patients lost to follow-up after randomization and the lack of blinding for subjective outcomes. In addition, no data on osteoarthritis were presented. The CTAF assessment concluded that the trial "presents evidence that the collagen meniscus implant offers no important clinical benefits, requires longer and more intensive post-operative rehabilitation, and some uncertainty remains about the potential for long-term harm from the device."

Meniscus Allograft Transplantation (MAT)

In a systematic review, Waugh et al. (2019) assessed the clinical effectiveness of MAT after meniscal injury and subsequent meniscectomy. Thirty-seven papers from 19 studies of MAT were included in the review. Cohort size in the included studies ranged from 30 to 313, with a total of 1731 people undergoing at least one MAT. There was considerable evidence from observational studies, of improvement in symptoms after meniscal allograft transplantation, but the authors found only one small pilot trial with a randomized comparison with a control group that received non-surgical care. MAT has not yet been proven to be chondroprotective. The authors concluded that the benefits of MAT include symptomatic relief and restoration of at least some previous activities, which will be reflected in utility values and hence in quality-adjusted life years, and in the longer term, prevention or delay of osteoarthritis, and avoidance or postponement of some knee replacements.

Elattar et al. (2011) conducted a meta-analysis of published trials reporting outcomes of meniscal allograft transplantation to establish its safety and reproducibility. The outcomes of 678 medial and 458 lateral grafts in 613 male, 265 female and 190 non-defined patients with a mean age of 34.8 years were included in the meta-analysis. According to the authors, all studies reported a continuously satisfactory outcome with restoration of working capacity in these active patients. The authors stated that meniscal allograft transplantation can be considered as safe and reliable for the treatment of refractory post-meniscectomy symptoms in selected patients.

Hergan et al. (2011) performed a systematic review evaluating MAT. Included in the review were 14 studies with at least 2 years' follow-up, studies with validated outcome measures, and studies in which the allograft meniscal horns were secured with bony fixation. Thirteen of the articles provided Level IV evidence, and one article provided Level III evidence. The authors concluded that good early and midterm results of cryopreserved or fresh-frozen, nonirradiated MAT can be achieved in a relatively young patient with only mild chondromalacia (lower than Outerbridge grade 3) who is not overweight and has a stable, mechanically aligned lower extremity, if the allograft is sized radiographically by use of anteroposterior and lateral films and the allograft meniscal horns have bony attachments and are fixed by bony techniques. Similar results can be expected if the transplant is performed alone or with a concomitant cartilage repair procedure; however, significant cartilage defects (Outerbridge grade 2 or greater) on both the femoral and tibial sides in the same compartment requiring autologous cartilage implantation result in a high failure rate. Good outcomes of MAT can be expected when performing a concomitant ligament reconstruction or malalignment procedure on the knee, unless greater than 3 concomitant procedures are performed. There is no significant difference in outcome between medial and lateral MAT. According to the authors, despite a growing body of knowledge on the topic, there remains a lack of consensus regarding optimal allograft sizing technique, allograft fixation techniques, tissue processing, indications, and long-term efficacy. The authors stated that a prospective, randomized trial comparing MAT in a meniscectomized knee with a control group is needed to determine the best technique and patient selection criteria.

Clinical Practice Guidelines

The American Academy of Orthopaedic Surgeons (AAOS)

The AAOS published an information statement regarding the use of musculoskeletal tissue allografts. (AAOS, 2011)

The AAOS supports the following:

- The use of musculoskeletal allograft as a therapeutic alternative to autograft use for appropriate patients. Allograft tissues should be acquired from facilities that demonstrate compliance, use well-accepted banking methodology and good tissue practices. The AAOS urges all tissue banks to follow rigorous national guidelines and standards.
- The AAOS strongly favors on-site inspection and recommends the use of tissue banks by the American Association of Tissue Banks (AATB). The AAOS supports informed consent, for both the donor family and the recipient of human tissue, in accordance with local, state and federal laws and regulations.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Transplantation of meniscal allografts is a surgical procedure and, as such, is not subject to regulation by the FDA. However, the FDA does regulate certain aspects of tissue banking, and tissues are subject to FDA registration and requirements for good tissue practices and infectious disease screening and testing, as well as to the good manufacturing practice requirements applicable to drugs and devices. According to current rules, FDA premarket review or marketing approval is not required for minimally processed tissues transplanted from one person to another for their normal structural functions; these criteria apply to meniscal allografts. Refer to the following website for more information:

<http://www.fda.gov/BiologicsBloodVaccines/TissueTissueProducts/default.htm>. (Accessed May 16, 2022)

Collagen meniscus implants, also known as collagen scaffold, are bioresorbable, primarily bovine type 1 collagen products that are designed as a tissue-engineered scaffold to support the generation of new meniscus-like tissue. For information on collagen meniscus implants, refer to the following FDA website for Premarket Approvals (use product code OLC):

<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed May 16, 2022)

References

The foregoing Oxford policy has been adapted from an existing UnitedHealthcare national policy that was researched, developed and approved by UnitedHealthcare Medical Technology Assessment Committee. [2022T05430]

American Academy of Orthopaedic Surgeons (AAOS). OrthInfo: Meniscal transplant surgery. 2021. Available at: <https://orthoinfo.aaos.org/en/treatment/meniscal-transplant-surgery/>. Accessed May 16, 2022.

American Academy of Orthopaedic Surgeons (AAOS). Use of musculoskeletal tissue allografts. 1991. Revised 2011.

Bulgheroni E, Grassi A, Bulgheroni P, et al. Long-term outcomes of medial CMI implant versus partial medial meniscectomy in patients with concomitant ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2015;23(11):3221-3227.

ECRI. Autologous osteochondral autologous and allogeneic grafts for repairing knee cartilage. Plymouth Meeting (PA): ECRI; 2020 Jul 30. (Clinical Evidence Assessment).

Elattar M, Dhollander A, Verdonk R, et al. Twenty-six years of meniscal allograft transplantation: is it still experimental? A meta-analysis of 44 trials. *Knee Surg Sports Traumatol Arthrosc.* 2011 Feb;19(2):147-57.

Grassi A, Lucidi GA, Filardo G, et al. Minimum 10-Year clinical outcome of lateral collagen meniscal implants for the replacement of partial lateral meniscal defects: further results from a prospective multicenter study. *Orthop J Sports Med.* 2021 May 25;9(5):2325967121994919.

Grassi A, Zaffagnini S, Marcheggiani, et al. Clinical outcomes and complications of a collagen meniscus implant: a systematic review. *Int Orthop.* 2014 Sep;38(9):1945-53.

Harston A, Nyland J, Brand E, et al. Collagen meniscus implantation: A systematic review including rehabilitation and return to sports activity. *Knee Surg Sports Traumatol Arthrosc.* 2012;20(1):135-146.

Hergan D, Thut D, Sherman O, et al. Meniscal allograft transplantation. *Arthroscopy.* 2011 Jan;27(1):101-12.

Lucidi GA, Grassi A, Al-Zu'bi BBH, et al. Satisfactory clinical results and low failure rate of medial collagen meniscus implant (CMI) at a minimum 20 years of follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2021 Dec;29(12):4270-4277.

Rosso F, Bisicchia S, Bonasia DE, et al. Meniscal allograft transplantation: a systematic review. *Am J Sports Med.* Apr 2015.

Tice J. Collagen meniscus implant for repair of medial meniscus injury of the knee. *California Technology Assessment Forum.* June, 2010.

Veronesi F, Di Matteo B, Vitale ND, et al. Biosynthetic scaffolds for partial meniscal loss: A systematic review from animal models to clinical practice. *Bioact Mater.* 2021 Apr 7;6(11):3782-3800.

Warth RJ, Rodkey WG. Resorbable collagen scaffolds for the treatment of meniscus defects: a systematic review. *Arthroscopy.* 2015 May;31(5):927-41.

Waugh N, Mistry H, Metcalfe A, et al. Meniscal allograft transplantation after meniscectomy: clinical effectiveness and cost-effectiveness. *Knee Surg Sports Traumatol Arthrosc.* 2019 Jun;27(6):1825-1839.

World Health Organization and World Bank. (WHO) World report on disability. 2011.

Zaffagnini S, Grassi A, Marcheggiani Muccioli GM, et al. Two-year clinical results of lateral collagen meniscus implant: A multicenter study. *Arthroscopy.* 2015 Jul;31(7):1269-78.

Zaffagnini S, Marcheggiani Muccioli GM, et al. Prospective long-term outcomes of the medial collagen meniscus implant versus partial medial meniscectomy: a minimum 10-year follow-up study. *Am J Sports Med.* 2011 May;39(5):977-85.

Policy History/Revision Information

Date	Summary of Changes
01/01/2023	Documentation Requirements <ul style="list-style-type: none">Added list of applicable CPT codes Supporting Information <ul style="list-style-type: none">Removed <i>Prior Authorization Requirements</i> sectionArchived previous policy version SURGERY 096.18 T2

Instructions for Use

This Clinical Policy provides assistance in interpreting UnitedHealthcare Oxford standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare Oxford reserves the right to modify its Policies as necessary. This Clinical Policy is provided for informational purposes. It does not constitute medical advice.

The term Oxford includes Oxford Health Plans, LLC and all of its subsidiaries as appropriate for these policies. Unless otherwise stated, Oxford policies do not apply to Medicare Advantage members.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. UnitedHealthcare Oxford Clinical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.