

BLEPHAROPLASTY, BLEPHAROPTOSIS AND BROW PTOSIS REPAIR

Guideline Number: MMG008.P

Effective Date: July 1, 2019

[Instructions for Use](#) ⓘ

Table of Contents	Page
COVERAGE RATIONALE	1
DOCUMENTATION REQUIREMENTS	4
DEFINITIONS	4
APPLICABLE CODES	5
REFERENCES	6
Guideline HISTORY/REVISION INFORMATION	7
INSTRUCTIONS FOR USE	7

Related Medical Management Guideline

- [Cosmetic and Reconstructive Procedures](#)

COVERAGE RATIONALE

Indications for Coverage

Some states require benefit coverage for services that UnitedHealthcare considers Cosmetic Services and Surgery, such as repair of external Congenital Defects in the absence of a Functional Impairment. Refer to the member specific benefit plan document.

Criteria for a Coverage Determination that Surgery is Reconstructive and Medically Necessary

The following must be available when requested by UnitedHealthcare:

- Best corrected visual acuity in both eyes, all members (except pediatrics)
- Eye exam (chief complaint, HPI)
- Clear, high-quality, clinical photographs (eye level, frontal with the member looking straight ahead, light reflex visible and centered)
- Peripheral or superior Visual Fields automated, reliable, un-taped/taped are preferable.
 - In situations where computerized Reliable Visual Field testing is not available, we will accept manual Reliable Visual Field testing.
 - In situations where Reliable Visual Field testing is not possible, see section below: "[When the Member is Not Capable of Reliable Visual Field Testing.](#)"

Note: The Visual Fields and high-quality, clinical photographs must be consistent. If multiple procedures are requested, the following criteria must be met:

- All criteria for each individual procedure must be met; and
- Reliable Visual Field testing shows visual impairment which can't be addressed by one procedure alone; and
- High-quality, clinical photograph findings are consistent with Visual Field findings.

Upper eyelid blepharoplasty (CPT 15822 and 15823) is considered reconstructive and medically necessary when the following criteria are present:

- Ptosis has been ruled out as the primary cause of Visual Field obstruction; and
- Clear, high-quality, clinical photographs must show that the extra skin is the primary cause of Visual Field obstruction; and
- The member must have a Functional or Physical Impairment complaint directly related to an abnormality of the eyelid(s); and
- Excess skin (dermatochalasis/blepharochalasis) touches the lashes; and
- Automated peripheral or superior Reliable Visual Field testing, with the eyelid skin taped and un-taped, showing improvement of 30% (or 12 degrees) or more
 - In situations where computerized Reliable Visual Field testing is not available we will accept manual Reliable Visual Field testing.
 - In situations where Reliable Visual Field testing is not possible, see section below, "[When the Member is Not Capable of Reliable Visual Field Testing.](#)"

Note: Extended blepharoplasty may be indicated for blepharospasm (eyelids are forced shut) when the following two criteria are met:

- Debilitating symptoms (e.g., pain); and
- Conservative treatment has been tried and failed, or is contraindicated (e.g., Botox®).

Upper eyelid blepharoptosis repair (CPT 67901– 67909) is considered reconstructive and medically necessary when the following criteria are present:

- The member must have a Functional or Physical Impairment complaint directly related to the position of the eyelid(s); and
- Other treatable causes of ptosis are ruled out (e.g., recent Botox® injections, myasthenia gravis when applicable); and
- Eyelid droop (upper eyelid ptosis) and a Marginal Reflex Distance – 1 (MRD-1) of 2.0 mm or less; and
- The MRD is documented in clear, high-quality, clinical photographs with the member looking straight ahead and light reflex centered on the pupil; and
- Automated peripheral or superior Reliable Visual Field testing, with the eyelids taped and un-taped, showing improvement of 30% (or 12 degrees) or more improvement in the number of points seen.
 - In situations where computerized Reliable Visual Field testing is not available, we will accept manual Reliable Visual Field testing.
 - In situations where Reliable Visual Field testing is not possible, see section below, "[When the Member is Not Capable of Reliable Visual Field Testing.](#)"

Note: For children under age 10 years, ptosis repair is covered to prevent amblyopia. Reliable Visual Field testing is not required, but high-quality, clinical photographs are required.

Brow ptosis (CPT 67900) is considered reconstructive and medically necessary when the following criteria are present:

- Other causes have been eliminated as the primary cause for the Visual Field obstruction (e.g., Botox® treatments within the past six (6) months); and
- The member must have a functional complaint related to brow ptosis. Brow ptosis must be documented in two high-quality, clinical photographs. One showing the eyebrow below the bony superior orbital rim, and a second photograph with the brow elevated that eliminates the Visual Field defect; and
 - Automated peripheral and superior Reliable Visual Field testing, with differential taping (eyebrow and eyebrow + eyelid) showing 30% (or 12 degrees) or more improvement in total number of points seen with the eyebrow taped up. In situations where computerized Reliable Visual Field testing is not available we will accept manual Reliable Visual Field testing.
 - In situations where Reliable Visual Field testing is not possible, see section below, "[When the Member is Not Capable of Reliable Visual Field Testing.](#)"
- Documentation indicating the specific brow lift procedure (e.g., supra-ciliary, midforehead or coronal, pretrichial, direct brow lift vs browpexy, internal brow lift).

Brow ptosis repair (CPT 67900) as an adjunct to upper eyelid blepharoplasty (CPT 15822 and 15823) is considered reconstructive and medically necessary when the criteria for each separate service are met (as per above) AND:

- Automated peripheral and superior Reliable Visual Field testing demonstrates the following:
 - Differential taping showing 30% (or 12 degrees) or more improvement in total number of points seen with the eyelid taped up and an additional 30% (or 12 degrees) or more improvement in total number of points seen with the eyelid + eyebrow taped up, confirming the contribution of brow ptosis to visual field obstruction.

Note: for Browpexy/internal brow lift, see [Coverage Limitations and Exclusions](#).

Eyelid surgery with an anophthalmic socket (has no eyeball) is considered reconstructive and medically necessary when both of the following criteria are present:

- The member has an anophthalmic condition; and
- The member is experiencing difficulties fitting or wearing an ocular prosthesis.

Lower eyelid blepharoplasty (CPT 15820 and 15821) is usually cosmetic, however, is considered reconstructive and medically necessary only when all of the following criteria are present:

- There is documented facial nerve damage; and
- Clear, high-quality, clinical photographs document the pathology; and
- The member is unable to close the eye due to the lower lid dysfunction; and
- Functional Impairment including both of the following:
 - Documented uncontrolled tearing or irritation; and

- Conservative treatments tried and failed.

Ectropion (eyelid turned outward) (CPT 67914 through 67917) or punctal eversion is considered reconstructive and medically necessary when all of the following criteria are present:

- Clear, high-quality, clinical photographs document the pathology.
- Corneal or conjunctival injury with both of the following criteria:
 - Subjective symptoms include either:
 - Pain or discomfort; or
 - Excess tearing; and
 - Any one of the following:
 - Exposure keratitis; and/or
 - Keratoconjunctivitis; and/or
 - Corneal ulcer.

Entropion (eyelid turned inward) (CPT 67921-67924) is considered reconstructive and medically necessary when all of the following criteria are present:

- Clear, high-quality, clinical photographs must document the following:
 - Lid turned inward; and
 - At least one of the following:
 - Trichiasis; or
 - Irritation of cornea or conjunctiva; and
 - Subjective symptoms including either of the following:
 - Excessive tearing; or
 - Pain or discomfort.

Lid Retraction Surgery (CPT 67911) Lid retraction surgery is considered reconstructive and medically necessary when all of the following criteria are present:

- Other causes have been eliminated as the reason for the lid retraction such as use of dilating eye drops, glaucoma medications; and
- Clear, high-quality, clinical photographs document the pathology; and
- There is Functional Impairment (such as 'dry eyes', pain/discomfort, tearing, blurred vision); and
- Tried and failed conservative treatments; and
- In cases of thyroid eye disease two or more Hertel measurements at least 6 months apart with the same base measurements are unchanged.

Canthoplasty/Canthopexy (CPT 21280, 21282, 67950, 67961, 67966) is considered reconstructive and medically necessary when all of the following criteria are present:

- Functional Impairment; and
- Clear, high-quality, clinical photographs document the pathology; and
- Repair of ectropion or entropion will not correct condition; and
- At least one of the following is present:
 - Epiphora (excess tearing) not resolved by conservative measures; or
 - Corneal dryness unresponsive to lubricants; or
 - Corneal ulcer

Repair of Floppy Eyelid Syndrome (FES) (CPT 67961 and 67966) is considered reconstructive and medically necessary when ALL of the following are present when documented and confirmed by history and examination:

- Subjective symptoms must include eyelids spontaneously "flipping over" when the member sleeps due to rubbing on the pillow, **and** one of the following:
 - Eye pain or discomfort; or
 - Excess tearing; or
 - Eye irritation, ocular redness and discharge.
- Physical Examination that documents the following:
 - Eyelash Ptosis; and
 - Significant upper eyelid laxity; and
 - Presence of Giant Papillary Conjunctivitis
 - or**
 - Corneal findings such as:
 - Superficial Punctate Erosions (SPK); or
 - Corneal abrasion (documentation of a history of corneal abrasion or recurrent erosion syndrome is considered sufficient); or
 - Microbial Keratitis.

- Clear, high-quality, clinical photographs that clearly document Floppy Eyelid Syndrome and demonstrate both of the following:
 - Lids must be everted in the photographs; and
 - Conjunctival surface (underbelly) of the lids must clearly demonstrate Giant Papillary Conjunctivitis
- Documentation that conservative treatment has been tried and failed, examples may include:
 - Ocular lubricants both drops (daytime) and ointments (bedtime); or
 - Short trial of antihistamines; or
 - Topical steroid drops; or
 - Eye shield and/or taping the lids at bedtime.
- Other causes of the eye findings have been ruled out, examples may include:
 - Allergic conjunctivitis
 - Atopic keratoconjunctivitis
 - Blepharitis
 - Contact lens (CL) complication
 - Dermatochalasis
 - Ectropion
 - Giant Papillary Conjunctivitis (GPC) that is not related to FES
 - Ptosis of the lid(s)
 - Superior limbic keratoconjunctivitis (SLK)

When the Member Is Not Capable of Reliable Visual Field Testing

Reliable Visual Field testing is not required when the member is not capable of performing a Visual Field test. The following are some examples:

- If the member is a child 12 years old or under
- If the member has intellectual disabilities (previously known as mental retardation) or some other severe neurologic disease

Coverage Limitations and Exclusions

UnitedHealthcare West excludes Cosmetic Services and Surgery from coverage including but not limited to the following:

- Procedures that do not meet the reconstructive criteria above in the [Indications for Coverage](#) section.
- Browpexy/internal brow lift is not designed to improve function.

DOCUMENTATION REQUIREMENTS

Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The documentation requirements outlined below are used to assess whether the member meets the clinical criteria for coverage but do not guarantee coverage of the service requested.

Required Clinical Information

Blepharoplasty, Blepharoptosis and Brow Ptosis Repair

Medical notes documenting **all** of the following:

- Goldman, manual or automated technique visual fields, taped and untaped
- Marginal reflex distance (MRD-1)
- Visual complaints, functional impairments and ruling out other causes
- High-quality photograph(s); all photos must be full face, labeled with the date taken and the applicable case number obtained at time of notification, or member's name and ID number on the photograph(s)

Note: Submission of color photos can be submitted via the external portal at www.uhcprovider.com/paan or via email at CCR@uhc.com; faxes of color photos will not be accepted.

DEFINITIONS

The following definitions may not apply to all plans. Refer to the member specific benefit plan document for applicable definitions.

Congenital Defect: A condition present at birth.

Cosmetic Services and Surgery (CA only): Cosmetic surgery and cosmetic services are not covered. Cosmetic surgery and cosmetic services are defined as surgery and services performed to alter or reshape normal structures of the body in order to improve appearance. Drugs, devices and procedures related to cosmetic surgery or cosmetic services are not covered. Surgeries or services that would ordinarily be classified as cosmetic will not be reclassified as reconstructive, based on a Member's dissatisfaction with his or her appearance.

Cosmetic Services and Surgery (WA, OR, TX and OK only): Cosmetic surgery and cosmetic services are not covered. Cosmetic surgery and cosmetic services are defined as surgery and services performed to alter or reshape normal structures of the body in order to improve appearance. Drugs, devices and procedures related to cosmetic surgery or cosmetic services are not covered. Surgeries or services that would ordinarily be classified as cosmetic will not be reclassified as reconstructive, based on a Member's dissatisfaction with his or her appearance, as influenced by that Member's underlying psychological makeup or psychiatric condition.

Floppy Eyelid Syndrome (FES): Characterized by significant upper eyelid laxity and chronic papillary conjunctivitis in upper palpebral conjunctiva that is poorly responsive to topical lubrication and steroids. FES is known to be associated with obesity, obstructive sleep apnea, Down syndrome, and keratoconus. Keratoconus can be linked to frequent rubbing and mechanical effect on the palpebral conjunctiva and cornea.

Functional or Physical or Physiological Impairment: Functional or Physical or Physiological Impairment causes deviation from the normal function of a tissue or organ. This results in a significantly limited, impaired, or delayed capacity to move, coordinate actions, or perform physical activities and is exhibited by difficulties in one or more of the following areas: physical and motor tasks; independent movement; performing basic life functions.

Giant Papillary Conjunctivitis: Is defined by exam findings of giant papillary hypertrophy primarily affecting the upper tarsal conjunctiva.

Marginal Reflex Distance – 1 (MRD-1): Measures the number of millimeters from the corneal light reflex or center of the pupil to the upper lid margin. (Note: the "-1" in MRD-1 refers to the upper lid and not the measurement in millimeters.)

Marginal Reflex Distance – 2 (MRD-2): Measures the number of millimeters from the corneal light reflex or center of the pupil to the lower lid margin. (Note: the "-2" in MRD-2 refers to the lower lid and not the measurement in millimeters.)

Reconstructive Surgery: Reconstructive surgery is covered to correct or repair abnormal structures of the body caused by congenital defects, developmental abnormalities, trauma, infection, tumors or disease. The purpose of reconstructive surgery is to correct abnormal structures of the body to improve function or create a normal appearance to the extent possible.

Reliable (Visual Fields): The reliability of the visual fields is indicated in the visual field report with the number of fixation losses, false negative and false positives. A reliable visual field has less than 30% or fewer of fixation losses, false negative and false positives.

APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this guideline does not imply that the service described by the code is a covered or non-covered health care service. Benefit coverage for health care services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

Note: The following codes may be cosmetic; review is required to determine if considered cosmetic or reconstructive.

CPT Code	Description
Blepharoplasty (Lower eyelid)	
15820	Blepharoplasty, lower eyelid;
15821	Blepharoplasty, lower eyelid; with extensive herniated fat pad
Blepharoplasty (Upper eyelid)	
15822	Blepharoplasty, upper eyelid;
15823	Blepharoplasty, upper eyelid; with excessive skin weighting down lid
Brow Ptosis Repair	
67900	Repair of brow ptosis (supraciliary, mid-forehead or coronal approach)
Upper Eyelid Blepharoptosis Repair	
67901	Repair of blepharoptosis; frontalis muscle technique with suture or other material (e.g., banked fascia)

CPT Code	Description
Upper Eyelid Blepharoptosis Repair	
67902	Repair of blepharoptosis; frontalis muscle technique with autologous fascial sling (includes obtaining fascia)
67903	Repair of blepharoptosis; (tarso) levator resection or advancement, internal approach
67904	Repair of blepharoptosis; (tarso) levator resection or advancement, external approach
67906	Repair of blepharoptosis; superior rectus technique with fascial sling (includes obtaining fascia)
67908	Repair of blepharoptosis; conjunctivo-tarso-Muller's muscle-levator resection (e.g., Fasanella-Servat type)
67909	Reduction of overcorrection of ptosis
Lid Retraction	
67911	Correction of lid retraction
Ectropion	
67914	Repair of ectropion; suture
67915	Repair of ectropion; thermocauterization
67916	Repair of ectropion; excision tarsal wedge
67917	Repair of ectropion; extensive (e.g., tarsal strip operations)
67921	Repair of entropion; suture
67922	Repair of entropion; thermocauterization
67923	Repair of entropion; excision tarsal wedge
67924	Repair of entropion; extensive (e.g., tarsal strip or capsulopalpebral fascia repairs operation)
Canthus Repair and Lid Repair	
21280	Medial canthopexy (separate procedure)
21282	Lateral canthopexy
67950	Canthoplasty (reconstruction of canthus)
67961	Excision and repair of eyelid, involving lid margin, tarsus, conjunctiva, canthus, or full thickness, may include preparation for skin graft or pedicle flap with adjacent tissue transfer or rearrangement; up to one-fourth of lid margin
67966	Excision and repair of eyelid, involving lid margin, tarsus, conjunctiva, canthus, or full thickness, may include preparation for skin graft or pedicle flap with adjacent tissue transfer or rearrangement; over one-fourth of lid margin
Floppy Eyelid Syndrome	
67961	Excision and repair of eyelid, involving lid margin, tarsus, conjunctiva, canthus, or full thickness, may include preparation for skin graft or pedicle flap with adjacent tissue transfer or rearrangement; up to one-fourth of lid margin
67966	Excision and repair of eyelid, involving lid margin, tarsus, conjunctiva, canthus, or full thickness, may include preparation for skin graft or pedicle flap with adjacent tissue transfer or rearrangement; over one-fourth of lid margin

CPT® is a registered trademark of the American Medical Association

REFERENCES

- American Society of Plastic Surgeons (ASPS); Practice Parameter for Blepharoplasty; Approved by the Executive Committee of the American Society of Plastic Surgeons®, March 2007. Available at: [Practice Parameter For Blepharoplasty – American Society of Plastic Surgeons](#). Accessed May 1, 2019.
- Burkat CN, Lemke BN. Acquired lax eyelid syndrome: an unrecognized cause of the chronically irritated eye. *Ophthalm Plast Reconstr Surg*. 2005 Jan;21(1):52-8.
- Chambe J, Laib S, Hubbard J, Erhardt C, Ruppert E, Schroder C, et al. Floppy eyelid syndrome is associated with obstructive sleep apnea: a prospective study on 127 patients. *J Sleep Res*. 2012 Jun;21(3):308-15.

Fowler AM, Dutton JJ Floppy eyelid syndrome as a subset of lax eyelid conditions: relationships and clinical relevance (an ASOPRS thesis). *Ophthal Plast Reconstr Surg.* 2010;26(3):195-204. doi: 10.1097/IOP.0b013e3181b9e37.

Korn BS, Chokthaweesak W, et al. *Video Atlas of Oculofacial Plastic and Reconstructive Surgery.* 2nd ed. Elsevier Inc. 2016. Chapter 21, Internal Brow Plasty; p.143-146.

MCG™ Care Guidelines, 23rd edition, 2019. Blepharoplasty, Canthoplasty, and Related Procedures ACG: A-0195 (AC). Orbit, Eyelids and Lacrimal System, Section 7. Basic and Clinical Science Course. San Francisco: American Academy of Ophthalmology; 2009.

Periman LM, Sires BS. Floppy eyelid syndrome: a modified surgical technique. *Ophthal Plast Reconstr Surg.* 2002 Sep;(18)5:370-2.

Trussler AP, Rohrich RJ. MOC-PSSM CME article: Blepharoplasty. *Plast Reconstr Surg.* 2008 Jan; 121 (1 Suppl): 1-10.

Valenzuela AA, Sullivan TJ. Medial upper eyelid shortening to correct medial eyelid laxity in floppy eyelid syndrome: a new surgical approach. *Ophthal Plast Reconstr Surg.* 2005;21(4):259-63.

Warren RJ, Neligan PC. *Plastic Surgery: Volume 2.* 3rd edition. Saunders. 2012. Chapter 7, Forehead Rejuvenation; p. 93-107.

GUIDELINE HISTORY/REVISION INFORMATION

Date	Action/Description
08/01/2019	Template Update <ul style="list-style-type: none"> • Added <i>Documentation Requirements</i> section
07/01/2019	Coverage Rationale <ul style="list-style-type: none"> • Simplified content (no change to guidelines) Definitions <ul style="list-style-type: none"> • Added definition of "Reconstructive Surgery" • Updated definition of "Congenital Defect" • Removed definition of "Reconstructive Procedures (CA only)" Supporting Information <ul style="list-style-type: none"> • Archived previous policy version MMG008.O

INSTRUCTIONS FOR USE

This Medical Management Guideline provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard benefit plan. In the event of a conflict, the member specific benefit plan document governs. Before using this guideline, please the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Management Guideline is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. UnitedHealthcare West Medical Management Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Member benefit coverage and limitations may vary based on the member's benefit plan Health Plan coverage provided by or through UnitedHealthcare of California, UnitedHealthcare Benefits Plan of California, UnitedHealthcare of Oklahoma, Inc., UnitedHealthcare of Oregon, Inc., UnitedHealthcare Benefits of Texas, Inc., or UnitedHealthcare of Washington, Inc.