

UnitedHealthcare Pharmacy
Clinical Pharmacy Programs

Program Number	2021 P 1172-7
Program	Prior Authorization – California, Maryland and New York Regulatory Program - Weight Loss
Medication	Includes both brand and generic versions and all formulations of the listed products unless otherwise noted Xenical (orlistat), benzphetamine, diethylpropion, phendimetrazine, phentermine (all brand products including Adipex-P and Lomaira) Contrave (naltrexone and bupropion), Imcivree (setmelanotide), Qsymia (phentermine and topiramate extended-release), and Saxenda (liraglutide)
P&T Approval Date	11/2015, 9/2016, 9/2018, 9/2019, 11/2019, 11/2020, 4/2021
Effective Date	7/1/2021; Oxford only: 7/1/2021

1. **Background:**

Anti-obesity agents are indicated in the management of obesity as an adjunct to lifestyle modifications including diet, exercise and behavioral modification. Medication therapy may provide modest weight reduction in conjunction with lifestyle modifications and therapy selection may be based on a specific medications side effects and warnings.

Body Mass Index (BMI) uses weight and height to create an index of underweight, overweight or obesity in adults. The international classification is as follows:

Classification	BMI(kg/m²)
Underweight	< 18.50
Normal range	18.50 - 24.99
Overweight	≥ 25.00
Obese	≥ 30.00
Obese class I	30.00 - 34.99
Obese class II	35.00 - 39.99
Obese class III	≥ 40.00

WHO Global Database on Body Mass Index

This program uses Obese Class III and Obese Class I (with weight related comorbidities) as markers for coverage and is designed to meet regulatory requirements for coverage of weight loss medications in Maryland and New York and morbid obesity in California.

2. **Coverage Criteria:**

A. benzphetamine, diethylpropion, phendimetrazine, phentermine (Includes both brand and generic versions and all formulations of the listed products unless otherwise noted):

1. **Initial Authorization:**

a. Coverage will be approved based on **all** of the following:

- (1) Treatment is being requested for weight loss
- (2) Patient is > 16 years of age
- (3) Failure to lose $\geq 5\%$ of body weight after at least 6 months of lifestyle modification alone (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program). Document lifestyle modifications employed and total weight loss
- (4) **One** of the following:
 - (a) Failure to lose $\geq 5\%$ of body weight after six months of treatment with OTC orlistat (Alli) (document date of trial of orlistat and total body weight lost)
 - (b) Contraindication (including age) or intolerance to OTC orlistat (Alli)
- (5) Used as an adjunct to lifestyle modification. Document which lifestyle modification will be employed.
- (6) **One** of the following:
 - (a) Body Mass Index (BMI) ≥ 40 kg/m² (Obesity Class III). Documentation of current height and weight required.

-OR-

 - (b) **Both** of the following:
 - i. BMI ≥ 30 kg/m² (Obesity Class I). Documentation of current height and weight required.
 - ii. Documentation of a weight-related comorbidity (examples include dyslipidemia, hypertension, type 2 diabetes, sleep apnea)

Authorization will be issued for 3 months.

2. **Reauthorization:**

a. Coverage will be approved based on **both** of the following:

- (1) Documentation of current weight showing a weight loss of $\geq 5\%$ of baseline body weight
- (2) Documentation of continuation of lifestyle modification

Authorization for will be issued for 6 months.

B. Xenical, Contrave, or Qsymia (Includes both brand and generic versions and all formulations of the listed products unless otherwise noted):

1. **Initial Authorization:**

a. Coverage will be approved based on **all** of the following:

- (1) Treatment is being requested for weight loss
- (2) **One** of the following:
 - (a) Patient is ≥ 12 years of age for Xenical
 - (b) Patient is ≥ 18 years of age for Contrave, Qsymia
- (3) Failure to lose $\geq 5\%$ of body weight through at least 6 months of lifestyle modification alone (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program). Document lifestyle modifications employed and total weight loss
- (4) **One** of the following:
 - (a) Failure to lose $\geq 5\%$ of body weight after six months of treatment with OTC orlistat (Alli) (document date of trial of orlistat and total body weight lost)
 - (b) Contraindication (including age) or intolerance to OTC orlistat (Alli)
- (5) Used as an adjunct to lifestyle modification. Document which lifestyle modification will be employed.
- (6) **One** of the following:
 - (a) Body Mass Index (BMI) ≥ 40 kg/m² (Obesity Class III) Documentation of current height and weight required.

-OR-

- (b) **Both** of the following:
 - i. BMI ≥ 30 kg/m² (Obesity Class I) Documentation of current height and weight required.
 - ii. Documentation of a weight-related comorbidity (examples include dyslipidemia, hypertension, type 2 diabetes, sleep apnea)

Contrave: Authorization will be issued for 4 months.

Qsymia, Xenical: Authorization will be issued for 6 months.

2. Reauthorization

a. Coverage will be approved based on **both** of the following:

- (1) Documentation of current weight showing a weight loss of $\geq 5\%$ of baseline body weight
- (2) Documentation of continuation of lifestyle modification

Authorization will be issued for 6 months.

C. Saxenda

1. Initial Authorization:

a. Coverage will be approved based on **all** of the following:

- (1) Treatment is being requested for weight loss
- (2) Patient is ≥ 18 years of age
- (3) Failure to lose $\geq 5\%$ of body weight through at least 6 months of lifestyle modification alone (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program). Document lifestyle modifications employed and total weight loss
- (4) **Both** of the following:
 - (a) **One** of the following:
 - i. Failure to lose $\geq 5\%$ of body weight after six months of treatment with OTC orlistat (Alli) (document date of trial of orlistat and total body weight lost)
 - ii. Contraindication (including age) or intolerance to OTC orlistat (Alli)
 - (b) Contraindication, intolerance or failure to lose and maintain $\geq 5\%$ body weight following 3 month trial EACH, of **two** of the following medications (document date of trial of each medication and total body weight lost):
 - i. Prescription Xenical
 - ii. Qsymia
 - iii. Contrave

(5) Used as an adjunct to lifestyle modification. Document which lifestyle modification will be employed

(6) **One** of the following:

(a) Body Mass Index (BMI) ≥ 40 kg/m² (Obesity Class III). Documentation of current height and weight required.

-OR-

(b) **Both** of the following:

i. BMI ≥ 30 kg/m² (Obesity Class I). Documentation of current height and weight required.

ii. Documentation of a weight-related comorbidity (examples include dyslipidemia, hypertension, type 2 diabetes, sleep apnea)

Authorization will be issued for 4 months.

2. Reauthorization

a. Coverage will be approved based on **both** of the following:

(1) Documentation of current weight showing a weight loss of $\geq 5\%$ of baseline body weight

(2) Documentation of continuation of lifestyle modification

Authorization will be issued for 6 months.

D. Imcivree

1. Initial Authorization:

a. Coverage will be approved based on **all** of the following:

(1) Diagnosis of obesity is due to POMC, PCSK1, or LEPR gene deficiency

-AND-

(2) **One** of the following:

(a) Adult patient with BMI ≥ 30 kg/m²

-OR-

(b) Pediatric patient with weight $>95^{\text{th}}$ percentile for age on growth chart assessment

-AND-

- (3) Genetic testing confirming variants in POMC, PCSK1, or LEPR genes interpreted as pathogenic, likely pathogenic, or of uncertain significance

-AND-

- (4) Patient is currently enrolled in or has history of a weight loss management program

Authorization will be issued for 6 months.

2. Reauthorization

- a. Coverage will be approved based on **one** the following criterion:

- (1) If on therapy for less than 12 months, documentation of a positive clinical response to Imcivree therapy defined as weight loss $\geq 5\%$ of baseline weight

-OR-

- (2) If on therapy for ≥ 12 months, documentation of a positive clinical response to Imcivree therapy defined as $\geq 10\%$ weight loss from baseline

Authorization will be issued for 12 months.

3. Additional Clinical Rules:

- Supply limits may be in place.

4. References:

1. AACE position statement on obesity and obesity medicine. September/October 2012.
2. National Institutes of Health, National Heart, Lung, and Blood Institute, and North American Association for the Study of Obesity. The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. 2000.
3. American Gastroenterological Association medical position statement on obesity. Gastroenterology 2002 Sep;123(3):879-81.
4. Bray, GA, Ryan, DH. Medical Therapy for the Patient with Obesity. Circulation. 2012;125:1695-1703.
5. Barlow, SE. Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report. Pediatrics 2007.
6. Benzphetamine [package insert]. HJ Harkins Company, Inc. Heritage Pharmaceuticals. December 2019,
7. Diethylpropion [package insert]. Philadelphia, PA: Lannett Company, Inc. December 2019.

8. Phendimetrazine Slow-Release Capsules [package insert]. Princeton, NJ: Sandoz Inc. ; November 2018
9. Adipex-P® [package insert]. Parsippany, NJ : Teva Pharmaceuticals USA, Inc. ; s ;March 2020.
10. Xenical [package insert]. South San Francisco, CA : Roche Pharmaceuticals ; September 2020
10. Qsymia [package insert]. Moutain View, CA. Vivus, Inc ; March 2018. 13.
Contrace [package insert]/ Deerfield, IL : Takeda Pharmaeuticals Inc ; September, 2014.
11. Saxenda [package insert]. Plainsboro, NJ : Novo Nordisk. March 2020.
12. Bray, GA. Obesity in Adults. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. (Accessed on September 14, 2015.)
13. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). JAMA. 2001;285:2486-2497.
14. Pharmacological Management of Obesity: An Endocrine Society Clinical Practice Guideline. The Journal of Clinical Endocrinology & Metabolism 2015 100:2, 342-362
15. AHA/ACC/TOS Prevention Guideline: 2013 AHA/ACC/TOS Guideline for the Management of Overweight and Obesity in Adults: A Report to the American College of Cardiology/American Health Association Task Force on Practice Guidelines and The Obesity Society. Circulation. 2014; 129:S102-138
16. World Health Organization. (2006). *Global Database on Body Mass Index*. Retrieved from http://apps.who.int/bmi/index.jsp?introPage=intro_3.html
17. Lomaira [package insert]. Newtown, PA : KVK-Tech, Inc. December 2018. .
18. Imcivree [package insert]. Boston, MA : Rhythm Pharmaceuticals, Inc ; November 2020.

Program	Prior Authorization - California and New York Regulatory Program - Weight Loss
Change Control	
Date	Change
11/2015	New program.
9/2016	Annual review. Added Belviq XR.
9/2018	Added Lomaira as in scope. Updated references. Formatting changes.
9/2019	Annual Review. Revised background section. Revised list of modifications of weight related comorbidity.
11/2019	Added Maryland as in scope per rider requirements.
11/2020	Annual review. Updated references. Removed Belviq/Belviq XR.
4/2021	Added Imcivree as in scope. Added Imcivree criteria. Updated references. Formatting changes.