

Peer Specialty: Endocrinology					
Quality of Care					
UnitedHealth Premium® methodology documents are located on unitedhealthpremium.uhc.com .					
Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Chronic Kidney Disease	Patient(s) compliant with prescribed SGLT2 inhibitor (minimum compliance 80%)	Patient was 80% or more compliant with prescribed SGLT2 inhibitor medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
Concurrent Use of Opioids and Benzodiazepines	Patient(s) did not have concurrent use of prescription opioids and benzodiazepines	Patient did not have prescription opioid and benzodiazepine medications concurrently dispensed	Safety	Prescribing	Contact Partnership for Quality Measurement
Coronary Artery Disease	Patient(s) compliant with prescribed beta-blocker-containing medication (minimum compliance 80%)	Patient was 80% or more compliant with prescribed beta-blocker-containing medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
Diabetes	Patient(s) compliant with prescribed ACE-inhibitor-containing medication (minimum compliance 80%)	Patient was 80% or more compliant with prescribed ACE-inhibitor-containing medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
	Patient(s) compliant with prescribed angiotensin receptor blocker-containing medication (minimum compliance 80%)	Patient was 80% or more compliant with prescribed angiotensin receptor blocker-containing medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
	Patient(s) compliant with prescribed sodium-glucose cotransporter 2 (SGLT2) inhibitor-containing medication (minimum compliance 80%)	Patient was 80% or more compliant with prescribed sodium-glucose cotransporter 2 (SGLT2) inhibitor-containing medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
	Patient(s) that had at least 2 HbA1c tests in last 12 reported months	Patient had at least 2 HbA1c tests in last 12 reported months	Guideline Concordance: Chronic Disease	Patient	Synopsis
	Patient(s) with a diagnosis of diabetic nephropathy, proteinuria, or chronic renal failure currently taking an ACE-inhibitor or angiotensin receptor blocker	Patient with diabetic nephropathy, proteinuria, or chronic renal failure had an ACE-inhibitor or angiotensin receptor antagonist medication dispensed	Guideline Concordance: Chronic Disease	Patient	Synopsis
Diabetes Medications-Part D Medication Adherence	Patient(s) compliant with all prescribed diabetes medications (minimum compliance 80% or higher) (Medicare only)	Patient was 80% or more compliant with all prescribed diabetes medications	Guideline Concordance: Chronic Disease	Patient	Contact Partnership for Quality Measurement
Head Imaging for Uncomplicated Headache	Patient(s) with uncomplicated headache that did not have imaging studies	Patient with uncomplicated headache did not have imaging studies	Low Value Care	Rendering	Synopsis

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Hyperlipidemia	Patient(s) compliant with prescribed bempedoic acid-containing medication (minimum compliance 80%)	Patient was 80% or more compliant with prescribed bempedoic acid-containing medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
Kidney Health Evaluation for Patients with Diabetes	Patient(s) 18-64 years with diabetes that had kidney health evaluation in last 12 reported months	Patient with diabetes had kidney evaluation	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
	Patient(s) 65-74 years with diabetes that had kidney health evaluation in last 12 reported months	Patient with diabetes had kidney evaluation	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
	Patient(s) 75-85 years with diabetes that had kidney health evaluation in last 12 reported months	Patient with diabetes had kidney evaluation	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
Medication Safety Monitoring	Older adult patients who had an accidental fall or hip fracture who did not use an antiepileptic, nonbenzodiazepine hypnotic, SSRI, SNRI, antipsychotic, benzodiazepine, or tricyclic antidepressant after the incident	Patient with an accidental fall or hip fracture did not have an antiepileptic, nonbenzodiazepine hypnotic, SSRI, SNRI, antipsychotic, benzodiazepine, or tricyclic antidepressant medication dispensed after the incident	Safety	Prescribing	Contact Partnership for Quality Measurement
	Older adult patients with chronic kidney disease who did not use a Cox-2 selective or nonaspirin NSAID after the earliest record of chronic kidney disease	Patient with chronic kidney disease did not have a Cox-2 selective or nonaspirin non-steroidal anti-inflammatory drug (NSAID) dispensed after the earliest record of chronic kidney disease	Safety	Prescribing	Contact Partnership for Quality Measurement
	Older adult patients with dementia who did not use an antipsychotic, benzodiazepine, tricyclic antidepressant, nonbenzodiazepine hypnotic or anticholinergic agent after the earliest record of dementia	Patient with dementia did not have an antipsychotic, benzodiazepine, tricyclic antidepressant, nonbenzodiazepine hypnotic or anticholinergic agent dispensed after the earliest record of dementia	Safety	Prescribing	Contact Partnership for Quality Measurement
Osteoporosis Management	Patient(s) compliant with prescribed oral bisphosphonate (minimum compliance 80%)	Patient was 80% or more compliant with prescribed oral bisphosphonate medication	Guideline Concordance: Chronic Disease	Patient	Synopsis
	Women 67-85 years who were treated or tested for osteoporosis within six months of a fracture	Patient was treated or tested for osteoporosis within six months of a fracture	Guideline Concordance: Chronic Disease	Patient	Contact Partnership for Quality Measurement
Renin Angiotensin System (RAS) Antagonists-Part D Medication Adherence	Patient(s) compliant with prescribed RAS antagonist medication (minimum compliance 80% or higher) (Medicare only)	Patient was 80% or more compliant with prescribed RAS antagonist medication	Guideline Concordance: Chronic Disease	Patient	Contact Partnership for Quality Measurement

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Risk of Continued Opioid Use	Patient(s) age 18-64 years who were opioid-naive and were not prescribed access to opioid medication for 15 or more days during the first 30 days following first opioid treatment initiation	Patient did not have opioid medication for 15 or more days during the first 30 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
	Patient(s) age 65 years and older who were opioid-naive and were not prescribed access to opioid medication for 15 or more days during the first 30 days following first opioid treatment initiation	Patient did not have opioid medication for 15 or more days during the first 30 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
	Patient(s) age 18-64 years who were opioid-naive and were not prescribed access to opioid medication for 31 or more days during the first 62 days following first opioid treatment initiation	Patient did not have opioid medication for 31 or more days during the first 62 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
	Patient(s) age 65 years and older who were opioid-naive and were not prescribed access to opioid medication for 31 or more days during the first 62 days following first opioid treatment initiation	Patient did not have opioid medication for 31 or more days during the first 62 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
Statin Therapy for Patients with Cardiovascular Disease	Men 21-75 years with cardiovascular disease that received a high-intensity or moderate-intensity statin medication	Patient with cardiovascular disease had a high or moderate-intensity statin medication dispensed	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
	Women 40-75 years with cardiovascular disease that received a high-intensity or moderate-intensity statin medication	Patient with cardiovascular disease had a high or moderate-intensity statin medication dispensed	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
	Men 21-75 years with statin adherence (proportion of days covered) at least 80% during the treatment period	Patient was 80% or more compliant with prescribed statin medication	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
	Women 40-75 years with statin adherence (proportion of days covered) at least 80% during the treatment period	Patient was 80% or more compliant with prescribed statin medication	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
Statin Therapy for Patients with Diabetes	Patient(s) 40-75 years with diabetes that received a statin medication	Patient with diabetes had a statin medication dispensed	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance
	Patient(s) with statin adherence (proportion of days covered) at least 80% during the treatment period	Patient was 80% or more compliant with prescribed statin medication	Guideline Concordance: Chronic Disease	Patient	Contact National Committee for Quality Assurance

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Statins-Part D Medication Adherence	Patient(s) compliant with prescribed statin medication (minimum compliance 80% or higher) (Medicare only)	Patient was 80% or more compliant with prescribed statin medication	Guideline Concordance: Chronic Disease	Patient	Contact Partnership for Quality Measurement
Use of Contrast Material in CT	Patient(s) with an abdomen CT test performed that did not have "combined studies" (with and without contrast material)	Patient did not have an abdomen CT test using combined studies (with and without contrast material)	Low Value Care	Ordering	Contact Centers for Medicare & Medicaid Services
Use of High-Risk Medications in Older Adults	Patients 67 years and older who did not receive two or more of the same high-risk medications except for appropriate diagnosis in the last 12 reported months	Patient did not have two or more of the same high-risk medications except for the appropriate diagnosis dispensed	Safety	Prescribing	Contact Partnership for Quality Measurement
	Patients 67 years and older who did not receive two or more of the same high-risk medications from the same drug class in the last 12 reported months	Patient did not have two or more of the same high-risk medications from the same drug class dispensed	Safety	Prescribing	Contact Partnership for Quality Measurement
Use of Opioid Medications	Patient(s) 18 years or older without an average morphine milligram equivalent (MME) \geq 90mg/day during the treatment period	Patient did not have an average morphine equivalent dose \geq 90 mg/day	Safety	Prescribing	Contact National Committee for Quality Assurance
Use of Opioids from Multiple Providers	Patient(s) 18 years or older that did not fill opioid prescriptions from four or more different prescribers	Patient did not have opioid medications from four or more different prescribers dispensed	Safety	Prescribing	Contact National Committee for Quality Assurance

Important notes about UnitedHealth Premium®

The information from UnitedHealth Premium is not an endorsement of a particular physician or health care professional's suitability for the health care needs of any member. UnitedHealthcare does not practice medicine nor provide health care services. Physicians are solely responsible for medical judgments and treatments.

A Premium Care Physician designation does not guarantee the quality or the outcome of any health care services members receive. The fact that a physician does not have a Premium Care Physician designation does not mean the physician does not provide quality health care services.

All physicians in the UnitedHealthcare Network have met certain minimum credentialing requirements. Regardless of whether a physician has received a Premium Care Physician designation, members have access to all physicians in the UnitedHealthcare Network as described in the member's benefit plan.

There are various reasons why a physician may not be designated as a Premium Care Physician. A physician may not receive a designation because that physician has not been evaluated. This occurs when a physician does not practice in a specialty or market that is evaluated by Premium, or the physician's evaluation is in process. This also occurs when there are not enough measures, patients, and/or episodes attributed to the physician for evaluation. This is not an indicator of the total number of patients treated by the physician, or the number of procedures performed by the physician.

UnitedHealthcare informs members that designations are intended only as a guide when choosing a physician and should not be the sole factor in selecting a physician. Members are encouraged to discuss designations with a physician before choosing them or consult with their current physician(s) for advice on selecting other physicians.

As with all programs that evaluate performance based on evaluation of a sample, there is a risk of error. There is a risk of error in the claims data used and in the way patient care is attributed to physicians. UnitedHealth Premium uses statistical testing to compare a physician's performance to benchmarks. There is a risk of error in statistical tests when applied to the data and a result based on statistical testing is not a guarantee of correct inference or classification. Physicians have the opportunity to review the data and evaluation results and may submit requests for changes and/or corrections.

The information contained in this document is subject to change.