

Peer Specialty: Ear, nose, and throat
Quality of Care

UnitedHealth Premium® methodology documents are located on unitedhealthpremium.uhc.com.

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Acute Otitis Externa	Patient(s) 2 years of age and older with acute otitis externa who were not prescribed systemic antimicrobial therapy	Patient with acute otitis externa did not have a systemic antimicrobial medication dispensed	Low Value Care	Rendering	Synopsis
Acute Sinusitis	Patient(s) treated with an antibiotic for acute sinusitis that received a first line antibiotic	The first antibiotic medication patient received for acute sinusitis was a first line antibiotic medication	Guideline Concordance: Acute Care	Rendering	Synopsis
Adenoidectomy	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Appropriate Treatment for Upper Respiratory Infection	Patient(s) 3 months to 17 years of age with a diagnosis of upper respiratory infection (URI) that did not have a prescription for an antibiotic on or within three days after the initiating visit	Patient with a diagnosis of upper respiratory infection (URI) did not have a prescription for an antibiotic on or within three days after the initiating visit	Low Value Care	Rendering	Contact National Committee for Quality Assurance
	Patient(s) 18-64 years of age with a diagnosis of upper respiratory infection (URI) that did not have a prescription for an antibiotic on or within three days after the initiating visit	Patient with a diagnosis of upper respiratory infection (URI) did not have a prescription for an antibiotic on or within three days after the initiating visit	Low Value Care	Rendering	Contact National Committee for Quality Assurance
	Patient(s) 65 years of age and older with a diagnosis of upper respiratory infection (URI) that did not have a prescription for an antibiotic on or within three days after the initiating visit	Patient with a diagnosis of upper respiratory infection (URI) did not have a prescription for an antibiotic on or within three days after the initiating visit	Low Value Care	Rendering	Contact National Committee for Quality Assurance
Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis	Patient(s) 3 months to 17 years of age with a diagnosis of acute bronchitis/bronchiolitis that did not have a prescription for an antibiotic on or within three days after the initiating visit	Patient with a diagnosis of acute bronchitis/bronchiolitis did not have a prescription for an antibiotic on or within three days after the initiating visit	Low Value Care	Rendering	Contact National Committee for Quality Assurance

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis	Patient(s) 18-64 years of age with a diagnosis of acute bronchitis/bronchiolitis that did not have a prescription for an antibiotic on or within three days after the initiating visit	Patient with a diagnosis of acute bronchitis/bronchiolitis did not have a prescription for an antibiotic on or within three days after the initiating visit	Low Value Care	Rendering	Contact National Committee for Quality Assurance
	Patient(s) 65 years of age and older with a diagnosis of acute bronchitis/bronchiolitis that did not have a prescription for an antibiotic on or within three days after the initiating visit	Patient with a diagnosis of acute bronchitis/bronchiolitis did not have a prescription for an antibiotic on or within three days after the initiating visit	Low Value Care	Rendering	Contact National Committee for Quality Assurance
Concurrent Use of Opioids and Benzodiazepines	Patient(s) did not have concurrent use of prescription opioids and benzodiazepines	Patient did not have prescription opioid and benzodiazepine medications concurrently dispensed	Safety	Prescribing	Contact Partnership for Quality Measurement
Head Imaging for Uncomplicated Headache	Patient(s) with uncomplicated headache that did not have imaging studies	Patient with uncomplicated headache did not have imaging studies	Low Value Care	Rendering	Synopsis
Laryngoscopy with Treatment, Direct	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 90 days after the assessed procedure	Patient did not have a redo procedure within 90 days after the assessed procedure	Outcomes	Rendering	Synopsis
Laryngoscopy with Treatment, Flexible	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 90 days after the assessed procedure	Patient did not have a redo procedure within 90 days after the assessed procedure	Outcomes	Rendering	Synopsis
Laryngoscopy with Treatment, Indirect	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 90 days after the assessed procedure	Patient did not have a redo procedure within 90 days after the assessed procedure	Outcomes	Rendering	Synopsis
Medication Safety Monitoring	Older adult patients with dementia who did not use an antipsychotic, benzodiazepine, tricyclic antidepressant, nonbenzodiazepine hypnotic or anticholinergic agent after the earliest record of dementia	Patient with dementia did not have an antipsychotic, benzodiazepine, tricyclic antidepressant, nonbenzodiazepine hypnotic or anticholinergic agent dispensed after the earliest record of dementia	Safety	Prescribing	Contact Partnership for Quality Measurement

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Myringotomy, with General Anesthesia	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) with post-procedure confirmatory hearing test within 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient had a post-procedure confirmatory hearing test within 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Myringotomy, without General Anesthesia	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) with post-procedure confirmatory hearing test within 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient had a post-procedure confirmatory hearing test within 365 days after the assessed procedure	Outcomes	Rendering	Synopsis

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Nasal Ablation, Submucosal	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
Nasal Ablation, Superficial	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
Nasal Endoscopy with Treatment, Standard	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Nasal Endoscopy with Treatment, with Balloon Dilation	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Nasal Endoscopy with Treatment, with Repair of CSF Leak	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Nasal Vestibule Repair	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Risk of Continued Opioid Use	Patient(s) age 18-64 years who were opioid-naive and were not prescribed access to opioid medication for 15 or more days during the first 30 days following first opioid treatment initiation	Patient did not have opioid medication for 15 or more days during the first 30 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
	Patient(s) age 65 years and older who were opioid-naive and were not prescribed access to opioid medication for 15 or more days during the first 30 days following first opioid treatment initiation	Patient did not have opioid medication for 15 or more days during the first 30 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
	Patient(s) age 18-64 years who were opioid-naive and were not prescribed access to opioid medication for 31 or more days during the first 62 days following first opioid treatment initiation	Patient did not have opioid medication for 31 or more days during the first 62 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Risk of Continued Opioid Use	Patient(s) age 65 years and older who were opioid-naive and were not prescribed access to opioid medication for 31 or more days during the first 62 days following first opioid treatment initiation	Patient did not have opioid medication for 31 or more days during the first 62 days following initial opioid treatment	Safety	Prescribing	Contact National Committee for Quality Assurance
Septoplasty	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Thyroidectomy - Bilateral	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
Thyroidectomy - Single Side	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
Tonsillectomy and Adenoidectomy, Age 12 and Above	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Tonsillectomy and Adenoidectomy, Under Age 12	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Tonsillectomy, Age 12 and Above	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Tonsillectomy, Age 12 and Above	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Tonsillectomy, Under Age 12	Patient(s) without post-procedure complications within 30 days of the assessed procedure	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Turbinates Excision	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Tympanic Membrane Repair	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Tympanic Membrane Repair	Patient(s) without redo procedure between 181 to 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 181 and 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) with post-procedure confirmatory hearing test within 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient had a post-procedure confirmatory hearing test within 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Tympanoplasty	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) with post-procedure confirmatory hearing test within 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient had a post-procedure confirmatory hearing test within 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Tympanostomy, with General Anesthesia	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) with post-procedure confirmatory hearing test within 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient had a post-procedure confirmatory hearing test within 365 days after the assessed procedure	Outcomes	Rendering	Synopsis

Condition/procedure	Measure	Compliance criteria	Measure type	Attribution method	Source
Tympanostomy, without General Anesthesia	Patient(s) without post-procedure complications within 30 days of the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a complication related to the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) without redo procedure within 180 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient did not have a redo procedure within 180 days after the assessed procedure	Outcomes	Rendering	Synopsis
	Patient(s) with post-procedure confirmatory hearing test within 365 days after the assessed procedure (Note: evaluation for single-side and bilateral procedures is performed separately)	Patient had a post-procedure confirmatory hearing test within 365 days after the assessed procedure	Outcomes	Rendering	Synopsis
Use of High-Risk Medications in Older Adults	Patients 67 years and older who did not receive two or more of the same high-risk medications except for appropriate diagnosis in the last 12 reported months	Patient did not have two or more of the same high-risk medications except for the appropriate diagnosis dispensed	Safety	Prescribing	Contact Partnership for Quality Measurement
	Patients 67 years and older who did not receive two or more of the same high-risk medications from the same drug class in the last 12 reported months	Patient did not have two or more of the same high-risk medications from the same drug class dispensed	Safety	Prescribing	Contact Partnership for Quality Measurement
Use of Opioid Medications	Patient(s) 18 years or older without an average morphine milligram equivalent (MME) \geq 90mg/day during the treatment period	Patient did not have an average morphine equivalent dose \geq 90 mg/day	Safety	Prescribing	Contact National Committee for Quality Assurance
Use of Opioids from Multiple Providers	Patient(s) 18 years or older that did not fill opioid prescriptions from four or more different prescribers	Patient did not have opioid medications from four or more different prescribers dispensed	Safety	Prescribing	Contact National Committee for Quality Assurance

Important notes about UnitedHealth Premium®

The information from UnitedHealth Premium is not an endorsement of a particular physician or health care professional's suitability for the health care needs of any member. UnitedHealthcare does not practice medicine nor provide health care services. Physicians are solely responsible for medical judgments and treatments.

A Premium Care Physician designation does not guarantee the quality or the outcome of any health care services members receive. The fact that a physician does not have a Premium Care Physician designation does not mean the physician does not provide quality health care services.

All physicians in the UnitedHealthcare Network have met certain minimum credentialing requirements. Regardless of whether a physician has received a Premium Care Physician designation, members have access to all physicians in the UnitedHealthcare Network as described in the member's benefit plan.

There are various reasons why a physician may not be designated as a Premium Care Physician. A physician may not receive a designation because that physician has not been evaluated. This occurs when a physician does not practice in a specialty or market that is evaluated by Premium, or the physician's evaluation is in process. This also occurs when there are not enough measures, patients, and/or episodes attributed to the physician for evaluation. This is not an indicator of the total number of patients treated by the physician, or the number of procedures performed by the physician.

UnitedHealthcare informs members that designations are intended only as a guide when choosing a physician and should not be the sole factor in selecting a physician. Members are encouraged to discuss designations with a physician before choosing them or consult with their current physician(s) for advice on selecting other physicians.

As with all programs that evaluate performance based on evaluation of a sample, there is a risk of error. There is a risk of error in the claims data used and in the way patient care is attributed to physicians. UnitedHealth Premium uses statistical testing to compare a physician's performance to benchmarks. There is a risk of error in statistical tests when applied to the data and a result based on statistical testing is not a guarantee of correct inference or classification. Physicians have the opportunity to review the data and evaluation results and may submit requests for changes and/or corrections.

The information contained in this document is subject to change.