

# Omvo<sup>™</sup> (Mirikizumab-Mrkz)

**Policy Number:** 2024D0129A  
**Effective Date:** April 1, 2024

[➔ Instructions for Use](#)

Table of Contents	Page
<a href="#">Coverage Rationale</a> .....	1
<a href="#">Applicable Codes</a> .....	2
<a href="#">Background</a> .....	3
<a href="#">Benefit Considerations</a> .....	3
<a href="#">Clinical Evidence</a> .....	4
<a href="#">U.S. Food and Drug Administration</a> .....	5
<a href="#">References</a> .....	5
<a href="#">Policy History/Revision Information</a> .....	6
<a href="#">Instructions for Use</a> .....	6

Related Commercial Policies
<ul style="list-style-type: none"> <li><a href="#">Maximum Dosage and Frequency</a></li> <li><a href="#">Provider Administered Drugs – Site of Care</a></li> <li><a href="#">Self-Administered Medications</a></li> </ul>
Community Plan Policy
<ul style="list-style-type: none"> <li><a href="#">Omvo<sup>™</sup> (Mirikizumab-Mrkz)</a></li> </ul>

## Coverage Rationale

[➔ See Benefit Considerations](#)

**This policy refers to Omvo (mirikizumab-mrkz) injection.** Omvo (mirikizumab-mrkz) for self-administered subcutaneous injection is obtained under the pharmacy benefit.

### Ulcerative Colitis (UC)

**Omvo is proven for the treatment of ulcerative colitis when all of the following criteria are met:**

- Diagnosis of moderately to severely active ulcerative colitis; **and**
- Omvo is to be administered as three intravenous induction doses; **and**
- Omvo induction dosing is in accordance with the United States Food and Drug Administration (FDA) labeled dosing for UC; **and**
- Patient is not receiving Omvo in combination with another targeted immunomodulator [e.g., Enbrel (etanercept), Cimzia (certolizumab), Simponi (golimumab), Orencia (abatacept), adalimumab, Xeljanz (tofacitinib), Olumiant (baricitinib), Rinvoq (upadacitinib), Stelara (ustekinumab), Skyrizi (risankizumab)]  
**and**
- Authorization will be issued for 3 induction doses

**Omvo is medically necessary for the treatment of ulcerative colitis when all of the following criteria are met:**

- Diagnosis of moderately to severely active ulcerative colitis; **and**
- **One** of the following:
  - Patient has had prior or concurrent inadequate response to a therapeutic course of oral corticosteroids and/or immunosuppressants (e.g., azathioprine, 6-mercaptopurine); **or**
  - Patient has been previously treated with a targeted immunomodulator FDA-approved for the treatment of ulcerative colitis [e.g., adalimumab, Simponi (golimumab), Stelara (ustekinumab), Xeljanz (tofacitinib), Rinvoq (upadacitinib)]**and**
- History of failure, contraindication, or intolerance to **two** of the following preferred products (document drug, date, and duration of trial) (for Medicare reviews, refer to the [CMS](#) section\*):

- One of the preferred adalimumab products (for a list of preferred adalimumab products, refer to the member specific benefit plan document for drug coverage guidelines)
- Simponi (golimumab)
- Stelara (ustekinumab)
- Xeljanz/Xeljanz XR (tofacitinib)
- Rinvoq (upadacitinib)
- Omvoh is to be administered as three intravenous induction doses; **and**
- Omvoh induction dosing is in accordance with the U.S FDA labeled dosing for UC; **and**
- Patient is not receiving Omvoh in combination with another targeted immunomodulator [e.g., Enbrel (etanercept), Cimzia (certolizumab), Simponi (golimumab), Orencia (abatacept), adalimumab, Xeljanz (tofacitinib), Olumiant (baricitinib), Rinvoq (upadacitinib), Stelara (ustekinumab), Skyrizi (risankizumab)]; **and**
- Prescribed by or in consultation with a gastroenterologist; **and**
- Authorization will be issued for 3 induction doses

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
C9168	Injection, mirikizumab-mrkz, 1 mg
J3490	Unclassified drug
J3590	Unclassified biologics

Diagnosis Code	Description
K51.00	Ulcerative (chronic) pancolitis without complications
K51.011	Ulcerative (chronic) pancolitis with rectal bleeding
K51.012	Ulcerative (chronic) pancolitis with intestinal obstruction
K51.013	Ulcerative (chronic) pancolitis with fistula
K51.014	Ulcerative (chronic) pancolitis with abscess
K51.018	Ulcerative (chronic) pancolitis with other complication
K51.019	Ulcerative (chronic) pancolitis with unspecified complications
K51.20	Ulcerative (chronic) proctitis without complications
K51.211	Ulcerative (chronic) proctitis with rectal bleeding
K51.212	Ulcerative (chronic) proctitis with intestinal obstruction
K51.213	Ulcerative (chronic) proctitis with fistula
K51.214	Ulcerative (chronic) proctitis with abscess
K51.218	Ulcerative (chronic) proctitis with other complication
K51.219	Ulcerative (chronic) proctitis with unspecified complications
K51.30	Ulcerative (chronic) recto sigmoiditis without complications
K51.311	Ulcerative (chronic) recto sigmoiditis with rectal bleeding
K51.312	Ulcerative (chronic) recto sigmoiditis with intestinal obstruction
K51.313	Ulcerative (chronic) recto sigmoiditis with fistula
K51.314	Ulcerative (chronic) recto sigmoiditis with abscess
K51.318	Ulcerative (chronic) recto sigmoiditis with other complication

Diagnosis Code	Description
K51.319	Ulcerative (chronic) recto sigmoiditis with unspecified complications
K51.40	Inflammatory polyps of colon without complications
K51.411	Inflammatory polyps of colon with rectal bleeding
K51.412	Inflammatory polyps of colon with intestinal obstruction
K51.413	Inflammatory polyps of colon with fistula
K51.414	Inflammatory polyps of colon with abscess
K51.418	Inflammatory polyps of colon with other complication
K51.419	Inflammatory polyps of colon with unspecified complications
K51.50	Left sided colitis without complications
K51.511	Left sided colitis with rectal bleeding
K51.512	Left sided colitis with intestinal obstruction
K51.513	Left sided colitis with fistula
K51.514	Left sided colitis with abscess
K51.518	Left sided colitis with other complication
K51.519	Left sided colitis with unspecified complications
K51.80	Other ulcerative colitis without complications
K51.811	Other ulcerative colitis with rectal bleeding
K51.812	Other ulcerative colitis with intestinal obstruction
K51.813	Other ulcerative colitis with fistula
K51.814	Other ulcerative colitis with abscess
K51.818	Other ulcerative colitis with other complication
K51.819	Other ulcerative colitis with unspecified complications
K51.90	Ulcerative colitis, unspecified, without complications
K51.911	Ulcerative colitis, unspecified with rectal bleeding
K51.912	Ulcerative colitis, unspecified with intestinal obstruction
K51.913	Ulcerative colitis, unspecified with fistula
K51.914	Ulcerative colitis, unspecified with abscess
K51.918	Ulcerative colitis, unspecified with other complication
K51.919	Ulcerative colitis, unspecified with unspecified complications
K52.1	Toxic gastroenteritis and colitis

## Background

Omvoh is a humanized IgG4 monoclonal antibody that selectively binds to the p19 subunit of human IL-23 cytokine and inhibits its interaction with the IL-23 receptor. IL-23 is involved in mucosal inflammation and affects the differentiation, expansion, and survival of T cell subsets, and innate immune cell subsets, which represent sources of pro-inflammatory cytokines.

## Benefit Considerations

Some Certificates of Coverage allow for coverage of experimental/investigational/unproven treatments for life-threatening illnesses when certain conditions are met. The member specific benefit plan document must be consulted to make coverage decisions for this service. Some states mandate benefit coverage for off-label use of medications for some diagnoses or under some circumstances when certain conditions are met. Where such mandates apply, they supersede language in the benefit document or in the medical or drug policy. Benefit coverage for an otherwise unproven service for the treatment of serious rare

diseases may occur when certain conditions are met. Refer to the Policy and Procedure addressing the treatment of serious rare diseases.

## Clinical Evidence

### Proven

#### *Ulcerative Colitis*

The safety and efficacy of mirikizumab-mrkz was evaluated in two randomized, double-blind, placebo-controlled clinical studies, one induction study [UC-1 (NCT03518086)] and one maintenance study [UC-2 (NCT03524092)], in adult subjects with moderately to severely active ulcerative colitis who had inadequate response, loss of response, or failed to tolerate any of the following: corticosteroids, 6-mercaptopurine, azathioprine, biologic therapy (TNF blocker, vedolizumab), or tofacitinib. The 12-week intravenous induction study (UC-1) was followed by the 40-week subcutaneous randomized withdrawal maintenance study (UC-2).

#### Study UC-1

In UC-1, efficacy was evaluated in 1062 subjects who were randomized 3:1 at Week 0 to receive 300 mg mirikizumab-mrkz or placebo by intravenous infusion at Week 0, Week 4, and Week 8. Subjects had a mean age of 43 years (range 18 to 79 years); 40% were female; and 71% identified as White, 25% as Asian, 1% as American Indian or Alaska Native, 1% as Black or African American, and < 2% as another racial group or did not report their racial group. Subjects were permitted to use stable doses of aminosalicylates, immunomodulators (6-mercaptopurine, azathioprine, methotrexate), and oral corticosteroids (prednisone  $\leq$  20 mg/day or equivalent, extended-release budesonide 9 mg/day, beclomethasone dipropionate 5 mg/day). At baseline, 41% of subjects were receiving oral corticosteroids, 24% were receiving immunomodulators, and 75% were receiving aminosalicylates.

At baseline, 57% were biologic and Janus Kinase inhibitor (JAKi) naive, 41% had failed at least one biologic, 3% had failed a JAKi, and 2% had previously received but had not failed a biologic or JAKi.

Disease activity was assessed based on the modified Mayo score (mMS), which ranges from 0 to 9 and has three subscores that are each scored from 0 (normal) to 3 (most severe): stool frequency, rectal bleeding, and findings on centrally read endoscopy subscore. At baseline, subjects had a mMS of 5 to 9, including a centrally read endoscopy subscore of 2 or 3. An endoscopy subscore of 2 was defined by marked erythema, absent vascular pattern, friability, and erosions; and a subscore of 3 was defined by spontaneous bleeding and ulceration. Subjects had a median mMS of 7, and 58% had severely active disease (mMS of 7 to 9).

The primary endpoint was clinical remission at Week 12. The secondary endpoints were clinical response, endoscopic improvement, and histologic-endoscopic mucosal improvement.

Study UC-1 was not designed to evaluate the relationship of histologic-endoscopic mucosal improvement at Week 12 to disease progression and long-term outcomes.

#### *Rectal Bleeding and Stool Frequency Subscores*

Decreases in rectal bleeding and stool frequency subscores were observed as early as Week 3 in subjects treated with OMVOH compared to subjects on placebo.

After 12 weeks of induction, 65% of patients achieved clinical response and 24% achieved clinical remission compared to placebo (43% and 15%, for clinical response and clinical remission, respectively). Decreases in rectal bleeding and stool frequency subscores were observed as early as week 3 in patients treated with mirikizumab compared to patients on placebo.

#### Study UC-2

The maintenance study (UC-2) evaluated 506 subjects who achieved clinical response at Week 12 in Study UC-1. These subjects were randomized 2:1 to receive 200 mg mirikizumab-mrkz or placebo subcutaneously every 4 weeks for 40 weeks in UC2, for a total of 52 weeks of treatment. Subjects who were on concomitant ulcerative colitis therapies during UC-1 were required to continue on stable doses of oral aminosalicylates and immunomodulators (6-mercaptopurine, azathioprine,

methotrexate). Corticosteroid tapering was required for subjects who were receiving corticosteroids at baseline and achieved clinical response in UC-1.

The primary endpoint was clinical remission at Week 40. The secondary endpoints were endoscopic improvement, maintenance of clinical remission in subjects who achieved clinical remission at Week 12, corticosteroid-free clinical remission, and histologic-endoscopic mucosal improvement.

Study UC-2 was not designed to evaluate the relationship of histologic-endoscopic mucosal improvement at Week 40 to disease progression and long-term outcomes.

### *Bowel Urgency*

Bowel urgency was assessed during UC-1 and UC-2 with an Urgency Numeric Rating Scale (NRS) of 0 to 10. A greater proportion of subjects with a baseline Urgency NRS weekly average score  $\geq 3$  treated with OMVOH compared to placebo reported an Urgency NRS weekly average score of 0 or 1 (39% versus 23%) at Week 40. Urgency NRS weekly average scores of 0 to 1 were also observed in a greater proportion of subjects treated with OMVOH compared to placebo at Week 12.

### *Endoscopic Assessment*

Normalization of the endoscopic appearance of the mucosa (endoscopic remission) was defined as a Mayo endoscopic subscore of 0. At Week 40 in UC-2, endoscopic remission was observed in a greater proportion of subjects treated with OMVOH compared to placebo (22% versus 14%).

Among those who achieved clinical response at 12 weeks, 51% of all patients and 45% of patients who failed prior treatment with a biologic or Janus kinase inhibitor achieved clinical remission at 52 weeks compared to placebo (27% and 15%, respectively). Of the patients who achieved clinical response at 12 weeks, 50% achieved steroid-free clinical remission at 52 weeks, compared to 27% of patients receiving placebo. Patients in steroid-free clinical remission were steroid-free for at least 12 weeks prior to the end of the 52-week assessment. Among patients who achieved clinical remission at 12 weeks, 66% of patients maintained clinical remission through 1 year of continuous treatment compared 40% of patients receiving placebo.

## **U.S. Food and Drug Administration (FDA)**

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

OmvoH is an interleukin-23 antagonist indicated for the treatment of moderately to severely active ulcerative colitis in adults.

## **Centers for Medicare and Medicaid Services (CMS)**

Medicare does not have a National Coverage Determination (NCD) for OmvoH™ (mirikizumab-mrkz). Local Coverage Determinations/Articles (LCDs)/LCAs that refer to the intravenous administration of OmvoH™ (mirikizumab-mrkz) do not exist.

In general, Medicare covers outpatient (Part B) drugs that are furnished "incident to" a physician's service provided that the drugs are not usually self-administered by the patients who take them. Refer to the [Medicare Benefit Policy Manual, Chapter 15, §50 - Drugs and Biologicals](#). (Accessed December 6, 2023)

\*Preferred therapy criteria for Medicare Advantage members, refer to [Medicare Part B Step Therapy Programs](#).

## **References**

1. OmvoH [package insert]. Indianapolis, IN: Eli Lilly and Company; October 2023.
2. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology*. 2020 Jan 13.
3. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol*. 2019 Mar;114(3):384-413.Yese.

## Policy History/Revision Information

Date	Summary of Changes
04/01/2024	<ul style="list-style-type: none"><li data-bbox="337 220 756 249">• New Medical Benefit Drug Policy</li></ul>

## Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

This Medical Benefit Drug Policy may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence ([Medicare IOM Pub. No. 100-16, Ch. 4, §90.5](#)).

UnitedHealthcare may also use tools developed by third parties, such as the InterQual<sup>®</sup> criteria, to assist us in administering health benefits. UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.