

PREVENTIVE CARE SERVICES

Guideline Number: CDG.016.27

Effective Date: October 1, 2019

[Instructions for Use](#) ⓘ

Table of Contents	Page
COVERAGE RATIONALE	1
FREQUENTLY ASKED QUESTIONS	3
DEFINITIONS	4
APPLICABLE CODES	4
REFERENCES	40
GUIDELINE HISTORY/REVISION INFORMATION	41
INSTRUCTIONS FOR USE	41

Related Commercial Policies

- [Breast Imaging for Screening and Diagnosing Cancer](#)
- [Cardiovascular Disease Risk Tests](#)
- [Computed Tomographic Colonography](#)
- [Consultation Services Policy](#)
- [Cytological Examination of Breast Fluids for Cancer Screening](#)
- [Genetic Testing for Hereditary Cancer](#)
- [Preventive Medicine and Screening Policy](#)
- [Vaccines](#)

COVERAGE RATIONALE

Indications for Coverage

Introduction

UnitedHealthcare covers certain medical services under the preventive care services benefit. The federal Patient Protection and Affordable Care Act (PPACA) requires non-grandfathered health plans to cover certain “recommended preventive services” as identified by PPACA under the preventive care services benefit, without cost sharing to members when provided by network physicians. This includes:

- Evidence-based items or services that have in effect a rating of “A” or “B” in the current recommendations of the United States Preventive Services Task Force.
- Immunizations for routine use in children, adolescents and adults that have in effect a recommendation from the *Advisory Committee on Immunization Practices of the Centers for Disease Control and Prevention*.
- With respect to infants, children and adolescents, evidence-informed preventive care and screenings provided for in the comprehensive guidelines supported by the *Health Resources and Services Administration*.
- With respect to women, such additional preventive care and screenings as provided for in comprehensive guidelines supported by the *Health Resources and Services Administration*.

Member Cost-Sharing

- **Non-Grandfathered Plans:**
 - Non-grandfathered plans provide coverage for preventive care services with no member cost sharing (i.e. covered at 100% of Allowed Amounts without deductible, coinsurance or copayment) when services are obtained from a Network provider.
 - Under PPACA, services obtained from an out-of-network provider are not required to be covered under a plan’s preventive benefit, and may be subject to member cost sharing. Refer to the member specific benefit plan document for out-of-network benefit information, if any.
- **Grandfathered Plans:**
 - Plans that maintain grandfathered status under PPACA are not required by law to provide coverage for these preventive services without member cost sharing; although a grandfathered plan may choose to voluntarily amend its plan document to include these preventive benefits.
 - Except where there are state mandates, a grandfathered plan might include member cost sharing, or exclude some of the preventive care services identified under PPACA.
 - Refer to the member specific benefit plan document for details on how benefits are covered under a grandfathered plan.

Preventive vs. Diagnostic Services

Certain services can be done for preventive or diagnostic reasons. When a service is performed for the purpose of preventive screening and is appropriately reported, it will be considered under the preventive care services benefit.

This includes services directly related to the performance of a covered preventive care service (see the Frequently Asked Questions section for additional information.)

Preventive services are those performed on a person who:

- has not had the preventive screening done before and does not have symptoms or other abnormal studies suggesting abnormalities; or
- has had screening done within the recommended interval with the findings considered normal; or
- has had diagnostic services results that were normal after which the physician recommendation would be for future preventive screening studies using the preventive services intervals.

When a service is done for diagnostic purposes it will be considered under the applicable non-preventive medical benefit. Diagnostic services are done on a person who:

- had abnormalities found on previous preventive or diagnostic studies that require further diagnostic studies; or
- had abnormalities found on previous preventive or diagnostic studies that would recommend a repeat of the same studies within shortened time intervals from the recommended preventive screening time intervals; or
- had a symptom(s) that required further diagnosis; or
- does not fall within the applicable population for a recommendation or guideline

Covered Breastfeeding Equipment

Personal-use electric breast pump:

- The purchase of a personal-use electric breast pump (HCPCS code E0603).
 - This benefit is limited to one pump per birth. In the case of a birth resulting in multiple infants, only one breast pump is covered.
 - A breast pump purchase includes the necessary supplies for the pump to operate.
- Replacement breast pump supplies necessary for the personal-use electric breast pump to operate. This includes: standard power adaptor, tubing adaptors, tubing, locking rings, bottles specific to breast pump operation, caps for bottles that are specific to the breast pump, valves, filters, and breast shield and/or splash protector for use with the breast pump.

Coverage Limitations and Exclusions

- Services not covered under the preventive care benefit may be covered under another portion of the medical benefit plan.
- The coverage outlined in this guideline does not address certain outpatient prescription medications, tobacco cessation drugs and/or over the counter items, as required by PPACA. These preventive benefits are administered by the member's pharmacy plan administrator. For details on coverage, refer to the member-specific pharmacy plan administrator.
- A vaccine (immunization) is not covered if it does not meet company vaccine policy requirements for FDA labeling and if it does not have explicit ACIP recommendations for routine use published in the Morbidity and Mortality Weekly Report (MMWR) of the Centers for Disease Control and Prevention (CDC).
- Examinations, screenings, testing, or vaccines (immunizations) are not covered when:
 - required solely for the purposes of career or employment, school or education, sports or camp, travel (including travel vaccines (immunizations)), insurance, marriage or adoption; or
 - related to judicial or administrative proceedings or orders; or
 - conducted for purposes of medical research; or
 - required to obtain or maintain a license of any type.
- Services that are investigational, experimental, unproven or not medically necessary are not covered.
- Breastfeeding equipment and supplies not listed above. This includes, but is not limited to:
 - Manual breast pumps and all related equipment and supplies.
 - Hospital-grade breast pumps and all related equipment and supplies.
 - Equipment and supplies not listed in the [Covered Breastfeeding Equipment](#) section above, including but not limited to:
 - Batteries, battery-powered adaptors, and battery packs.
 - Electrical power adapters for travel.
 - Bottles which are not specific to breast pump operation. This includes the associated bottle nipples, caps and lids.
 - Travel bags, and other similar travel or carrying accessories.
 - Breast pump cleaning supplies including soap, sprays, wipes, steam cleaning bags and other similar products.
 - Baby weight scales.
 - Garments or other products that allow hands-free pump operation.
 - Breast milk storage bags, ice-packs, labels, labeling lids, and other similar products.
 - Nursing bras, bra pads, breast shells, nipple shields, and other similar products.

- Creams, ointments, and other products that relieve breastfeeding related symptoms or conditions of the breasts or nipples.

Note: See the [Indications for Coverage](#) section above for covered breastfeeding equipment.

FREQUENTLY ASKED QUESTIONS

1	Q:	If woman has an abnormal finding on a preventive screening mammography and the follow up mammogram was found to be normal, will UnitedHealthcare cover her future mammograms under the preventive care services benefit?
	A:	Yes, if the member was returned to normal mammography screening protocol, her future mammography screenings would be considered under the preventive care services benefit.
2	Q:	If a polyp is encountered during a preventive screening colonoscopy, are future colonoscopies considered under the preventive care services benefit?
	A:	No. If a polyp is removed during a preventive screening colonoscopy, future colonoscopies would normally be considered to be diagnostic because the time intervals between future colonoscopies would be shortened.
3	Q:	If a member had elevated cholesterol on a prior preventive screening, are future cholesterol tests considered under the preventive care services benefit?
	A:	Once the diagnosis has been made, further testing is considered diagnostic rather than preventive. This is true whether or not the member is receiving pharmacotherapy.
4	Q:	Are the related therapeutic services for a preventive colonoscopy covered under the preventive care benefit?
	A:	Yes, related services integral to a colonoscopy are covered under the preventive care services benefit including: pre-operative examination, the associated facility, anesthesia, polyp removal (if necessary), pathologist and physician fees. However, the preventive benefit does not include a post-operative examination.
5	Q:	Are the related services for a woman's outpatient sterilization procedure covered under the preventive care benefit?
	A:	Yes, related services for a woman's outpatient sterilization are covered under the preventive care services benefit including: associated implantable devices, facility fee, anesthesia, and surgeon/physician fees. Note the following clarifications: <ul style="list-style-type: none"> • The preventive benefit does not include a pre- or post-operative examination. • If a woman is admitted to an inpatient facility for another reason, and has a sterilization performed during that admission, the sterilization fees (surgical fee, device fee, anesthesia, pathologist and physician fees), are covered under the preventive benefit. However, the facility fees are not covered under the preventive care benefit since the sterilization is incidental to, and is not the primary reason, for the admission. • For hysteroscopic fallopian tube occlusion sterilization procedures, the preventive benefit includes an outpatient, followup hysterosalpingogram to confirm that the fallopian tubes are completely blocked.
6	Q:	Are blood draws/venipunctures included in the preventive care benefit?
	A:	Yes, blood draws/venipunctures are considered under the preventive benefit if billed for a covered preventive lab services that requires a blood draw.
7	Q:	Do any preventive care services require prior-authorization?
	A:	Certain services require prior-authorization on most benefit plans. This includes, but may not be limited to: BRCA lab screening, computed tomographic colonography (virtual colonoscopy), and screening for lung cancer with low-dose computed tomography.
8	Q:	Is a newly-combined vaccine (a vaccine with several individual vaccines combined into one) covered under preventive care benefits?
	A:	A new vaccine that is pending ACIP recommendations, but is a combination of previously approved individual components, may be eligible under the preventive care benefit.
9	Q:	Are preventive care services affected by other policies?
	A:	Yes, including for example, the Reimbursement Policy titled Preventive Medicine and Screening Policy describes situations which may affect reimbursement of preventive care services.

10	Q:	Are travel vaccines covered under preventive care benefits?
	A:	Benefits for preventive care services include vaccines for routine use in children, adolescents and adults that have in effect a recommendation from ACIP with respect to the individual involved. Vaccines that are specific to travel (e.g. typhoid, yellow fever, cholera, plague, and Japanese encephalitis virus) are excluded from the preventive care services benefit.
11	Q:	For preventive services that have a diagnosis code requirement, does the listed diagnosis code need to be the primary diagnosis on the claim?
	A:	In general, most preventive services do not require the preventive diagnosis code to be in the primary position. However, certain preventive services do require the diagnosis code to be in the primary position, which include: (1) Chemoprevention of Breast Cancer (Counseling), and (2) Genetic Counseling and Evaluation for BRCA Testing.
12	Q:	Does the preventive care services benefit include prescription or over the counter (OTC) items?
	A:	Refer to the plan's pharmacy benefit plan administrator for details on prescription medications and OTCs available under the plan's preventive benefit.

DEFINITIONS

The following definitions may not apply to all plans. Refer to the member specific benefit plan document for applicable definitions.

Modifier 33: Preventive service; when the primary purpose of the service is the delivery of an evidence based service in accordance with a US Preventive Services Task Force A or B rating in effect and other preventive services identified in preventive services mandates (legislative or regulatory), the service may be identified by adding 33 to the procedure. For separately reported services specifically identified as preventive, the modifier should not be used.

Note: UnitedHealthcare considers the procedures and diagnostic codes and Preventive Benefit Instructions listed in the table below in determining whether preventive care benefits apply. While Modifier 33 may be reported, it is not used in making preventive care benefit determinations.

Acronyms

Throughout this document the following acronyms are used:

- USPSTF: United States Preventive Services Task Force
- PPACA: Patient Protection and Affordable Care Act of 2010
- ACIP: Advisory Committee on Immunization Practices
- HRSA: Health Resources and Services Administration

APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this guideline does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Coverage Determination Guidelines may apply.

CPT® is a registered trademark of the American Medical Association

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
Abdominal Aortic Aneurysm Screening <u>USPSTF Rating (June 2014): B</u> The USPSTF recommends one-time screening for abdominal	Procedure Code(s): <i>Ultrasound Screening Study for Abdominal Aortic Aneurysm:</i> 76706	Age 65 through 75 (ends on 76 th birthday). Requires at least one of the diagnosis codes listed in this row.

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
aortic aneurysm (AAA) by ultrasonography in men ages 65 to 75 years who have ever smoked.	Diagnosis Code(s): F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891	
Bacteriuria Screening <u>USPSTF Rating (July 2008): A</u> Screening for asymptomatic bacteriuria with urine culture for pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	Procedure Code(s): 81007, 87086, 87088 Diagnosis Code(s): Pregnancy Diagnosis Codes	Requires a Pregnancy Diagnosis Code .
Chlamydia Infection Screening <u>USPSTF Rating (Sept. 2014): B</u> The USPSTF recommends screening for chlamydia in sexually active women age 24 years and younger and in older women who are at increased risk for infection. Notes: This recommendation applies to all sexually active adolescents and adult women, including pregnant women. Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11 years – 21 years.	Procedure Code(s): <i>Chlamydia Infection Screening:</i> 86631, 86632, 87110, 87270, 87320, 87490, 87491, 87492, 87801, 87810 <i>Blood Draw:</i> 36415, 36416 Blood draw codes only apply to lab codes 86631 or 86632 Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes OR <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Other: Z11.3, Z11.8, Z11.9, Z20.2	<i>Chlamydia Infection Screening:</i> Requires a Pregnancy Diagnosis Code OR one of the Screening diagnosis codes listed in this row. <i>Blood Draw:</i> Required to be billed with 86631 or 86632 AND <ul style="list-style-type: none"> One of the Screening diagnosis codes listed in this row OR With a Pregnancy Diagnosis Code.
Gonorrhea Screening <u>USPSTF Rating (Sept. 2014): B</u> The USPSTF recommends screening for gonorrhea in sexually active women age 24 years and younger and in older women who are at increased risk for infection. Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11 years – 21 years.	Procedure Code(s): 87590, 87591, 87592, 87801, 87850 Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes OR <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Other: Z11.3, Z11.9, Z20.2	Requires either a Pregnancy Diagnosis Code OR one of the Screening diagnosis codes listed in this row.
Hepatitis B Virus Infection Screening <i>Pregnant Women:</i> <u>USPSTF Rating (June 2009): A</u> Screening for hepatitis B virus	Procedure Code(s): <i>Hepatitis B Virus Infection Screening:</i> 87340, 87341, G0499	<i>Hepatitis B Virus Infection Screening:</i> Requires a Pregnancy Diagnosis Code OR one of the Screening diagnosis codes listed in this row.

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>(HBV) infection in pregnant women at their first prenatal visit.</p> <p><i>Persons at High Risk:</i> <u>USPSTF Rating (May 2014): B</u> The USPSTF recommends screening for hepatitis B virus (HBV) infection in persons at high risk for infection.</p> <p>Also see the Medical Policy titled Hepatitis Screening.</p>	<p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes OR <i>Screening:</i> Z00.00, Z00.01, Z11.59, Z57.8</p>	<p><i>Blood Draw:</i> Requires one of the listed Hepatitis B Virus Infection Screening procedure codes listed in this row AND</p> <ul style="list-style-type: none"> • A Pregnancy Diagnosis Code OR • One of the Screening diagnosis codes listed in this row.
<p>Hepatitis C Virus Infection Screening</p> <p><u>USPSTF Rating (June 2013): B</u> The USPSTF recommends screening for hepatitis C virus (HCV) infection in persons at high risk for infection. The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965.</p> <p>Also see the Medical Policy titled Hepatitis Screening.</p>	<p>Procedure Code(s): <i>Hepatitis C Virus Infection Screening:</i> 86803, 86804, G0472</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Hepatitis C Virus Infection Diagnosis Codes</p>	<p><i>Hepatitis C Virus Infection Screening:</i> Requires one of the Hepatitis C Virus Infection Diagnosis Codes.</p> <p><i>Blood Draw:</i> Requires one of the Hepatitis C Virus Infection Screening procedure codes listed in this row AND a Hepatitis C Virus Infection Diagnosis Code.</p>
<p>HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults</p> <p><u>USPSTF Rating (June 2019): A</u> The USPSTF recommends that clinicians screen for HIV infection in:</p> <ul style="list-style-type: none"> • Adolescents and adults aged 15 to 65 years. Younger adolescents and older adults who are at increased risk of infection should also be screened. • All pregnant persons, including those who present in labor or at delivery whose HIV status is unknown. <p>Note: Bright Futures recommends HIV screening lab work be conducted once between ages 15–18 years. Also recommended anytime between ages 11–14 years, and 19–21 years when a risk assessment is positive.</p>	<p>Procedure Code(s): <i>HIV (Human Immunodeficiency Virus) Screening:</i> 86689, 86701, 86702, 86703, 87389, 87390, 87391, 87806, G0432, G0433, G0435, G0475, S3645</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes OR <i>Screening:</i> <i>Adult: Z00.00, Z00.01</i> <i>Child: Z00.121, Z00.129,</i> <i>Other: Z11.3, Z11.4, Z11.59, Z11.9, Z20.6, Z22.6, Z22.8, Z22.9</i></p> <p>Also see Expanded Women's Preventive Health section.</p>	<p>No age limits.</p> <p><i>HIV – Human Immunodeficiency Virus – Screening:</i> Requires a Pregnancy Diagnosis Code OR one of the Screening diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Requires both of the following:</p> <ul style="list-style-type: none"> • One of the listed HIV Screening procedure codes listed in this row AND • One of the Screening diagnosis codes listed in this row OR a Pregnancy Diagnosis Code

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>RH Incompatibility Screening</p> <p><u>USPSTF Rating (Feb. 2004): A</u> Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.</p> <p><u>USPSTF Rating (Feb. 2004): B</u> Repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.</p>	<p>Procedure Code(s): <i>RH Incompatibility Screening:</i> 86901</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Pregnancy Diagnosis Codes</p>	<p><i>RH Incompatibility Screening:</i> Requires a Pregnancy Diagnosis Code.</p> <p><i>Blood Draw:</i> Required to be billed with 86901 AND with a Pregnancy Diagnosis Code.</p>
<p>Syphilis Screening</p> <p><i>Non-Pregnant Adults and Adolescents at Increased Risk:</i> <u>USPSTF Rating (June 2016): A</u> The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection (asymptomatic, nonpregnant adults and adolescents who are at increased risk for syphilis infection).</p> <p><i>Pregnant Women:</i> <u>USPSTF Rating (Sept. 2018): A</u> The USPSTF recommends early screening for syphilis infection in all pregnant women.</p> <p>Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.</p>	<p>Procedure Code(s): <i>Syphilis Screening:</i> 86592, 86593</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes OR <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Other: Z11.2, Z11.3, Z11.9, Z20.2</p>	<p><i>Syphilis Screening:</i> Requires a Pregnancy Diagnosis Code OR one of the Screening diagnosis code listed in this row.</p> <p><i>Blood Draw:</i> Requires both of the following:</p> <ul style="list-style-type: none"> • One of the listed Syphilis Screening procedure codes listed in this row AND • One of the Screening diagnosis codes listed in this row OR a Pregnancy Diagnosis Code.
<p>Genetic Counseling and Evaluation for BRCA Testing; and BRCA Lab Screening</p> <p><u>USPSTF Rating (Dec. 2013): B</u> The USPSTF recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with one of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast</p>	<p>Genetic Counseling and Evaluation</p> <p>Procedure Code(s): <i>Medical Genetics and Genetic Counseling Services:</i> 96040, S0265</p> <p><i>Evaluation and Management (Office Visits):</i> 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397, G0463</p>	<p>Genetic Counseling and Evaluation</p> <p>*Medical Necessity plans require genetic counseling before BRCA Lab Screening.</p> <p>Requires one of the Genetic Counseling and Evaluation diagnosis codes listed in this row in the primary position.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>cancer susceptibility genes (<i>BRCA1</i> or <i>BRCA2</i>). Women with positive screening results should receive genetic counseling and, if indicated after counseling, BRCA testing.</p> <p>See the Medical Policy titled Genetic Testing for Hereditary Cancer.</p>	<p>Diagnosis Code(s): Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43</p> <p>BRCA Lab Screening Procedure Code(s): 81162, 81163, 81164, 81165, 81166, 81167, 81212, 81215, 81216, 81217</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Family History or Personal History of breast cancer and/or ovarian cancer:</i> Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43</p>	<p>BRCA Lab Screening</p> <p>*Prior authorization requirements apply to BRCA lab screening.</p> <p>Applies to age 18+ when billed with one of the BRCA Lab Screening diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Requires one of the BRCA Lab Screening procedure codes listed in this row AND one of the BRCA Lab Screening diagnosis codes listed in this row.</p>
<p>Diabetes Screening</p> <p><u>USPSTF Rating (Oct. 2015): B</u> The USPSTF recommends screening for abnormal blood glucose as part of cardiovascular risk assessment in adults aged 40 to 70 years who are overweight or obese. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity.</p> <p>See Behavioral Counseling in Primary Care to Promote a Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors for intensive behavioral counseling interventions.</p> <p>For additional diabetes screening benefits, also see the <i>Expanded Women's Preventive Health</i> section for Screening for Gestational Diabetes Mellitus and Screening for Diabetes Mellitus After Pregnancy.</p>	<p>Procedure Code(s): <i>Diabetes Screening:</i> 82947, 82948, 82950, 82951, 82952, 83036</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Required Diagnosis Codes (requires at least one):</i> Z00.00, Z00.01, Z13.1</p> <p>AND One of the following additional diagnosis codes as follows: <i>Additional Diagnosis Codes (requires at least one):</i> <i>Overweight:</i> E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29</p> <p><i>Obesity :</i> E66.01, E66.09, E66.1, E66.8, E66.9, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p> <p><i>Body Mass Index 30.0 – 39.9:</i> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39</p> <p><i>Body Mass Index 40.0 and Over:</i> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p>	<p>Limited to age 40-70 years (ends on 71st birthday).</p> <p><i>Diabetes Screening:</i> Requires one of the Required Diagnosis Codes listed in this row AND one of the listed Additional Diagnosis Codes in this row.</p> <p><i>Blood Draw:</i> Requires ALL of the following:</p> <ul style="list-style-type: none"> • One of the listed Diabetes Screening procedure codes listed in this row AND • One of the listed Required Diagnosis Codes AND • One of the listed Additional Diagnosis Codes. <p>Preventive Benefit Does Not Apply: If a Diabetes Diagnosis Code is present in any position, the preventive benefit does not apply; see the Diabetes Diagnosis Code List.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
	<p><i>Essential Hypertension:</i> I10</p> <p><i>Hypertensive Heart Disease:</i> I11.0, I11.9</p> <p><i>Hypertensive Chronic Kidney Disease:</i> I12.0, I12.9</p> <p><i>Hypertensive Heart and Chronic Kidney Disease:</i> I13.0, I13.10, I13.11, I13.2</p> <p><i>Secondary Hypertension:</i> I15.0, I15.1, I15.2, I15.8, I15.9, N26.2</p> <p><i>Hypertension Complicating Pregnancy, Childbirth and the Puerperium:</i> O10.011, O10.012, O10.013, O10.019, O10.02, O10.03, O10.111, O10.112, O10.113, O10.119, O10.12, O10.13, O10.211, O10.212, O10.213, O10.219, O10.22, O10.23, O10.311, O10.312, O10.313, O10.319, O10.32, O10.33, O10.411, O10.412, O10.413, O10.419, O10.42, O10.43, O10.911, O10.912, O10.913, O10.919, O10.92, O10.93, O11.1, O11.2, O11.3, O11.4, O11.5, O11.9, O13.1, O13.2, O13.3, O13.4, O13.5, O13.9, O16.1, O16.2, O16.3, O16.4, O16.5, O16.9</p> <p><i>Urgent/Emergency/Crisis Hypertension</i> I16.0, I16.1, I16.9</p> <p>See the <i>Expanded Women's Preventive Health</i> section for Screening for Gestational Diabetes Mellitus and Screening for Diabetes Mellitus After Pregnancy.</p>	
<p>Gestational Diabetes Mellitus Screening</p> <p><u>USPSTF Rating (Jan. 2014): B</u> The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant women after 24 weeks of gestation.</p>	<p>See the <i>Expanded Women's Preventive Health</i> section for Screening for Gestational Diabetes Mellitus codes.</p>	<p>See the <i>Expanded Women's Preventive Health</i> section for Screening for Gestational Diabetes Mellitus preventive benefit instructions.</p> <p>Note: This benefit applies regardless of the gestational week.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
For additional diabetes screening benefits, also see the Diabetes Screening row. Also see the <i>Expanded Women's Preventive Health</i> section for Screening for Gestational Diabetes Mellitus and Screening for Diabetes Mellitus After Pregnancy .		
<p>Screening Mammography</p> <p><u>USPSTF Rating (2002): B</u> The USPSTF recommends screening mammography, with or without clinical breast examination (CBE), every 1-2 years for women aged 40 and older.</p> <p>Also see the Medical Policy titled Breast Imaging for Screening and Diagnosing Cancer.</p> <p>Also see the Breast Cancer Screening for Average-Risk Women recommendation in the <i>Expanded Women's Preventive Health</i> section.</p>	<p>Procedure Code(s): 77063, 77067</p> <p>Revenue Code: 0403</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>No age limits.</p> <p>Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>Note: This benefit only applies to screening mammography.</p>
<p>Cervical Cancer Screening</p> <p><u>USPSTF Rating (Aug. 2018): A</u> The USPSTF recommends screening for cervical cancer every 3 years with cervical cytology alone in women aged 21 to 29 years.</p> <p>For women aged 30 to 65 years, the USPSTF recommends:</p> <ul style="list-style-type: none"> • Screening every 3 years with cervical cytology alone, • Every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or • Every 5 years with hrHPV testing in combination with cytology (co-testing). <p><u>Bright Futures, March 2014:</u> Adolescents should no longer be routinely screened for cervical dysplasia until age 21.</p> <p>Also see Screening for Cervical Cancer in the <i>Expanded Women's</i></p>	<p><i>Human Papillomavirus DNA Testing (HPV):</i></p> <p>Procedure Code(s): 0500T, 87624, 87625, G0476</p> <p>Diagnosis Code(s): Z00.00, Z00.01, Z01.411, Z01.419, Z11.51, Z12.4</p>	<p><i>Human Papillomavirus DNA Testing (HPV):</i> Age 30 years and up. Requires one of the diagnosis codes listed in this row.</p>
	<p><i>Cervical Cytology (Pap Test)</i></p> <p>Code Group 1 Procedure Code(s): G0101, G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, Q0091, P3000, P3001</p> <p>Code Group 1 Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p>	<p><i>Cervical Cytology (Pap Test)</i></p> <p>Code Group 1: Limited to age 21–65 years (ends on 66th birthday). Does not have diagnosis code requirements for preventive benefits to apply.</p>
	<p><i>Cervical Cytology (Pap Test)</i></p> <p>Code Group 2 Procedure Code(s): 88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, ,</p>	<p><i>Cervical Cytology (Pap Test)</i></p> <p>Code Group 2: Limited to age 21–65 years (ends on 66th birthday). Requires one of the Code Group 2</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<i>Preventive Health section.</i>	88155, 88164, 88165, 88166, 88167, 88174, 88175 Code Group 2 Diagnosis Code(s): Z00.00, Z00.01, Z01.411, Z01.419, Z12.4	diagnosis codes listed in this row.
<p>Cholesterol Screening (Lipid Disorders Screening)</p> <p><u>USPSTF Rating (Nov. 2016): B</u> <i>Statin Use for the Primary Prevention of Cardiovascular Disease in Adults</i></p> <p>The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (i.e., symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met:</p> <ol style="list-style-type: none"> 1. They are aged 40 to 75 years; 2. They have 1 or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and 3. They have a calculated 10-year risk of a cardiovascular event of 10% or greater. <p>Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults aged 40 to 75 years.</p> <p>Notes:</p> <ul style="list-style-type: none"> • For statin medications benefits, refer to the pharmacy plan administrator. • See Dyslipidemia Screening (Bright Futures) for recommendations for children. 	<p>Procedure Code(s): <i>Cholesterol Screening:</i> 80061, 82465, 83718, 83719, 83721, 83722, 84478</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Z00.00, Z00.01, Z13.220</p>	<p><i>Cholesterol Screening:</i> Ages 40–75 years (ends on 76th birthday).</p> <p>Requires one of the diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Ages 40-75 years (ends on 76th birthday): Requires one of the listed Cholesterol Screening procedure codes AND one of the Diagnosis Codes listed in this row.</p> <p>Preventive Benefit Does Not Apply: For all ages above, if any of the following lipid disorders diagnosis codes are present in any position, the preventive benefit does not apply: E71.30, E75.5, E78.00, E78.01, E78.2, E78.3, E78.41, E78.49, E78.5, E78.79, E78.81, E78.89, E88.2, E88.89</p>
<p>Colorectal Cancer Screening</p> <p><u>USPSTF Rating (June 2016): A</u> The USPSTF recommends screening for colorectal cancer starting at age 50 years and continuing until age 75 years. The risks and benefits of different screening methods vary.</p>	<p>Fecal Occult Blood Testing (FOBT), Fecal Immunochemical Test (FIT), Fecal DNA, Sigmoidoscopy, or Colonoscopy</p> <p>Procedure Code(s):</p>	<p>Age Limits for Colorectal Cancer Screenings: 50-75 years (ends on 76th birthday).</p> <p>Fecal Occult Blood Testing (FOBT), Fecal Immunochemical Test (FIT), Fecal DNA, Sigmoidoscopy, or Colonoscopy:</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
	<p>Code Group 1: Sigmoidoscopy: G0104, G0106 Colonoscopy: G0105, G0120, G0121, G0122 FOBT and FIT: G0328 Colonoscopy Pre-op Consultation: S0285</p>	<p>Code Group 1: Does not have diagnosis code requirements for preventive benefits to apply.</p>
	<p>Code Group 2: Sigmoidoscopy: 45330, 45331, 45333, 45338, 45346 Colonoscopy: 44388, 44389, 44392, 44394, 45378, 45380, 45381, 45384, 45385, 45388 FOBT and FIT: 82270, 82274</p>	<p>Code Group 2: Requires one of the diagnosis codes listed in this row OR one of the procedure codes from Code Group 1, regardless of diagnosis.</p>
	<p>Code Group 3: Pathology: 88304, 88305</p> <p>Code Group 4: Anesthesia: 00812, 99152, 99153, 99156, 99157, G0500</p>	<p>Code Group 3 (Pathology) and Code Group 4 (Anesthesia): Requires one of the diagnosis codes listed in this row AND one of the procedure codes from Code Group 1 or Code Group 2.</p> <p>Code Group 3 and 4: Note: Preventive when performed for a colorectal cancer screening. Preventive benefits only apply when the surgeon's claim is preventive.</p>
	<p>Code Group 5: Pre-op/Consultation: 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241*, 99242*, 99243*, 99244*, 99245*</p> <p>*For additional information on the reimbursement of consultation codes 99241-99245, refer to the Reimbursement Policy titled Consultation Services Policy.</p>	<p>Code Group 5 : Requires one of the Code Group 5 diagnosis codes.</p>
	<p>Code Group 6: Fecal DNA: 81528</p> <p>Does not have diagnosis code requirements for preventive benefits to apply.</p>	<p>Code Group 6 (Fecal DNA): Benefit is limited to once every 3 years.</p> <p>Does not have diagnosis code requirements for preventive benefits to apply.</p>
	<p>Diagnosis Code(s): Code Groups 2, 3, and 4: Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.71, Z83.79</p> <p>Code Group 5: Z12.10, Z12.11, Z12.12, Z80.0, Z83.71, Z83.79</p>	

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
	<p>Computed Tomographic Colonography (Virtual Colonoscopy) Procedure Code(s): 74263</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p>	<p>Computed Tomographic Colonography (Virtual Colonoscopy) Does not have diagnosis code requirements for preventive benefit to apply.</p> <p>Prior authorization requirements may apply, depending on plan.</p>
<p>Wellness Examinations (well baby, well child, well adult)</p> <p><u>USPSTF Rating:</u> None UnitedHealthcare supports AAP and AAFP age and frequency guidelines.</p> <p><u>HRSA Requirements:</u> The Wellness Examinations codes include the following HRSA requirements for Women:</p> <ul style="list-style-type: none"> • Breastfeeding support and counseling • Contraceptive methods counseling and followup care • Domestic violence screening • Annual HIV counseling • Sexually transmitted infections counseling • Well-woman visits • Screening for urinary incontinence 	<p>Procedure Code(s): <i>Medicare wellness exams:</i> G0402, G0438, G0439 <i>STIs behavioral counseling:</i> G0445 <i>Annual gynecological exams:</i> S0610, S0612, S0613</p> <p><i>Preventive medicine services (evaluation and management):</i> 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397</p> <p><i>Preventive medicine, individual counseling:</i> 99401, 99402, 99403, 99404</p> <p><i>Preventive medicine, group counseling:</i> 99411, 99412</p> <p><i>Newborn Care (evaluation and management):</i> 99461</p> <p><i>Counseling Visit (to discuss the need for Lung Cancer Screening (LDCT) using Low Dose CT Scan):</i> G0296</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>Also see the Expanded Women's Preventive Health section</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>G0445 is limited to twice per year.</p> <p>G0296 is limited to age 55 to 80 years (ends on 81st birthday).</p>
<p>Vaccines (Immunizations)</p> <p><u>USPSTF Rating:</u> None An immunization that does not fall under one of the exclusions in the Certificate of Coverage is considered covered after the following conditions are satisfied:</p>	See the Preventive Vaccines (Immunizations) section	See the Preventive Vaccines (Immunizations) section

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>1. FDA approval; 2. Explicit ACIP recommendations for routine use published in the Morbidity & Mortality Weekly Report (MMWR) of the Centers for Disease Control and Prevention (CDC).</p> <p>Implementation will typically occur within 60 days after publication in the MMWR.</p>		
<p>Newborn Screenings All newborns</p> <p><u>USPSTF Rating (July 2008): B</u> Hearing Screening: Screening for hearing loss in all newborn infants.</p> <p><u>USPSTF Rating (March 2008): A</u> Hypothyroidism Screening: Screening for congenital hypothyroidism in newborns.</p> <p><u>USPSTF Rating (March 2008): A</u> Phenylketonuria Screening: Screening for phenylketonuria (PKU) in newborns.</p> <p><u>USPSTF Rating (Sept. 2007): A</u> Sickle Cell Screening: Screening for sickle cell disease in newborns.</p> <p>Note: For Bright Futures hearing screening, see Hearing Tests (Bright Futures).</p>	<p>Procedure Code(s): <i>Hearing Screening:</i> 92551, 92558, 92585, 92586, 92587, 92588, V5008</p> <p><i>Hypothyroidism Screening:</i> 84437, 84443</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p><i>Phenylketonuria Screening:</i> 84030, S3620</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p><i>Sickle Cell Screening:</i> 83020, 83021, 83030, 83033, 83051, S3850</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p><i>Newborn Screenings:</i> Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p><i>Blood Draw:</i> Age 0-90 days, requires one of the listed Hypothyroidism Screening, Phenylketonuria Screening, or Sickle Cell Screening procedure codes.</p>
<p>Metabolic Screening Panel (Newborns)</p>	<p>Procedure Code(s): <i>Metabolic Screening Panel:</i> 82017, 82136, 82261, 82775, 83020, 83498, 83516, 84030, 84437, 84443, S3620</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p><i>Metabolic Screening Panel:</i> Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p><i>Blood Draw:</i> Age 0-90 days. Requires one of the listed Metabolic Screening Panel procedure codes listed in this row.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>Osteoporosis Screening</p> <p><u>USPSTF Rating (June 2018): B</u> Women 65 and older: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women 65 years and older.</p> <p><u>USPSTF Rating (June 2018): B</u> Postmenopausal women younger than 65 years at increased risk of osteoporosis: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in postmenopausal women younger than 65 years who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.</p>	<p>Procedure Code(s): 76977, 77078, 77080, 77081 G0130</p> <p>Diagnosis Code(s): Z00.00, Z00.01, Z13.820, Z82.62</p>	Requires one of the diagnosis codes listed in this row.
<p>Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults</p> <p><u>USPSTF Rating (Nov. 2018): B</u> The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use.</p> <p><u>Bright Futures (April 2017):</u> Bright Futures recommends alcohol or drug use assessments from age 11-21 years.</p>	<p>Procedure Code(s): <i>Alcohol or Drug Use Screening:</i> 99408, 99409</p> <p><i>Annual Alcohol Screening: G0442</i></p> <p><i>Brief Counseling for Alcohol: G0443</i></p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p>	Does not have diagnosis code requirements for preventive benefits to apply.
<p>High Blood Pressure in Adults – Screening</p> <p><u>USPSTF Rating (Oct. 2015):A</u> The USPSTF recommends screening for high blood pressure in adults aged 18 years or older. The USPSTF recommends</p>	<p>Blood Pressure Measurement in a Clinical Setting: N/A</p>	<p>Blood Pressure Measurement in a Clinical Setting: This service is included in a preventive care wellness examination.</p>
	<p>Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting):</p>	<p>Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting):</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	Procedure Code(s): <i>Ambulatory Blood Pressure Measurement:</i> 93784, 93786, 93788 or 93790 Diagnosis Code(s): <i>Abnormal Blood-Pressure Reading Without Diagnosis of Hypertension:</i> R03.0	Age 18 years and up. Requires the diagnosis code listed in this row.
Breast Cancer: Medications for Risk Reduction <u>USPSTF Rating (Sept. 2013): B</u> The USPSTF recommends that clinicians engage in shared, informed decision making with women who are at increased risk for breast cancer about medications to reduce their risk. For women who are at increased risk for breast cancer and at low risk for adverse medication effects, clinicians should offer to prescribe risk-reducing medications, such as tamoxifen or raloxifene.	Procedure Code(s): <i>Evaluation and Management (Office Visits):</i> 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397, G0463 Diagnosis Code(s): Z80.3, Z80.41, Z15.01, Z15.02	Requires one of the diagnosis codes listed in this row in the primary position.
Primary Care Interventions to Promote Breastfeeding <u>USPSTF Rating (Oct. 2016): B</u> The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	N/A Also see the Expanded Women's Preventive Health section	Included in primary care or OB/GYN office visits
Screening for Depression in Adults <u>USPSTF Rating (Jan. 2016): B</u> Recommendation: The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	Procedure Code(s): 96127, G0444 Diagnosis Code(s): Required for 96127 Only: <i>Encounter for screening for depression:</i> Z13.31, Z13.32	Requires one of the diagnosis code listed in this row, for 96127. The diagnosis codes listed in this row are not required, for G0444.

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>Depression in Children and Adolescents (Screening)</p> <p><u>USPSTF Rating (Feb. 2016): B</u> The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.</p> <p>Note: The Bright Futures Periodicity Schedule recommends depression screening begin at age 12 through 21 years.</p>	<p>Procedure Code(s): 96127, G0444</p> <p>Diagnosis Code(s): Required for 96127 Only: <i>Encounter for screening for depression: Z13.31, Z13.32</i></p>	<p>Requires one of the diagnosis codes listed in this row, for 96127.</p> <p>The diagnosis codes listed in this row are not required for G0444.</p>
<p>Behavioral Counseling in Primary Care to Promote a Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors</p> <p><u>USPSTF Rating (Aug. 2014): B</u> The USPSTF recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.</p>	<p>Procedure Code(s): <i>Medical Nutrition Therapy or Counseling:</i> 97802, 97803, 97804, G0270, G0271, S9470</p> <p><i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404</p> <p><i>Behavioral Counseling or Therapy:</i> 0403T, G0446, G0447, G0473</p> <p>Diagnosis Code(s): Screening: Z13.220</p> <p><i>History:</i> F17.210, F17.211, F17.213, F17.218, F17.219, Z72.0, Z87.891, Z82.49</p> <p><i>Overweight:</i> E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29</p> <p><i>Body Mass Index 30.0 – 39.9:</i> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39</p> <p><i>Body Mass Index 40.0 and Over:</i> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p> <p><i>Impaired Fasting Glucose: R73.01</i></p>	<p>Requires one of the diagnosis codes listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and S9470.</p> <p>The diagnosis code listed in this row are not required for G0446, G0447, and G0473.</p> <p>G0446 is limited to once per year.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
	<p><i>Metabolic Syndrome: E88.81</i></p> <p><i>Hyperlipidemia / Dyslipidemia: E78.00, E78.01, E78.1, E78.2, E78.3, E78.41, E78.49, E78.5</i></p> <p><i>Obesity: E66.01, E66.09, E66.1, E66.8, E66.9, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</i></p> <p><i>Essential Hypertension: I10</i></p> <p><i>Secondary Hypertension: I15.0, I15.1, I15.2, I15.8, I15.9, N26.2</i></p> <p><i>Hypertension Complicating Pregnancy, Childbirth and the Puerperium: O10.011, O10.012, O10.013, O10.019, O10.02, O10.03, O10.111, O10.112, O10.113, O10.119, O10.12, O10.13, O10.211, O10.212, O10.213, O10.219, O10.22, O10.23, O10.311, O10.312, O10.313, O10.319, O10.32, O10.33, O10.411, O10.412, O10.413, O10.419, O10.42, O10.43, O10.911, O10.912, O10.913, O10.919, O10.92, O10.93, O11.1, O11.2, O11.3, O11.4, O11.5, O11.9, O13.1, O13.2, O13.3, O13.4, O13.5, O13.9, O16.1, O16.2, O16.3, O16.4, O16.5, O16.9</i></p> <p><i>Urgent/Emergency/Crisis Hypertension: I16.0, I16.1, I16.9</i></p> <p><i>Diabetes: Diabetes Diagnosis Code List</i></p> <p><i>Atherosclerosis: Atherosclerosis Diagnosis Code List</i></p> <p><i>Coronary Atherosclerosis: I25.10, I25.110, I25.111, I25.118, I25.119, I25.700, I25.701, I25.708, I25.709, I25.710, I25.711, I25.718, I25.719, I25.720, I25.721, I25.728, I25.729, I25.730, I25.731, I25.738, I25.739, I25.750, I25.751, I25.758, I25.759, I25.760, I25.761, I25.768, I25.769, I25.790, I25.791, I25.798, I25.799, I25.810, I25.811, I25.812</i></p>	

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>Weight Loss to Prevent Obesity-Related Morbidity and Mortality in Adults: Behavioral Interventions</p> <p><u>USPSTF Rating (Sept. 2018): B</u> The USPSTF recommends that clinicians offer or refer adults with a body mass index (BMI) of 30 or higher (calculated as weight in kilograms divided by height in meters squared) to intensive multicomponent behavioral interventions.</p>	<p>Procedure Code(s): <i>Medical Nutrition Therapy:</i> 97802, 97803, 97804</p> <p><i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404</p> <p><i>Behavioral Counseling or Therapy:</i> G0446, G0447, G0473</p> <p>Diagnosis Code(s): <i>Body Mass Index 30.0 – 39.9:</i> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39</p> <p><i>Body Mass Index 40.0 and over:</i> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p> <p><i>Obesity:</i> E66.01, E66.09, E66.1, E66.8, E66.9</p>	<p>G0446 is limited to once per year.</p> <p>Requires one of the diagnosis codes listed in this row for 97802-97804 and 99401-99404.</p> <p>The diagnosis codes listed in this row are not required for G0446, G0447 and G0473.</p>
<p>Screening for Obesity in Children and Adolescents</p> <p><u>USPSTF Rating (June 2017): B</u> The USPSTF recommends that clinicians screen for obesity in children and adolescents 6 years and older and offer or refer them to comprehensive, intensive behavioral interventions to promote improvements in weight status.</p>	<p>Procedure Code(s): <i>Medical Nutrition Therapy:</i> 97802, 97803, 97804</p> <p><i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404</p> <p><i>Behavioral Counseling or Therapy:</i> G0446, G0447, G0473</p> <p>Also see the codes in the Wellness Examinations row above.</p> <p>Diagnosis Code(s): <i>Obesity:</i> E66.01, E66.09, E66.1, E66.8, E66.9</p>	<p>G0446 is limited to once per year.</p> <p>Requires one of the diagnosis codes listed in this row for 97802-97804 and 99401-99404.</p> <p>The diagnosis codes listed in this row are not required for G0446, G0447, and G0473.</p>
<p>Behavioral Counseling to Prevent Sexually Transmitted Infections</p> <p><u>USPSTF Rating (Sept. 2014): B</u> The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs).</p>	<p>Procedure Code(s): 99401, 99402, 99403, 99404, G0445</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>G0445 is limited to twice per year.</p> <p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>Tobacco Smoking Cessation in Adults, including Pregnant Women: Behavioral and Pharmacotherapy Interventions</p> <p><u>USPSTF Rating (Sept. 2015): A</u> The USPSTF recommends that clinicians ask all pregnant women about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant women who use tobacco.</p> <p>The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)-approved pharmacotherapy for cessation to adults who use tobacco.</p>	<p>Procedure Code(s): <i>Behavioral Interventions:</i> 99406, 99407 99401, 99402, 99403, 99404</p> <p>Also see the codes in the Wellness Examinations row above.</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>
<p>Primary Care Interventions To Prevent Tobacco Use In Children And Adolescents</p> <p><u>USPSTF Rating (Aug. 2013): B</u> The USPSTF recommends that primary care clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use among school-aged children and adolescents.</p> <p><u>Bright Futures (April 2017):</u> Bright Futures recommends tobacco use assessments from age 11 years – 21 years.</p>	<p>Procedure Code(s): <i>Smoking and Tobacco Use Cessation Counseling Visit:</i> 99406, 99407, 99401, 99402, 99403, 99404</p> <p>Also see the codes in the Wellness Examinations row above.</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>
<p>Screening for Visual Impairment in Children</p> <p><u>USPSTF Rating (Sept. 2017): B</u> The USPSTF recommends vision screening at least once in all children aged 3 to 5 years to detect amblyopia or its risk factors.</p> <p><u>Bright Futures:</u> Instrument-based screening recommended for:</p> <ul style="list-style-type: none"> • Age 1–5 years: if the 	<p>Procedure Code(s): 99173, 99174, 99177</p> <p>Diagnosis Code(s): See the Preventive Benefit Instructions.</p>	<p>Age Limit (99173, 99174 and 99177): Up to age 21 years (ends on 22nd birthday).</p> <p>Code 99173: Does not have diagnosis code requirements for preventive benefits to apply.</p> <p>Code 99174 and 99177: See the Medical Policy titled Omnibus Codes for allowable diagnoses.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
screening is available. <ul style="list-style-type: none"> Age 6 years and up: if unable to test visual acuity monocularly with age appropriate optotypes. 		
Behavioral Counseling to Prevent Skin Cancer <u>USPSTF Rating (March 2018): B</u> The USPSTF recommends counseling young adults, adolescents, children and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons ages 6 months to 24 years with fair skin types to reduce their risk of skin cancer.	N/A	This service is included in a preventive care wellness examination or focused E&M visit.
Prevention of Falls in Community-Dwelling Older Adults <u>USPSTF Rating (April 2018): B</u> The USPSTF recommends exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls.	N/A	This service is included in a preventive care wellness examination or focused E&M visit.
Screening for Intimate Partner Violence <u>USPSTF Rating (Oct. 2018): B</u> The USPSTF recommends that clinicians screen for intimate partner violence in women of reproductive age and provide or refer women who screen positive to ongoing support services. Also see Screening and Counseling for Interpersonal and Domestic Violence in the <i>Expanded Women's Preventive Health</i> section.	N/A	This service is included in a preventive care wellness examination.
Screening for Lung Cancer with Low-Dose Computed Tomography <u>USPSTF Rating (Dec. 2013): B</u> The USPSTF recommends annual screening for lung cancer with low-dose computed tomography in adults ages 55 to 80 years who have a 30 pack-year smoking	Procedure Code(s): G0297 Diagnosis Code(s): F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891 Codes for Reporting Purposes: G9275, G9276, G9458, G9459, G9460	Requires one of the diagnosis codes listed in this row. Limitations: <ul style="list-style-type: none"> Limited to one per year, AND All of the following criteria: <ul style="list-style-type: none"> Age 55 to 80 years (ends on 81st birthday), and At least 30 pack-years* of smoking history, and

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.	Note: Codes G9275, G9276, G9458, G9459, and G9460 are for reporting purposes only, if applicable. These codes are not separately reimbursable.	<ul style="list-style-type: none"> ○ Either a current smoker, or, have quit within the past 15 years <p>Note: Prior authorization requirements may apply, depending on plan.</p> <p>*A pack-year is a way to measure the amount a person has smoked over a long period of time. It is calculated by multiplying the number of packs of cigarettes smoked per day by the number of years the person has smoked. For example, 1 pack year is equal to smoking 1 pack per day for 1 year, or 2 packs per day for half a year, and so on. Source: National Institutes of Health, National Cancer Institute Dictionary of Cancer Terms, pack year definition web page. http://www.cancer.gov/dictionary?Cd=306510</p>
<p>Fluoride Application in Primary Care</p> <p><u>USPSTF Rating (May 2014): B</u> Children From Birth Through Age 5 Years. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption.</p> <p><u>Bright Futures (April 2017):</u> For those at high risk, consider application of fluoride varnish for caries prevention every 3 to 6 months between ages 6 months to 5 years.</p>	<p>Procedure Code(s): <i>Application of Topical Fluoride by Physician or Other Qualified Health Care Professional: 99188</i></p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Age 0-5years (ends on 6th birthday).</p> <p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>
<p>Latent Tuberculosis Infection: Screening, Adults</p> <p><u>USPSTF Rating (Sept. 2016): B</u> The USPSTF recommends screening for latent tuberculosis infection (LTBI) in populations at increased risk. This recommendation applies to asymptomatic adults 18 years and older at increased risk for tuberculosis.</p>	<p>Procedure Code(s): <i>Screening:</i> 86480, 86481, 86580 <i>Followup Visit to Check Results:</i> 99211 <i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): R76.11, R76.12, Z00.00, Z00.01, Z11.1, Z11.7, Z20.1</p>	<p><i>Screening:</i> Ages 18 years and up.</p> <p>Requires one of the diagnosis codes listed in this row for CPT code 86480, 86481, and 86580.</p> <p><i>Followup Visit to Check Results (99211):</i> CPT code 99211 requires diagnosis code R76.11 or R76.12.</p> <p><i>Blood Draw:</i> Ages 18 years and up.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
	Note for age 18-21 years (ends on 22nd birthday): In addition to the codes in this row, the preventive benefit also applies to the diagnosis codes listed in the Bright Futures row: Tuberculosis (TB) Testing	Required to be billed with 86480 or 86481 AND one of the diagnosis codes listed in this row.
Preeclampsia Screening <u>USPSTF Rating (April 2017): B</u> The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy.	Preeclampsia screening by blood pressure measurement is included in the code for a prenatal care office visit. See the following code groups in the <i>Expanded Women's Preventive Health</i> section: <ul style="list-style-type: none"> • Prenatal Office Visits • Prenatal Care Visits • Global Obstetrical Codes 	See the following code groups in the <i>Expanded Women's Preventive Health</i> section: <ul style="list-style-type: none"> • Prenatal Office Visits • Prenatal Care Visits • Global Obstetrical Codes
Bright Futures		
Anemia Screening in Children (Bright Futures)	Procedure Code(s): <i>Anemia Screening in Children:</i> 85014, 85018 <i>Blood Draw:</i> 36415, 36416 Diagnosis Code(s): Z00.110, Z00.111, Z00.121, Z00.129, Z13.0	<i>Anemia Screening in Children:</i> Ages prenatal to 21 (ends on 22 nd birthday). No frequency limit. Requires one of the diagnosis codes listed in this row. <i>Blood Draw:</i> Ages prenatal to 21 (ends on 22 nd birthday). Required to be billed with 85014 or 85018 AND one of the diagnosis codes listed in this row.
Hearing Tests <u>Bright Futures (April 2017):</u> <i>Hearing Tests:</i> Recommended at ages: Newborn; Between 3-5 days to 2 months; 4 years; 5 years, 6 years; 8 years; 10 years; Once between age 11 – 14 years; Once between age 15 – 17 years; Once between age 18 – 21 years; Also recommended for those that have a positive risk assessment. <i>Risk Assessment:</i> Recommended at ages: 4 mo, 6 mo, 9 mo, 12 mo, 15 mo, 18 mo, 24 mo, 30 mo, 3 years, 7 years, and 9 years.	Procedure Code(s): <i>Hearing Tests:</i> 92551, 92552, 92553 Diagnosis Code(s): Z00.121, Z00.129, Z01.10 Note: A risk assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above	Ages prenatal to 21 (ends on 22 nd birthday). Limit of once per year. Requires one of the diagnosis codes listed in this row.
Screening for Visual Impairment in Children (Bright Futures)	See row above for Screening for Visual Impairment in Children	See row above Screening for Visual Impairment in Children .

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>Formal Developmental / Autism Screening</p> <p><u>Bright Futures:</u></p> <ul style="list-style-type: none"> • A formal, standardized developmental screen is recommended during the 9 month visit. • A formal, standardized developmental screen is recommended during the 18 month visit, including a formal autism screen. • A formal, standardized autism screen is recommended during the 24 month visit. • A formal, standardized developmental screen is recommended during the 30 month visit. 	<p>Procedure Code(s): 96110</p> <p>Diagnosis Code(s): Z00.121, Z00.129, Z13.40, Z13.41, Z13.42, Z13.49</p>	<p>Ages prenatal to 2 years (ends on 3rd birthday).</p> <p>No frequency limit.</p> <p>Requires one of the diagnosis codes listed in this row.</p>
<p>Lead Screening</p> <p><u>Bright Futures:</u> <i>Screening Lab Work:</i> Conduct risk assessment or screening, as appropriate, at the following intervals: 12 mo and 24 mo.</p> <p><i>Risk Assessment, and Screening if positive:</i> Recommended at 6 mo, 9 mo, 12 mo, 18 mo, 24 mo, 3 years, 4 years, 5 years and 6 years.</p>	<p>Procedure Code(s): <i>Lead Screening:</i> 83655</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Z00.121, Z00.129, Z77.011</p>	<p><i>Lead Screening:</i> Ages 6 months through age 6 years (ends on 7th birthday). No frequency limit.</p> <p>Requires one of the diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Ages 6 months through age 6 years (ends on 7th birthday).</p> <p>Required to be billed with 83655 AND one of the diagnosis codes in this row.</p>
<p>Tuberculosis (TB) Testing</p> <p><u>Bright Futures</u> For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults.</p>	<p>Procedure Code(s): <i>Screening:</i> 86580</p> <p><i>Followup visit to check results:</i> 99211</p> <p>Diagnosis Code(s): R76.11, R76.12, Z20.1, Z00.121, Z00.129, Z11.1, Z11.7</p> <p>Note: For age 18 years and older: In addition to these codes, the preventive benefit also applies to all codes listed in the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults</p>	<p>Ages prenatal to 21(ends on 22nd birthday).</p> <p>Note: For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults</p> <p>No frequency limit.</p> <p>CPT code 86580 requires one of the diagnosis codes listed in this row.</p> <p>CPT code 99211 requires diagnosis code R76.11, R76.12, or Z11.1.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health section](#).
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
<p>Dyslipidemia Screening</p> <p><u>Bright Futures (April 2014):</u> <i>Risk Assessment:</i> Recommended at 24 mo, 4 years, 6 years, 8 years, 12 years, 13 years, 14 years, 15 years, 16 years.</p> <p><i>Screening Lab Work:</i> Conduct if risk assessment is positive, or, at the following intervals: once between age 9 – 11 years; once between age 17 – 21 years</p>	<p>Procedure Code(s): <i>Dyslipidemia Screening Lab Work:</i> 80061, 82465, 83718, 83719, 83721, 83722, 84478</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Z00.121, Z00.129, Z13.220</p> <p>Note: A risk assessment is included in the code for a wellness examination visit; see the Wellness Examinations row above.</p>	<p><i>Dyslipidemia Screening Lab Work:</i> Ages 24 months to 21 years (ends on 22nd birthday). Requires one of the diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Ages 24 months to 21 years (ends on 22nd birthday).</p> <p>Requires one of the listed Dyslipidemia Screening procedure codes listed in this row AND one of the diagnosis codes listed in this row.</p>
<p>Tobacco, Alcohol or Drug Use Assessment</p> <p><u>Bright Futures (April 2017):</u> Bright Futures recommends tobacco, alcohol or drug use assessment from age 11 years – 21 years.</p>	<p>See codes in the rows above:</p> <ul style="list-style-type: none"> • Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents • Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults 	<p>See the rows above:</p> <ul style="list-style-type: none"> • Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents • Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults
<p>Psychosocial / Behavioral Assessment</p> <p><u>Bright Futures (April 2017):</u> Bright Futures recommends physicians conduct psychosocial / behavioral assessment at each of the recommended visits between newborn – 21 years.</p>	<p>An assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.</p>	<p>See the Wellness Examinations row above.</p>
<p>Depression Screening</p> <p><u>Bright Futures (April 2017):</u> Bright Futures recommends depression screening at each of the recommended visits between age 12-21 years.</p>	<p>See the codes in the Depression in Children and Adolescents (Screening) row above.</p>	<p>See the Depression in Children and Adolescents (Screening) row above.</p>
<p>Sexually Transmitted Infections (STI)</p> <p><u>Bright Futures (April 2017):</u> Bright Futures recommends the following: <i>STI Risk Assessment:</i> Conduct risk assessment at each of the recommended visits between 11 years – 21 years. <i>STI Lab Work:</i> Conduct if risk assessment is positive.</p>	<p>See the codes in the Chlamydia Infection Screening and Gonorrhea Screening rows above.</p>	<p>See the Chlamydia Infection Screening and Gonorrhea Screening rows above.</p>

Preventive Care Services

*Also see the [Expanded Women's Preventive Health](#) section.
Certain codes may not be payable in all circumstances due to other policies or guidelines.
For preventive care medications, refer to the pharmacy plan administrator.*

Service: <i>A date in this column is when the listed rating was released, not when the benefit is effective.</i>	Code(s):	Preventive Benefit Instructions:
HIV Screening <u>Bright Futures (April 2017):</u> <i>HIV Risk Assessment:</i> Conduct risk assessment at age 11 years, 12 years, 13 years, 14 years, 19 years, 20 years and 21 years. <i>HIV Screening Lab Work:</i> Conduct once between age 15-18 years. Also recommended anytime between ages 11-14 years, and 19-21 years when a risk assessment is positive.	See the codes in the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row above.	See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row above.

PREVENTIVE VACCINES (IMMUNIZATIONS)

A vaccine (immunization) that does not fall under one of the exclusions in the Certificate of Coverage is considered covered after the following conditions are satisfied: (1) FDA approval; (2) explicit ACIP recommendations for routine use published in the Morbidity & Mortality Weekly Report (MMWR) of the Centers for Disease Control and Prevention (CDC). Implementation will typically occur within 60 days after publication in the MMWR. In the case of a public health emergency (as defined by the Centers for Disease Control or state or local public health departments) UnitedHealthcare may choose to apply preventive benefits to a new vaccine if the vaccine has FDA approval, even if an ACIP recommendation has not been announced.

Notes:

- **Trade Name(s) column:** Brand names/trade names are included, when available, as examples for convenience only. Coverage pursuant to this Coverage Determination Guideline is based solely on the procedure codes.
- **Age Group column:** This column is provided for informational use only. For purposes of this document: Adult means age 18 years and up; Pediatric means age 0-18 years.
- **Benefit Limits column:** Benefit Limits in **bold text** are from FDA labeling and ACIP recommendations. Codes that indicate "For applicable age see code description" are limited to the age(s) listed in the code description.

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
Immunization Administration Preventive when included as part of a preventive immunization.	90460	Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional; first or only component of each vaccine or toxoid administered	N/A	Pediatric	For applicable age see code description.

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
	90461	Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional; each additional vaccine or toxoid component administered (List separately in addition to code for primary procedure)	N/A	Pediatric	For applicable age see code description.
	90471	Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); one vaccine (single or combination vaccine/toxoid)	N/A	Both	-
	90472	Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure)	N/A	Both	-
	90473	Immunization administration by intranasal or oral route; one vaccine (single or combination vaccine/toxoid)	N/A	Both	-
	90474	Immunization administration by intranasal or oral route; each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure)	N/A	Both	-
	G0008	Administration of influenza virus vaccine	N/A	Both	-
	G0009	Administration of pneumococcal vaccine	N/A	Both	-
	G0010	Administration of hepatitis B vaccine	N/A	Both	-
	0771 (revenue code)	Vaccine administration	N/A	Both	-
Meningococcal (MenB; MenB-4C; MenB-FHbp; Hib-MenCY; MPSV4; MCV4; MenACWY-CRM)	90620	Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B (MenB-4C), 2 dose schedule, for intramuscular use	Bexsero®	Both	Benefit Limit: Age 10 and up
	90621	Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2 or 3 dose schedule, for intramuscular use	Trumenba®	Both	Benefit Limit: Age 10 and up

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
	90644	Meningococcal conjugate vaccine, serogroups C & Y and Haemophilus influenzae b vaccine (Hib-MenCY), 4 dose schedule, when administered to children 2-15 months of age , for intramuscular use	MenHibrix®	Pediatric	For applicable age see code description.
	90733	Meningococcal polysaccharide vaccine, serogroups A, C, Y, W-135, quadrivalent (MPSV4) for subcutaneous use	Menomune®	Both	-
	90734	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent diphtheria toxoid carrier (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for intramuscular use	Menactra® Menveo®	Both	-
Hepatitis A	90632	Hepatitis A vaccine (HepA), adult dosage , for intramuscular use	Havrix® VAQTA®	Adult	For applicable age see code description.
	90633	Hepatitis A vaccine (HepA), pediatric/adolescent dosage -2 dose schedule, for intramuscular use	Havrix® VAQTA®	Pediatric	For applicable age see code description.
	90634	Hepatitis A vaccine (HepA), pediatric/adolescent dosage -3 dose schedule, for intramuscular use	Havrix®	Pediatric	For applicable age see code description.
	90636	Hepatitis A and hepatitis B vaccine (HepA-HepB), adult dosage , for intramuscular use	Twinrix®	Adult	For applicable age see code description.
Haemophilus influenzae b (Hib)	90647	Haemophilus influenzae b vaccine (Hib), PRP-OMP conjugate, 3 dose schedule, for intramuscular use	PedvaxHIB®	Both	-
	90648	Haemophilus influenzae b vaccine (Hib), PRP-T conjugate, 4 dose schedule, for intramuscular use	ActHIB® Hiberix®	Both	-
Human Papilloma Virus (HPV)	90649	Human Papilloma virus vaccine, types 6, 11, 16, 18, quadrivalent (HPV4), 3 dose schedule, for intramuscular use	Gardasil4®	Both	Benefit Limit: Ages 9-26 years (ends on 27th birthday)
	90650	Human Papilloma virus vaccine, types 16, 18, bivalent (HPV2), 3 dose schedule, for intramuscular use	N/A	Both	Benefit Limit: Ages 9-26 years (ends on 27th birthday)
	90651	Human Papillomavirus vaccine types 6, 11, 16, 18, 31, 33, 45, 52, 58, nonavalent (9vHPV), 2 or 3 dose schedule, for intramuscular use	Gardasil9®	Both	Benefit Limit: Ages 9-45 years (ends on 46th birthday)

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
Seasonal Influenza ('flu') <i>Note: Additional new seasonal flu immunization codes that are recently FDA-approved, but are not listed here, may be eligible for preventive benefits as of the FDA approval date.</i>	90630	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, for intradermal use	Fluzone® Intradermal Quadrivalent	Both	-
	90653	Influenza vaccine, inactivated (IIV), subunit, adjuvanted, for intramuscular use	Fluad®	Both	-
	90654	Influenza virus vaccine, trivalent (IIV3), split virus, preservative-free, for intradermal use	Fluzone® Intradermal Trivalent	Adult	Benefit Limit: 18-64 years (ends on 65th birthday)
	90655	Influenza virus vaccine, trivalent (IIV3), split virus, preservative free, 0.25 mL dosage, for intramuscular use	Fluzone® No Preservative Pediatric	Pediatric	Benefit Limit: 6-35 months old
	90656	Influenza virus vaccine, trivalent (IIV3), split virus, preservative free, 0.5 mL dosage, for intramuscular use	Afluria® Fluzone® No preservative Fluvirin® Fluarix® Flulaval®	Both	Benefit Limit: 3 years and up
	90657	Influenza virus vaccine, trivalent (IIV3), split virus, 0.25 mL dosage, for intramuscular use	Fluzone®	Pediatric	Benefit Limit: 6-35 months old
	90658	Influenza virus vaccine, trivalent (IIV3), split virus, 0.5 mL dosage, for intramuscular use	Afluria® Flulaval® Fluvirin® Fluzone®	Both	Benefit Limit: 3 years and up
	90660	Influenza virus vaccine, trivalent, live (LAIV3), for intranasal use	Flumist®	Both	Benefit Limit: Ages 2-49 years (ends on 50th birthday)
	90661	Influenza virus vaccine, trivalent (ccIIV3), derived from cell cultures, subunit, preservative and antibiotic free, 0.5 mL dosage, for intramuscular use	Flucelvax™	Adult	Benefit Limit: Ages 4 years and up
	90662	Influenza virus vaccine (IIV), split virus, preservative free, enhanced immunogenicity via increased antigen content, for intramuscular use	High Dose Fluzone®	Adult	Benefit Limit: Ages 65 years and up
90664	Influenza virus vaccine, live (LAIV), pandemic formulation, for intranasal use	Flumist®	Both	Benefit Limit: Ages 2-49 years (ends on 50th birthday)	
90666	Influenza virus vaccine (IIV), pandemic formulation, split virus, preservative free, for intramuscular use	N/A	Both	-	

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
	90667	Influenza virus vaccine (IIV), pandemic formulation, split virus, adjuvanted, for intramuscular use	N/A	Both	-
	90668	Influenza virus vaccine (IIV), pandemic formulation, split virus, for intramuscular use	N/A	Both	-
	90672	Influenza virus vaccine, quadrivalent, live (LAIV4), for intranasal use	Flumist® (LAIV4)	Both	Benefit Limit: Ages 2-49 years (ends on 50th birthday)
	90673	Influenza virus vaccine, trivalent (RIV3), derived from recombinant DNA (RIV3), hemagglutinin (HA) protein only, preservative and antibiotic free, for intramuscular use	Flublok®	Adult	Benefit Limit: Age 18 years and up
	90674	Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit, preservative and antibiotic free, 0.5 mL dosage, for intramuscular use	Flucelvax® Quadrivalent	Both	Benefit Limit: Age 4 years and up
	90682	Influenza virus vaccine, quadrivalent (RIV4), derived from recombinant DNA, hemagglutinin (HA) protein only, preservative and antibiotic free, for intramuscular use	Flublok Quadrivalent®	Adult	Benefit Limit: Age 18 years and up
	90685	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.25 mL, for intramuscular use	Fluzone Quadrivalent®	Pediatric	Benefit Limit: 6-35 months old
	90686	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.5 mL dosage, for intramuscular use	Afluria® Quadrivalent Fluarix® FluLaval Quadrivalent® Fluzone Quadrivalent®	Both	Benefit Limit: Ages 6 months and up
	90687	Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.25 mL dosage, for intramuscular use	Fluzone Quadrivalent®	Pediatric	Benefit Limit: 6-35 months old
	90688	Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.5 mL dosage, for intramuscular use	Afluria® Quadrivalent FluLaval Quadrivalent® Fluzone Quadrivalent®	Both	Benefit Limit: Ages 6 months and up

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
	90689	Influenza virus vaccine quadrivalent (IIV4), inactivated, adjuvanted, preservative free, 0.25mL dosage, for intramuscular use	-	Both	-
	90756	Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit, antibiotic free, 0.5mL dosage, for intramuscular use	Flucelvax Quadrivalent® (non-preservative free)	Both	-
	Q2034	Influenza virus vaccine, split virus, for intramuscular use (Agriflu)	Agriflu®	Adult	Benefit Limit: Ages 18 years and up
	Q2035	Influenza virus vaccine, split virus, when administered to individuals 3 years of age and older , for intramuscular use (AFLURIA)	Afluria®	Both	For applicable age see code description.
	Q2036	Influenza virus vaccine, split virus, when administered to individuals 3 years of age and older , for intramuscular use (FLULAVAL)	Flulaval®	Both	For applicable age see code description.
	Q2037	Influenza virus vaccine, split virus, when administered to individuals 3 years of age and older , for intramuscular use (FLUVIRIN)	Fluvirin®	Both	For applicable age see code description.
	Q2038	Influenza virus vaccine, split virus, when administered to individuals 3 years of age and older , for intramuscular use (Fluzone)	Fluzone®	Both	For applicable age see code description.
	Q2039	Influenza virus vaccine, not otherwise specified	N/A	Both	-
Pneumococcal polysaccharide (PPSV23)	90732	Pneumococcal polysaccharide vaccine, 23-valent (PPSV23), adult or immunosuppressed patient dosage, when administered to individuals 2 years or older , for subcutaneous or intramuscular use	Pneumovax 23®	Both	For applicable age see code description.
Pneumococcal conjugate	90670	Pneumococcal conjugate vaccine, 13 valent (PCV13), for intramuscular use	Prevnar 13® (PCV13)	Both	-
Rotavirus (RV1, RV5)	90680	Rotavirus vaccine, pentavalent (RV5), 3 dose schedule, live, for oral use	Rotateq®	Pediatric	Benefit Limit: 0-8 months old
	90681	Rotavirus vaccine, human, attenuated (RV1), 2 dose schedule, live, for oral use	Rotarix®	Pediatric	Benefit Limit: 0-8 months old

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
Diphtheria, tetanus toxoids, acellular pertussis and polio inactive (DTap-IPV)	90696	Diphtheria, tetanus toxoids, acellular pertussis vaccine and inactivated poliovirus vaccine (DTaP-IPV), when administered to children 4 through 6 years of age , for intramuscular use	Kinrix® Quadracel®	Pediatric	For applicable age see code description.
Diphtheria, tetanus toxoids, acellular pertussis, haemophilus influenza B, and polio inactive (DTap-IPV/Hib)	90698	Diphtheria, tetanus toxoids, acellular pertussis vaccine, Haemophilus influenzae type b, and inactivated poliovirus vaccine, (DTaP-IPV/Hib), for intramuscular use	Pentacel®	Pediatric	Benefit Limit: Ages 0-4 years (ends on 5th birthday)
Diphtheria, tetanus, acellular pertussis (DTap)	90700	Diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP), when administered to individuals younger than 7 years , for intramuscular use	Daptacel® Infanrix®	Pediatric	For applicable age see code description.
Diphtheria and tetanus (DT)	90702	Diphtheria and tetanus toxoids adsorbed (DT) when administered to individuals younger than 7 years , for intramuscular use	N/A	Pediatric	For applicable age see code description.
Measles, Mumps, Rubella (MMR)	90707	Measles, mumps and rubella virus vaccine (MMR), live, for subcutaneous use	MMR II®	Both	-
	90710	Measles, mumps, rubella, and varicella vaccine (MMRV), live, for subcutaneous use	ProQuad®	Pediatric	Benefit Limit: Ages 1-12 years (ends on 13th birthday)
Polio (IPV)	90713	Poliovirus vaccine, inactivated (IPV), for subcutaneous or intramuscular use	Ipol®	Both	-
Tetanus and diphtheria (Td)	90714	Tetanus and diphtheria toxoids adsorbed (Td), preservative free, when administered to individuals 7 years or older , for intramuscular use	Tenivac® Decavac®	Both	For applicable age see code description.
Tetanus, diphtheria toxoids and acellular pertussis (Tdap)	90715	Tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap), when administered to individuals 7 years or older , for intramuscular use	Adacel® Boostrix®	Both	For applicable age see code description.
Varicella (VAR) ('chicken pox')	90716	Varicella virus vaccine (VAR), live, for subcutaneous use	Varivax®	Both	-
Diphtheria, tetanus and acellular pertussis, hep B, and polio inactive (DTaP-HepB-IPV)	90723	Diphtheria, tetanus toxoids, acellular pertussis vaccine, hepatitis B, and inactivated poliovirus vaccine (DTaP-HepB-IPV), for intramuscular use	Pediarix®	Both	Benefit Limit: Ages 0-6 years (ends on 7th birthday)

PREVENTIVE VACCINES (IMMUNIZATIONS)

These codes do **not** have a diagnosis code requirement for preventive benefits to apply.

Category:	Code(s):	Description:	Trade Name(s) (See Note above)	Age Group: (Pediatric, Adult, or Both)	Benefit Limits: Age / Other (See Note above)
Zoster / Shingles (HZV/ZVL, RZV)	90736	Zoster (shingles) vaccine (HZV), live, for subcutaneous injection	Zostavax®	Adult	Benefit Limit: Age 60 years and up
	90750	Zoster (shingles) vaccine (HZV), recombinant, subunit, adjuvanted, for intramuscular use	Shingrix®	Adult	Benefit Limit: Age 50 years and up
Hepatitis B	90739	Hepatitis B vaccine (HepB), adult dosage, 2 dose schedule, for intramuscular use	HEPLISAV-B®	Adult	Benefit Limit: Age 18 and up
	90740	Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 3 dose schedule, for intramuscular use	Recombivax HB®	Both	-
	90743	Hepatitis B vaccine (HepB), adolescent , 2 dose schedule, for intramuscular use	Recombivax HB®	Pediatric (adolescent only)	For applicable age see code description.
	90744	Hepatitis B vaccine (HepB), pediatric/adolescent dosage , 3 dose schedule, for intramuscular use	Recombivax HB® Engerix-B®	Pediatric	For applicable age see code description.
	90746	Hepatitis B vaccine (HepB), adult dosage , 3 dose schedule, for intramuscular use	Recombivax HB® Engerix-B®	Adult	For applicable age see code description.
	90747	Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 4 dose schedule, for intramuscular use	Engerix-B®	Both	-
	90748	Hepatitis B and Haemophilus influenza b vaccine (Hib-HepB), for intramuscular use	N/A	Both	-

Expanded Women’s Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
Well-Woman Visits HRSA Requirement (Dec. 2016): Recommends that women receive at least one preventive care visit per year beginning in adolescence and continuing across the lifespan to ensure that the recommended preventive services including preconception, and many services necessary for prenatal and interconception care are obtained.	Procedure Code(s): <i>Well-Woman Visits:</i> See the Wellness Examinations row in the <i>Preventive Care Services</i> section.	<i>Well-Woman Visits:</i> See the Wellness Examinations row in the <i>Preventive Care Services</i> section.
	<i>Prenatal Office Visits:</i> Evaluation and Management (Office Visits): 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, G0463	<i>Prenatal Office Visits:</i> Requires a Pregnancy Diagnosis Code .

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
<p>The primary purpose of these visits should be the delivery and coordination of recommended preventive services as determined by age and risk factors.</p> <p>Also see Wellness Examinations and Preeclampsia Screening in the Preventive Care Services section.</p>	<p>Physician Prenatal Education, Group Setting: 99078</p> <p><i>Prenatal Care Visits:</i> 59425, 59426</p>	<p><i>Prenatal Care Visits:</i> Does not have diagnosis code requirements for the preventive benefit to apply.</p>
	<p><i>Global Obstetrical Codes:</i> 59400, 59510, 59610, 59618</p>	
	<p>Diagnosis Code(s): See the Pregnancy Diagnosis Codes.</p>	
<p>Screening for Gestational Diabetes Mellitus</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends screening pregnant women for gestational diabetes mellitus after 24 weeks of gestation (preferably between 24 and 28 weeks of gestation) in order to prevent adverse birth outcomes. Screening with a 50-g oral glucose challenge test (followed by a 3-hour 100-g oral glucose tolerance test if results on the initial oral glucose challenge test are abnormal) is preferred because of its high sensitivity and specificity. This recommendation also suggests that women with risk factors for diabetes mellitus be screened for preexisting diabetes before 24 weeks of gestation—ideally at the first prenatal visit, based on current clinical best practices.</p> <p>Also see the Diabetes Screening and Gestational Diabetes Mellitus Screening sections of the Preventive Care Services section, and the Screening for Diabetes Mellitus After Pregnancy section.</p>	<p>Procedure Code(s): <i>Diabetes Screening:</i> 82947, 82948, 82950, 82951, 82952, 83036</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): See the Pregnancy Diagnosis Codes.</p>	<p><i>Diabetes Screening:</i> Requires a Pregnancy Diagnosis Code (regardless of gestational week).</p> <p><i>Blood Draw:</i> Requires one of the diabetes screening procedure codes listed in this row AND one of the Pregnancy Diagnosis Codes.</p> <p>Note: If a diabetes diagnosis code is present in any position, the preventive benefit will not be applied. See the Diabetes Diagnosis Code List.</p>
<p>Screening for Diabetes Mellitus After Pregnancy</p> <p><u>HRSA Requirement (Dec. 2017)</u> The Women's Preventive Services</p>	<p>Procedure Code(s): <i>Diabetes Screening:</i> 82947, 82948, 82950, 82951, 82952, 83036</p>	<p><i>Diabetes Screening:</i> Requires one of the Required Screening diagnosis codes listed in this row AND Z86.32.</p>

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
<p>Initiative recommends women with a history of gestational diabetes mellitus (GDM) who are not currently pregnant and who have not previously been diagnosed with type 2 diabetes mellitus should be screened for diabetes mellitus. Initial testing should ideally occur within the first year postpartum and can be conducted as early as 4-6 weeks postpartum. Women with a negative initial postpartum screening test result should be rescreened at least every 3 years for a minimum of 10 years after pregnancy.</p> <p>Also see Gestational Diabetes Mellitus Screening and Diabetes Screening in the <i>Preventive Care Services</i> section, and the Screening for Gestational Diabetes Mellitus section.</p>	<p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Required Screening Diagnosis Codes (requires at least one): Z00.00, Z00.01, Z13.1 AND requires the following additional code: Additional Diagnosis Code Required: Z86.32 (personal history of gestational diabetes)</p>	<p>No age limit.</p> <p><i>Blood Draw:</i> Requires one of the Diabetes Screening procedure codes listed in this row AND one of the Required Screening diagnosis codes listed in this row AND Z86.32.</p> <p>Note: If a diabetes diagnosis code is present in any position, the preventive benefit will not be applied. See the Diabetes Diagnosis Code List.</p>
<p>Screening for Urinary Incontinence</p> <p>The Women's Preventive Services Initiative recommends screening women for urinary incontinence annually.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>
<p>Counseling for Sexually Transmitted Infections (STIs)</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends directed behavioral counseling by a health care provider or other appropriately trained individual for sexually active adolescent and adult women at an increased risk for STIs.</p> <p>This recommendation further recommends that health care providers use a woman's sexual history and risk factors to help identify those at an increased risk of STIs. Risk factors may include age younger than 25, a recent history of an STI, a new sex partner, multiple partners, a partner with concurrent partners, a partner with an STI, and a lack of or inconsistent condom use. For</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
adolescents and women not identified as high risk, counseling to reduce the risk of STIs should be considered, as determined by clinical judgement.		
<p>Education, Risk Assessment, and Screening for Human Immunodeficiency Virus Infection</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends prevention education and risk assessment for human immunodeficiency virus (HIV) infection in adolescents and women at least annually throughout the lifespan. All women should be tested for HIV at least once during their lifetime. Additional screening should be based on risk, and screening annually or more often may be appropriate for adolescents and women with an increased risk of HIV infection.</p> <p>Screening for HIV is recommended for all pregnant women upon initiation of prenatal care with retesting during pregnancy based on risk factors. Rapid HIV testing is recommended for pregnant women who present in active labor with an undocumented HIV status. Screening during pregnancy enables prevention of vertical transmission.</p>	<p>Education and Risk Assessment: See the Wellness Examinations row in the <i>Preventive Care Services</i> section above</p> <p>Screening Tests: See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the <i>Preventive Care Services</i> section above</p>	<p>Education and Risk Assessment: See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p> <p>Screening Tests: See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the <i>Preventive Care Services</i> section above.</p>
<p>Contraceptive Methods (Including Sterilizations)</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends that adolescent and adult women have access to the full range of female-controlled contraceptives to prevent unintended pregnancy and improve birth outcomes. Contraceptive care should include contraceptive counseling, initiation of contraceptive use, and follow-up care (e.g., management, and evaluation as well as changes to</p>	<p>Code Group 1 Procedure Code(s): <i>Sterilizations:</i> Tubal Ligation, Oviduct Occlusion: 58565, 58600, 58605, 58611, 58615, 58670, 58671, A4264 (See Code Group 4 below for Tubal Ligation Followup)</p> <p><i>Contraceptive Methods:</i> Diaphragm or Cervical Cap: 57170, A4261, A4266 IUD (copper): J7300 IUD (Skyla®): J7301 IUD (Liletta®): J7297</p>	<p>Code Group 1: Does not have diagnosis code requirements for preventive benefits to apply.</p>

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service:	Code(s):	Preventive Benefit Instructions:
<p><i>A date in this column reflects when the listed rating was issued.</i></p> <p>and removal or discontinuation of the contraceptive method). The Women's Preventive Services Initiative recommends that the full range of female-controlled U.S. Food and Drug Administration-approved contraceptive methods, effective family planning practices, and sterilization procedures be available as part of contraceptive care.</p> <p>Additionally, instruction in fertility awareness-based methods, including the lactation amenorrhea method, although less effective, should be provided for women desiring an alternative method.</p> <p>For counseling and followup care, see the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p> <p>Notes: Certain employers may qualify for an exemption from covering contraceptive methods and sterilizations on account of religious objections. Refer to the Outpatient Prescription Drug Rider, or SPD for self-funded plans, for specific prescription drug product coverage and exclusion terms, and myuhc.com for information regarding coverage for contraceptive drugs.</p>	<p>IUD (Kyleena[®]): J7296 (See Code Group 2 below for additional IUD codes)</p>	
	<p>Code Group 2 Procedure Code(s): <i>Contraceptive Methods:</i> Implantable Devices: J7306, J7307 11976 (capsule removal) 11981 (implant insertion) 11982 (implant removal) 11983 (removal with reinsertion)</p> <p>IUDs: J7298 (Mirena[®]) S4989 58300, S4981 (insertion) 58301 (removal) (See Code Group 1 above for additional IUD codes)</p> <p><i>Injections:</i> J1050 (injection) 96372 (administration)</p>	<p>Code Group 2: Requires one of the Code Group 2 diagnosis codes listed in this row.</p>
	<p>Code Group 2 Diagnosis Code(s): These are required for Code Group 2. <i>Contraceptive Management:</i> Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9</p>	
	<p>Code Group 3 Procedure Code(s): <i>Anesthesia for Sterilization:</i> 00851, 00940, 00942, 00950, 00952, 01960, 01961, 01965, 01966, 01967, 01968</p> <p>Code Group 3 Diagnosis Code(s): <i>Sterilization:</i> Z30.2</p>	<p>Code Group 3: Requires one of the Code Group 3 diagnosis code listed in this row.</p>
	<p>Code Group 4 Procedure Code(s): Tubal Ligation Followup Hysterosalpingogram: <i>Catheterization and Introduction of Saline or Contrast Material:</i> 58340 <i>Hysterosalpingography:</i> 74740 <i>Contrast Material:</i> Q9967</p> <p>Code Group 4 Diagnosis Code(s): <i>Tubal Ligation Status:</i> Z98.51</p>	<p>Code Group 4: Requires one of the Code Group 4 diagnosis code listed in this row.</p>

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
	<p>Code Group 5 Procedure Code(s): <i>IUD Followup Visit:</i> 99211, 99212</p> <p>Code Group 5 Diagnosis Code(s): Z30.431</p>	<p>Code Group 5: Requires one of the Code Group 5 diagnosis code listed in this row.</p>
<p>Breastfeeding Services and Supplies</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends comprehensive lactation support services (including counseling, education, and breastfeeding equipment and supplies) during the antenatal, perinatal, and postpartum periods to ensure the successful initiation and maintenance of breastfeeding.</p>	<p>Counseling and Education Procedure Code(s): 99241*, 99242*, 99243*, 99244*, 99245*, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, S9443</p> <p>Also see the codes in the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p> <p>Counseling and Education Diagnosis Code(s): Z39.1</p> <p>*For additional information on the reimbursement of consultation codes 99241-99245, refer to the Reimbursement Policy titled Consultation Services Policy.</p>	<p>Counseling and Education: Diagnosis code Z39.1 is required for 99241-99245, 99341-99345, and 99347-99350.</p> <p>Diagnosis code Z39.1 is not required for S9443.</p>
	<p>Breastfeeding Equipment & Supplies Procedure Code(s): <i>Personal Use Electric Breast Pump:</i> E0603</p> <p><i>Breast Pump Supplies:</i> A4281, A4282, A4283, A4284, A4285, A4286</p> <p>Breastfeeding Equipment & Supplies Diagnosis Code(s): Pregnancy Diagnosis Codes OR Z39.1.</p>	<p>Breastfeeding Equipment & Supplies: E0603 is limited to one purchase per birth.</p> <p>E0603 and A4281-A4286 require at least one of the diagnosis codes listed in this row.</p>
<p>Screening and Counseling for Interpersonal and Domestic Violence</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends screening adolescents and women for interpersonal and domestic violence, at least annually, and, when needed, providing or referring for initial intervention services. Interpersonal and domestic violence includes physical violence, sexual violence,</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
<p>stalking and psychological aggression (including coercion), reproductive coercion, neglect, and the threat of violence, abuse, or both. Intervention services include, but are not limited to, counseling, education, harm reduction strategies, and referral to appropriate supportive services.</p> <p>Also see the Screening for Intimate Partner Violence row in the <i>Preventive Care Services</i> section above.</p>		
<p>Breast Cancer Screening for Average-Risk Women</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends that average-risk women initiate mammography screening no earlier than age 40 and no later than age 50. Screening mammography should occur at least biennially and as frequently as annually. Screening should continue through at least age 74 and age alone should not be the basis to discontinue screening. These screening recommendations are for women at average risk of breast cancer. Women at increased risk should also undergo periodic mammography screening; however, recommendations for additional services are beyond the scope of this recommendation.</p>	<p>See the Screening Mammography row in the <i>Preventive Care Services</i> section above.</p>	<p>See the Screening Mammography row in the <i>Preventive Care Services</i> section above.</p>
<p>Screening for Cervical Cancer</p> <p><u>HRSA Requirement (Dec. 2016):</u> Recommends cervical cancer screening for average-risk women aged 21 to 65 years. For women aged 21 to 29 years recommends cervical cancer screening using cervical cytology (Pap test) every 3 years. Co-testing with cytology and human papillomavirus testing is not recommended for women younger than 30 years. Women aged 30 to 65 years should be screened with cytology and human papillomavirus testing every 5 years or cytology alone every 3</p>	<p>Human Papillomavirus DNA Testing (HPV)</p> <p>See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p>	<p>Human Papillomavirus DNA Testing (HPV)</p> <p>See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p>
	<p>Cervical Cytology (Pap Test):</p> <p>See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p>	<p>Cervical Cytology (Pap Test):</p> <p>See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

*These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.*

Service: <i>A date in this column reflects when the listed rating was issued.</i>	Code(s):	Preventive Benefit Instructions:
years. Women who are at average risk should not be screened more than once every 3 years.		

Revenue Codes

See the [Screening Mammography](#) and [Preventive Vaccines \(Immunizations\)](#) sections above for the applicable revenue codes.

Diagnosis Codes

[Preventive Care Services: ICD-10 Diagnosis Codes](#)

REFERENCES

ACIP Abbreviations for Vaccines (including Trade Names): <https://www.cdc.gov/vaccines/hcp/acip-recs/vac-abbrev.html>. Accessed July 12, 2019.

ACIP Vaccine Recommendations and Guidelines: <https://www.cdc.gov/vaccines/hcp/acip-recs/index.html>. Accessed July 12, 2019.

American Academy of Family Physicians (AAFP) Clinical Preventive Services Recommendations <https://www.aafp.org/patient-care/browse/type.tag-clinical-preventive-services-recommendations.html>. Accessed July 12, 2019.

American Academy of Pediatrics / Bright Futures / Recommendations for Pediatric Preventive Healthcare. (For ages 0–21): https://www.aap.org/en-us/Documents/periodicity_schedule.pdf. Accessed July 12, 2019.

American Academy of Pediatrics, Bright Futures Guidelines, 4th edition, Evidence and Rationale chapter https://brightfutures.aap.org/Bright%20Futures%20Documents/BF4_Evidence_Rationale.pdf. Accessed July 12, 2019.

American Academy of Pediatrics: <http://www.aap.org/>. Accessed July 12, 2019.

Centers for Disease Control and Prevention / Immunization Schedules: <http://www.cdc.gov/vaccines/schedules/index.html>. Accessed July 12, 2019.

Federal Register / Vol. 83, No. 39/Tuesday, February 27, 2018 / Notices: <https://www.gpo.gov/fdsys/pkg/FR-2018-02-27/pdf/2018-03943.pdf>. Accessed July 12, 2019.

Grade Definitions for USPSTF Recommendations <http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>. Accessed July 12, 2019.

July 19, 2010 IRS Interim Rules: http://www.irs.gov/irb/2010-29_IRB/index.html. Accessed July 12, 2019.

Published Recommendations, U.S. Preventive Services Task Force: <http://www.uspreventiveservicestaskforce.org/BrowseRec/Index/browse-recommendations>. Accessed July 12, 2019.

U.S. Food and Drug Administration (FDA), Vaccines Licensed for Use in the United States: <http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM093833>. Accessed July 12, 2019.

Women's Preventive Services Guidelines <https://www.hrsa.gov/womens-guidelines-2016/index.html>. Accessed July 12, 2019.

Women's Preventive Services Initiative (WPSI) <https://www.womenspreventivehealth.org/>. Accessed July 12, 2019.

GUIDELINE HISTORY/REVISION INFORMATION

Date	Action/Description
10/01/2019	<p>Frequently Asked Questions</p> <ul style="list-style-type: none"> • Updated <i>Answer #8</i> to clarify a new vaccine that is pending ACIP recommendations, but is a combination of previously approved individual components, <i>may be</i> eligible under the preventive care benefit <p>Applicable Codes</p> <ul style="list-style-type: none"> • Updated service description for HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults: <ul style="list-style-type: none"> ○ Removed April 2013 USPSTF 'A' rating ○ Added June 2019 USPSTF 'A' rating to indicate the USPSTF recommends that clinicians screen for HIV infection in: <ul style="list-style-type: none"> ▪ Adolescents and adults aged 15 to 65 years; younger adolescents and older adults who are at increased risk of infection should also be screened ▪ All pregnant persons, including those who present in labor or at delivery whose HIV status is unknown • Updated lists of applicable ICD-10 diagnosis codes to reflect annual code edits for: <ul style="list-style-type: none"> ○ Latent Tuberculosis Infection: Screening, Adults and Tuberculosis (TB) Testing (Bright Futures): Added Z11.7 ○ Atherosclerosis: Revised description for I70.238 and I70.248 • Updated benefit age limit for Gardasil9® [Human Papilloma Virus (HPV): CPT code 90651]; replaced "ages 9-26 years (ends on 27th birthday)" with "ages 9-45 years (ends on 46th birthday)" <p>Supporting Information</p> <ul style="list-style-type: none"> • Updated <i>References</i> section to reflect the most current information • Archived previous policy version CDG.016.26

INSTRUCTIONS FOR USE

This Coverage Determination Guideline provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this guideline, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Coverage Determination Guideline is provided for informational purposes. It does not constitute medical advice.

This Coverage Determination Guideline may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence ([Medicare IOM Pub. No. 100-16, Ch. 4, §90.5](#)).

UnitedHealthcare may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. UnitedHealthcare Coverage Determination Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

For self-funded plans with SPD language other than fully-insured Generic COC language, refer to the member specific benefit plan document for coverage.